


**NOTIFICATION PURSUANT TO
SECTION 6 OF DSHEA
AND 21 CFR § 161.93**



This notification is being filed on behalf of Basic Reseat Evanstown Research Labs that is the Distributor of the product(s) which bear the statements identified in this notification. Its business address is: 402 West 5050 North, Provo, Utah 84604. This notification is being made pursuant to Section 6 of DSHEA and Rule 21 C.F.R. § 101.93. The dietary supplement product on whose label or labeling the statements appear is **CarboLean™ – 90 ct. Box.**

The text of each structure-function statement for which notification is now being given is:

- (Statement 1): "Weight control"
- (Statement 2): "Intended to enhance: low carbohydrate weight loss programs"
- (Statement 3): "Adding CarboLean™ to your present low-carb diet plan can: Increase loss of body fat without changing what you eat."
- (Statement 4): "Adding CarboLean™ to your present low-carb diet plan can: Inhibit production of prostaglandins that impede fat loss at the cellular level"
- (Statement 5): "Adding CarboLean™ to your present low-carb diet plan can: Increase loss of body fat without altering your present caloric intake or low-carb diet regimen"
- (Statement 6): "Adding CarboLean™ to your present low-carb diet plan can: Dramatically elevate your energy levels"
- (Statement 7): "Adding CarboLean™ to your present low-carb diet plan can: Eliminate low-carb diet fatigue"

The following summary identifies the dietary ingredient(s) or supplement(s) for which a statement has been made:

<u>Statement Number</u>	<u>Identity of Dietary Ingredient(s) or Supplement(s) that is the Subject of the Statement:</u>
1	CarboLean™ – 90 ct. Box
2	CarboLean™ – 90 ct. Box
3	CarboLean™ – 90 ct. Box
4	CarboLean™ – 90 ct. Box
5	CarboLean™ – 90 ct. Box
6	CarboLean™ – 90 ct. Box
7	CarboLean™ – 90 ct. Box

97S 0162

LET 7475


74961

The following identifies the brand name of each supplement for which a statement is made:

<u>Statement Number</u>	<u>Brand Name</u>	<u>Label or Labeling?</u>
1	Basic Research, L.L.C, dba Evanstown Research Labs	both
2	Basic Research, L.L.C, dba Evanstown Research Labs	both
3	Basic Research, L.L.C, dba Evanstown Research Labs	both
4	Basic Research, L.L.C, dba Evanstown Research Labs	both
5	Basic Research, L.L.C, dba Evanstown Research Labs	both
6	Basic Research, L.L.C, dba Evanstown Research Labs	both
7	Basic Research, L.L.C, dba Evanstown Research Labs	both

I, Daniel B. Mowrey, Ph.D., am authorized to certify this Notification on behalf of Basic Research, L.L.C. dba Evanstown Research Labs. I certify that the information presented and contained in this Notification is complete and accurate, and that Basic Research, L.L.C. dba Evanstown Research Labs has substantiation that each structure-function statement is truthful and not misleading.

Dated this 4 day of March, 2001.

By: 
Daniel B. Mowrey, Ph.D.,
Consultant to Basic Research, L.L.C.
dba Evanstown Research Labs

Having a Hard Time on Your Low Carb Diet?

Adding CarboLean™ to your present low-carb diet plan can:

- Increase loss of body fat without changing what you eat.*
- Inhibit production of prostaglandins that impede fat loss at the cellular level.**
- Increase loss of body fat without altering your present caloric intake or low-carb diet regimen.**
- Dramatically elevate your energy levels.*
- Eliminate low-carb diet fatigue.**

WARNING: Maximum recommended total amount of ephedrine for a healthy adult human no more than 100mg in a 24 hour period for not more than 12 weeks.

WARNING: Do not take this product with any other product containing Ephedrine, Ephedra, or Ma Huang.

If you experience adverse effects after consumption of this product and you live in Texas, you can call the Texas Department of Health. If you live outside of Texas, call your local Poison Control Center; or, if an emergency call 911.

For any other product information, call customer service at 1-800-898-5153.

WEIGHT CONTROL*

Carbo Lean™

Metabolic Adjunct

*In tended to Enhance;
Low Carbohydra te
Weight Loss Programs**



©2000 Evanstow
Printed on Recycled Paperboard
(Minimum of 35% Post Consumer Content)

DIETARY
SUPPLEMENT
90 CAPSULES

Active Ingredients: Ephedrine, Caffeine, Acetylsalicylic Acid

Supplement Facts

Serving Size 2 capsules
Servings Per Container 45

	Amount Per Serving	% Daily Value*
Ephedrine alkaloids (from standardized whole plant ma huang extract (250 mg))	20mg	†
Caffeine (26mg from standardized bitter melon (212 mg))	200mg	†
Acetylsalicylic acid	324mg	†
Proprietary Blend	241mg	
Glycemic Complex		
Schizonepeta (spical) Extract (5:1)		†
Forsythia (fruit) Extract (5:1)		†
Gymnema (leaves, stem)		†
Bitter Melon (seeds)		†
Stevia (leaf)		†
Chromium polynicotinate		†

* Percent Daily Values are based on a 2,000 calorie diet
† Daily Value not established

Other Ingredients: rice flour

This product has 20 mg. of ephedrine group alkaloids per serving in the form of herbal extracts may cause serious adverse health effects. Read the label and follow directions.

*No statement on this package has been reviewed by the Food and Drug Administration. Neither the specified products nor any of these writings are intended to diagnose, treat, cure, or prevent any disease.

Distributed by

Evanstow
RESEARCH LABS

0500

Provo, UT 84604 • 1-800-898-5153 • www.carbolean.com
Manufactured in the U.S.A.

Carbo Lean™ Directions: Take two capsules with a full glass of water, three times per day: morning, noon and early evening. Do not exceed 6 capsules per day. NOTE: Jitters, anxiety, dry mouth or insomnia may occur during the first week of use. Reducing number of capsules per day during the first week should eliminate these mild effects. For additional information on this or any other Evanstow tabs product, please call our experienced customer service representatives at 1-800-898-5153.

WARNINGS: Do not take if sensitive to or if allergic to aspirin. Do not use if pregnant (it is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a physician because it may cause problems in the unborn child or complications during pregnancy), nursing, have asthma, or have stomach problems (such as heartburn, upset stomach or stomach pain) that persist or recur, or if you have ulcers or bleeding problems, unless directed by a physician. If ringing in the ears or loss of hearing occurs, consult a physician before taking again. **DRUG INTERACTION PRECAUTION:** Do not take this product if you are taking MAO inhibitors or a prescription drug for anticoagulation (thinning of the blood), diabetes, gout or arthritis or any other prescription drug unless directed by a physician. Do not take this product if you have any of the following conditions: high blood pressure, cardiovascular disease, thyroid disease; prostate problems or diabetes. Improper use may be hazardous to your health. This product is not intended for pain relief, fever, toothache or any other analgesic effect. Keep out of the reach of children. Children and teenagers should not use this product for chicken pox or flu symptoms before a physician is consulted about Reye Syndrome, a rare but serious illness reported to be associated with aspirin. In case of overdose, get medical help or contact a Poison Control Center right away. Consult a physician or licensed health care professional before using this product if you have, or have a family history of, heart disease; thyroid disease, diabetes, high blood pressure, recurrent headaches, depression or other psychiatric condition, glaucoma, difficulty in urinating, prostate enlargement, or seizure disorder, if you are using any other dietary supplement, or over-the-counter drug containing ephedrine, pseudoephedrine, or phenylpropanolamine (ingredients found in certain allergy, asthma, cough and weight control products). Exceeding recommended serving may cause serious adverse health effects including heart attacks and stroke. Discontinue use and call physician or licensed health care professional immediately if you experience rapid heartbeat, dizziness, severe headache, shortness of breath or similar symptoms. Consuming additional caffeine with this product may cause serious adverse health effects.

ALCOHOL WARNING: If you consume 3 or more alcoholic drinks everyday, ask your physician whether you should take aspirin. Aspirin may cause stomach bleeding.

Do not take if under 18 years of age.