

LACHMAN CONSULTANT SERVICES, INC.
Westbury, NY 11590



ATTACHMENT D



OXYCODONE AND ACETAMINOPHEN TABLETS, USP

R_x Only

Code 000000

Rev. 03/2002

DESCRIPTION


Each tablet, for oral administration, contains oxycodone hydrochloride and acetaminophen in the following strengths:

Oxycodone hydrochloride*	5 mg	7.5 mg	10 mg
WARNING: May be habit forming.			
Acetaminophen, USP	300 mg	300 mg	300 mg

- * 5 mg oxycodone HCl is equivalent to 4.4815 mg of oxycodone.
- 7.5 mg oxycodone HCl is equivalent to 6.7228 mg of oxycodone.
- 10 mg oxycodone HCl is equivalent to 8.9637 mg of oxycodone.

All strengths of Oxycodone and Acetaminophen Tablets, USP, also contain the following inactive ingredients: To be determined.


Acetaminophen, 4'-hydroxyacetanilide, a slightly bitter, white, odorless, crystalline powder, is a non-opiate, non-salicylate analgesic and antipyretic. It has the following structural formula:



C₈H₉NO₂

MW = 151.16

The oxycodone component is 14-hydroxydihydrocodeinone, a white, odorless, crystalline powder having a saline, bitter taste. It is derived from the opium alkaloid thebaine, and may be represented by the following structural formula:



C₁₈H₂₁NO₄

MW = 315.37

CLINICAL PHARMACOLOGY

The principal ingredient, oxycodone, is a semisynthetic narcotic analgesic with multiple actions qualitatively similar to those of morphine; the most prominent of these involve the central nervous system and organs composed of smooth muscle. The principle actions of therapeutic value of the oxycodone in oxycodone and acetaminophen tablets are analgesia and sedation.

Oxycodone is similar to codeine and methadone in that it retains at least one-half of its analgesic activity when administered orally.

Acetaminophen is a non-opiate, non-salicylate analgesic and antipyretic.

INDICATIONS AND USAGE

Oxycodone and acetaminophen tablets are indicated for the relief of moderate to moderately severe pain.

CONTRAINDICATIONS

Oxycodone and acetaminophen tablets should not be administered to patients who are hypersensitive to oxycodone or acetaminophen.

WARNINGS

Drug Dependence: Oxycodone can produce drug dependence of the morphine type and, therefore, has the potential for being abused. Psychic dependence, physical dependence and tolerance may develop upon repeated administration of oxycodone and acetaminophen tablets, and it should be prescribed and administered with the same degree of caution appropriate to the use of other oral narcotic-containing medications. Like other narcotic-containing medications, oxycodone and acetaminophen tablets are subject to the Federal Controlled Substances Act (Schedule II).

PRECAUTIONS

General

Head Injury and Increased Intracranial Pressure: The respiratory depressant effects of narcotics and their capacity to elevate cerebrospinal fluid pressure may be markedly exaggerated in the presence of head injury, other intracranial lesions or a pre-existing increase in intracranial pressure. Furthermore, narcotics produce adverse reactions which may obscure the clinical course of patients with head injuries.

Acute Abdominal Conditions: The administration of oxycodone and acetaminophen tablets or other narcotics may obscure the diagnosis or clinical course in patients with acute abdominal conditions.

Special Risk Patients: Oxycodone and acetaminophen tablets should be given with caution to certain patients such as the elderly or debilitated, and those with severe impairment of hepatic or renal function, hypothyroidism, Addison's disease, and prostatic hypertrophy or urethral stricture.

Information for Patients

Oxycodone may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery. The patient using oxycodone and acetaminophen tablets should be cautioned accordingly.

Drug Interactions

Patient receiving other narcotic analgesics, general anesthetics, phenothiazines, other tranquilizers, sedative-hypnotics or other CNS depressants (including alcohol) concomitantly with oxycodone and acetaminophen tablets may exhibit an additive CNS depression. When such combined therapy is contemplated, the dose of one or both agents should be reduced.

The use of MAO inhibitors or tricyclic antidepressants with oxycodone preparations may increase the effect of either the antidepressant or oxycodone.

The concurrent use of anticholinergics with narcotics may produce paralytic ileus.

Usage in Pregnancy

Pregnancy Category C: Animal reproductive studies have not been conducted with oxycodone and acetaminophen tablets. It is also not known whether oxycodone and acetaminophen tablets can cause fetal harm when administered to a pregnant woman or can affect reproductive capacity. Oxycodone and acetaminophen tablets should not be given to a pregnant woman unless in the judgment of the physician, the potential benefits outweigh the possible hazards.

Nonteratogenic Effects: Use of narcotics during pregnancy may produce physical dependence in the neonate.

Labor and Delivery: As with all narcotics, administration of oxycodone and acetaminophen tablets to the mother shortly before delivery may result in some degree of respiratory depression in the newborn and the mother, especially if higher doses are used.

Nursing Mothers

It is not known whether oxycodone and acetaminophen are excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when oxycodone and acetaminophen tablets are administered to a nursing woman.

Pediatric Use

Safety and effectiveness in children have not been established.

ADVERSE REACTIONS

The most frequently observed adverse reactions include lightheadedness, dizziness, sedation, nausea and vomiting. These effects seem to be more prominent in ambulatory than in nonambulatory patients, and some of these adverse reactions may be alleviated if the patient lies down.

Other adverse reactions include euphoria, dysphoria, constipation, skin rash and pruritus. At higher doses, oxycodone has most of the disadvantages of morphine including respiratory depression.

DRUG ABUSE AND DEPENDENCE

Oxycodone and acetaminophen tablets are a Schedule II controlled substance.

Oxycodone can produce drug dependence and has the potential for being abused (See WARNINGS).

OVERDOSAGE

Acetaminophen

Signs and Symptoms: In acute acetaminophen overdosage, dose-dependent, potentially fatal hepatic necrosis is the most serious adverse effect. Renal tubular necrosis, hypoglycemic coma and thrombocytopenia may also occur.

In adults, hepatic toxicity has rarely been reported with acute overdoses of less than 10 grams and fatalities with less than 15 grams. Importantly, young children seem to be more resistant than adults to the hepatotoxic effect of an acetaminophen overdose. Despite this, the measures outlined below should be initiated in any adult or child suspected of having ingested an acetaminophen overdose.

Early symptoms following a potentially hepatotoxic overdose may include: nausea, vomiting, diaphoresis and general malaise. Clinical and laboratory evidence of hepatic toxicity may not be apparent until 48 to 72 hours post-ingestion.

Treatment: The stomach should be emptied promptly by lavage or by induction of emesis with syrup of ipecac. Patient's estimates of the quantity of a drug ingested are notoriously unreliable. Therefore, if an acetaminophen overdose is suspected, a serum acetaminophen assay should be obtained as early as possible, but no sooner than four hours following ingestion. Liver function studies should be obtained initially and repeated at 24-hour intervals.

The antidote, N-acetylcysteine, should be administered as early as possible, preferably within 16 hours of the overdose ingestion for optimal results, but in any case, within 24 hours. Following recovery, there are no residual, structural, or functional hepatic abnormalities.

Oxycodone

Signs and Symptoms: Serious overdosage with oxycodone is characterized by respiratory depression (a decrease in respiratory rate and/or tidal volume, Cheyne-Stokes respiration, cyanosis), extreme somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, and sometimes bradycardia and hypotension. In severe overdosage, apnea, circulatory collapse, cardiac arrest and death may occur.

Treatment: Primary attention should be given to the reestablishment of adequate respiratory exchange through provision of a patent airway and the institution of assisted or controlled ventilation. The narcotic antagonist naloxone hydrochloride is a specific antidote against respiratory depression which may result from overdosage or unusual sensitivity to narcotics, including oxycodone. Therefore, an appropriate dose of naloxone hydrochloride (usual initial adult dose 0.4 mg to 2 mg) should be administered preferably by the intravenous route, and simultaneously with efforts at respiratory resuscitation. Since the duration of action of oxycodone may exceed that of the antagonist, the patient should be kept under continued surveillance and repeated doses of the antagonist should be administered as needed to maintain adequate respiration.

An antagonist should not be administered in the absence of clinically significant respiratory or cardiovascular depression. Oxygen, intravenous fluids, vasopressors and other supportive measures should be employed as indicated.

Gastric emptying may be useful in removing unabsorbed drug.

DOSAGE AND ADMINISTRATION

Dosage should be adjusted according to the severity of the pain and the response of the patient. It may occasionally be necessary to exceed the usual dosage recommended below in cases of more severe pain or in those patients who have become tolerant to the analgesic effect of narcotics. Oxycodone and acetaminophen tablets are given orally. The usual adult dosage is one tablet every 6 hours as needed for pain. The total daily dose of acetaminophen should not exceed 4 grams.

HOW SUPPLIED

Oxycodone and Acetaminophen Tablets, USP are supplied as:

Oxycodone and Acetaminophen Tablets, USP 5 mg/300 mg; NDC# 00000-000-00
Oxycodone and Acetaminophen Tablets, USP 7.5 mg/300 mg; NDC# 00000-000-00
Oxycodone and Acetaminophen Tablets, USP 10 mg/300 mg; NDC# 00000-000-00

Tablet description and packaging information are to be determined.

STORAGE: Store at controlled room temperature 15° to 30°C (59° to 86°F) [See USP].

PHARMACIST: Dispense in a tight, light-resistant container with a child-resistant closure.

DEA Order Form Required.

Manufactured by:
Manufacturer

Code 000000

Rev. 03/2002

62
100

FedEx USA Airbill
Express

FedEx Tracking Number

8322 6920 8642

1 From This portion can be removed for Recipient's records.
 Date 3/12/02 FedEx Tracking Number 832269208642
 Sender's Name R. POLLUCK Phone 516 222-6222
 Company LACHMAN CONSULTANT SERVICE INC
 Address 1600 STEWART AVE STE 604 Dept./Floor/Suite/Room
 City WESTBURY State NY ZIP 11590

2 Your Internal Billing Reference

3 To Recipient's Name DOCKETS MANAGEMENT CONSULTANTS Phone
 Company FOOD & DRUG ADMINISTRATION
 Address 5133 FISHERS LANE We cannot deliver to P.O. boxes or P.O. ZIP codes.
 Address ROOM 1061 Dept./Floor/Suite/Room
 City ROCKVILLE State MD ZIP 20852



0207798553

FedEx
emp# 313236 12MAR02


TRK# 8322 6920 8642 FORM 0215

STANDARD OVERNIGHT WED

Deliver By:
13 MAR 02
A2

20852 -MD-US

IAD
ZM GAIA



5 Packaging
 FedEx Envelope*
 FedEx Pak* Includes FedEx Small Pak, FedEx Large Pak
 Other

6 Special Handling


SATURDAY Delivery
Available only for FedEx Priority Overnight and FedEx 2Day to select ZIP codes

Does this ship
 No Yes

Dangerous Goods

7 Payment

Sender's Acct. No. I will bill



Total Paid


*Our liability is limited

8 Release Signature

By signing you authorize and agree to indemnify us.
Questions? Visit us
 or call 1.800.Go.FedEx® 800.463.3333
SRS • Rev. Date 10/01 • Part # 576105 • ©1994-2001 FedEx

THIS PACKAGE
 HAS BEEN
 X-RAYED
 FDA

447



62

100

FedEx USA Airbill
Express

FedEx Tracking Number

8322 6920 8642

1 From This portion can be removed for Recipient's records.

Date 3/12/02 FedEx Tracking Number 832269208642

Sender's Name R. POLLOCK Phone 516 222-6222

Company LACHMAN CONSULTANT SERVICE INC

Address 1600 STEWART AVE STE 604 Dept./Floor/Suite/Room

City WESTBURY State NY ZIP 11590

2 Your Internal Billing Reference

3 To Recipient's Name DOCKETS MANAGEMENT BRANCH Phone

Company FOOD + DRUG ADMINISTRATION

Address 5630 FISHERS LANE We cannot deliver to F.O. boxes or P.O. ZIP codes.

Address ROOM 1061


City ROCKVILLE State MD ZIP 20852



FedEx STANDARD OVERNIGHT WED
 emp# 313236 12MAR02
 TRK# 8322 6920 8642 FORM 0215
 Deliver By: 13MAR02 A2

20852 -MD-US

IAD
ZM GAIA



RECIPIENT: PEEL HERE

FedEx 1Day Freight* Next business day
 FedEx 2Day Freight Second business day
 FedEx 3Day Freight Third business day

* Call for Confirmation: Delivery commitment may be later in some areas.

5 Packaging
 FedEx Envelope* *Declared value limit \$500


6 Special Handling
 SATURDAY Available only Overnight and to select ZIP
 No

7 Payment
 Sender's Acct. No. in 1 will be billed

Total Packages _____ Charges _____
 Credit Card Auth. _____

8 Release Signature Sign to authorize _____

By signing you authorize us to deliver this shipment without obtaining a signature and agree to indemnify and hold us harmless from any resulting claims.
Questions? Visit our Web site at fedax.com
 or call 1.800.Go.FedEx® 800.463.3339.
 SHS • Rev. Date 10/01 • Part #1576105 • ©1994-2001 FedEx • PRINTED IN U.S.A.



447

0207798553

Recycled