

QUALIFIED HEALTH CLAIM PETITION

**100% WHEY PROTEIN PARTIALLY HYDROLYZED
in Infant Formula and
REDUCING THE RISK OF ALLERGY IN INFANTS**

TABLE OF CONTENTS & BIBLIOGRAPHY

<u>VOLUME I</u>	<u>TAB</u>
EXECUTIVE SUMMARY	A
PRELIMINARY REQUIREMENTS FOR HEALTH CLAIM PETITIONS <i>100% Whey Protein Partially Hydrolyzed in infant formula meets all the preliminary requirements of 21 CFR 101.70</i>	B
SUMMARY OF SCIENTIFIC DATA <i>Definition of terms The substance/disease relationship Review and rating of the scientific evidence Certification of ethical compliance Review of selected clinical trials Summary of meta-analyses, review papers and non-peer-reviewed reports</i>	C
ANALYTICAL DATA – SUBSTANCE CHARACTERIZATION <i>Technical attributes of the substance & of products eligible to make the proposed claim</i>	D
MODEL QUALIFIED HEALTH CLAIM <i>Proposed message Consumer research Expert opinions</i>	E
ENVIRONMENTAL IMPACT ASSESSMENT	F
CERTIFICATION OF COMPLETENESS	G
REFERENCE LIST – VOLUME I: ALL SECTIONS	H
<u>VOLUMES II and III</u>	
COPIES OF LISTED REFERENCES – VOLUME I	I
<u>VOLUMES IV and V</u>	
COMPREHENSIVE LITERATURE SEARCH – ANNOTATED	J
COPIES OF ADDITIONAL ARTICLES NOT REFERENCED IN VOLUME I	K

**BIBLIOGRAPHY OF CLINICAL TRIALS
RELIED UPON IN SUPPORT OF CLAIM**

STUDY DESIGN - TYPE 1

Barberi I, Salpietro DC, Catalioto G, Fulia F. Clinical trial of a new hypoallergenic formula in risk infants. 2nd World Congress of Perinatal Medicine, Rome and Florence 1993 (September).

Becker A, Watson W, Ferguson A, Dimich-Ward H, Chan-Yeung M. The Canadian asthma primary prevention study: outcomes at 2 years of age. *J Allergy Clin Immunol* 2004;113:650-6.

Chan YH, Shek LPC, Aw M, Quak SH, Lee BW. Use of hypoallergenic formula in the prevention of atopic disease among Asian children. *J Paediatr Child Health* 2002;38:84-88.

Chan-Yeung M, Manfreda J, Dimich-Ward H, Ferguson A, Watson W, Becker A. A randomized controlled study on the effectiveness of a multifaceted intervention program in the primary prevention of asthma in high-risk infants. *Arch Pediatr Adolesc Med* 2000;154:657-63.

Chandra RK, Singh G, Shridhara B. Effect of feeding whey hydrolysate, soy and conventional cow milk formulas on incidence of atopic disease in high risk infants. *Ann Allergy* 1989;63:102-6.

Chandra RK, Hamed A. Cumulative incidence of atopic disorders in high risk infants fed whey hydrolysate, soy, and conventional cow milk formulas. *Ann Allergy* 1991;67:129-32.

Chandra RK, Hamed A, Prasad C, Singh GK. Cumulative incidence of allergic disorders in high risk infants fed whey hydrolysate, soy and conventional cow's milk formulas. *Am J Clin Nutr* 1992;56:758 (Abstract).

Chandra, RK. Five-year follow-up of high-risk infants with family history of allergy who were exclusively breast-fed or fed partial whey hydrolysate, soy, and conventional cow's milk formulas. *J Pediatr Gastroenterol Nutr* 1997;24:380-8.

De Seta L, Siani P, Cirillo G, Di Gruttola M, Cimaduomo L, Coletta S. Prevention of allergic disease by an hypoallergenic formula: preliminary results at the 24 months follow-up. *Med Surg Ped* 1994;16:251-4.

Lam BCC, Yeung CY. The effect of breast milk, infant formula and hypoallergenic formula on incidence of atopic manifestation in high risk infants. 6th Annual Scientific Meeting of Hong Kong College of Physicians and Hong Kong College of Paediatricians, Hong Kong 1992 (October).

Lam BCC, Yeung CY. The effect of breast milk, infant formula and hypoallergenic formula on incidence of atopic manifestation. Internal report, Nestec SA 1992.

Iikura Y, Akimoto K, Ebisawa M, Onda T, Akazawa A, Saito H, Kimura T, Ishizawa K, Koya N. Effect of hydrolyzed whey protein formula for babies on development of allergic symptoms during infancy. Nestlé Nutrition Workshop Intestinal Immunology and Food Allergy, Raven Press 1995;34:269-75.

Marini A, Agosti M, Motta G, Mosca F. Effects of a dietary and environmental prevention programme on the incidence of allergic symptoms in high atopic risk infants: three years follow-up. *Acta Paediatr Suppl* 1996;414:1-21.

Martinez Valverde A. Study on the assessment of serum IgE and specific IgE to Alpha lactalbumin, Beta lactoglobulin, and casein in infants fed hypoallergenic formula and other forms of feeding. Thesis, Universidad de Malaga, Spain, 1993.

Silva Rey AL, Garcia G, Nogales A. Preventive effect of a partially hydrolysed cow milk infant formula: clinical and biological 24 month follow-up study. Thesis, Spain, 1996.

Tsai Y, Chou C, Hsieh K. The effect of hypoallergenic formula on the occurrence of allergic diseases in high risk infants. *Acta Paed Sin* 1991;32:137-44.

Vandenplas Y, Hauser B, Van den Borre C, Sacre L, Dab I. Effect of a whey hydrolysate prophylaxis of atopic disease. *Ann Allergy* 1992;68:419-24.

Vandenplas Y. Atopy at 3 years in high-risk infants fed whey hydrolysate or conventional formula. *Lancet* 1992;339:1118. Letter to the Editor.

Vandenplas Y, Hauser B, Van den Borre C, Clybouw C, Mahler T, Hachimi-Idrissi S, Deraeve L, Malfroot A, Dab I. The long-term effect of a partial whey hydrolysate formula on the prophylaxis of atopic disease. *Eur J Pediatr* 1995;154:488-94.

Von Berg A, Koletzko S, Grübl A, Filipiak-Pittroff B, Wichmann H, Bauer CP, Reinhardt D, Berdel D, German Infant Nutritional-Intervention Study Group. The effect of hydrolyzed cow's milk formula for allergy prevention in the first year of life: The German Infant Nutritional Intervention Study, a randomized double-blind trial. *J Allergy Clin Immunol* 2003;111:533-40.

STUDY DESIGN TYPE 3

D'Agata A, Betta P, Sciacca P, Morano C, Practico G, Curreri R, Quattrocchi O, Sciacca F. The role of preventing diet in newborn at risk of atopy. Follow-up results. *Med Surg Ped* 1996;18:469-72.

Exl BM, Deland U, Wall M, Preysch U, Secretin MC, Shmerling DH. Zug-Frauenfeld nutritional survey (“Zuff Study”): allergen-reduced nutrition in a normal infant population and its health –related effects: results at the age of six months. *Nutr Res* 1998;18:1443-62.

Exl BM, Deland U, Secretin MC, Preysch U, Wall M, Shmerling DH. Improved general health status in an unselected infant population following an allergen-reduced dietary intervention programme: the ZUFF-STUDY-PROGRAMME. Part II: infant growth and health status to age 6 months. *Eur J Nutr* 2000;39:145-56.

Schmidt E, Eden-Köhler J, Tonkaboni F, Tölle J. Alimentary allergy prevention in infants with familial increased allergic risk: The effect of different feeding regimens in the first 6 months on atopic manifestations during the first year of life. A large-scale feeding trial. Nestlé Nutrition Workshop Intestinal Immunology and Food Allergy, Raven Press 1995;34:231-48.

Vandenplas Y, Deneyer M, Sacre L, Loeb H. Preliminary data on a field study with a new hypo-allergenic formula. *Eur J Pediatr* 1988;148:274-7.

Vandenplas Y, Malfroot A, Dab I. Short-term prevention of cow’s milk protein allergy in infants. *Immunology & Allergy Practice* 1989;11:430-7.

Willems R, Duchateau J, Magrez P, Denis R, Casimir G. Influence of hypoallergenic milk formula on the incidence of early allergic manifestations in infants predisposed to atopic diseases. *Ann Allergy* 1993;71:147-50.

REVIEW ARTICLES AND META ANALYSES

Baumgartner M, Brown CA, Exl BM, Secretin MC, van’t Hof M, Haschke F. Controlled trials investigating the use of one partially hydrolyzed whey formula for dietary prevention of atopic manifestations until 60 months of age: an overview using meta-analytical techniques. *Nutr Res* 1998;18:1425-42.

Chandra RK. Breast feeding, hydrolysate formulas and delayed introduction of selected foods in the prevention of food hypersensitivity and allergic disease. *Nutr Res* 2002;21:125-35.

Halken S, Jacobsen HP, Host A, Holmenlund D. The effect of hypo-allergenic formulas in infants at risk of allergic disease. *Eur J Clin Nutr* 1995;49:S77-83.

Osborn DA, Sinn J. Formulas containing hydrolysed protein for prevention of allergy and food intolerance in infants (Cochrane Review). In: *The Cochrane Library*, Issue 4, 2003. Chichester, UK: John Wiley & Sons, Ltd.

Ram FSF, Ducharme FM, Scarlett J. Cow’s milk protein avoidance and development of childhood wheeze in children with a family history of atopy (Cochrane Review) In: *The Cochrane Library*, Issue 2, 2003. Oxford: Update Software.

Schoetzau A, Gehring U, Wichmann HE. Prospective cohort studies using hydrolysed formulas for allergy prevention in atopy-prone newborns: a systematic review. *Eur J Pediatr* 2001;160:323-32.

Zeiger RS. Dietary aspects of food allergy prevention in infants and children. *J Pediatr Gastroenterol Nutr* 2000;39:S77-86.