

Drug Facts
 Active ingredient (in each gelcap)
 Famotidine 10mg
 Purpose
 Acid reducer

Uses
 • relieves heartburn associated with acid indigestion and sour stomach
 • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
 Allergy alert: Do not use if you are allergic to famotidine or other acid reducers
 Do not use
 • if you have trouble swallowing
 • with other acid reducers
 Stop use and ask a doctor if
 • stomach pain continues
 • you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 • adults and children 12 years and over:
 • To relieve symptoms, swallow 1 gelcap with a glass of water from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
 • To prevent symptoms, swallow 1 gelcap with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
 • do not use more than 2 gelcaps in 24 hours
 • children under 12 years: ask a doctor



Pepcid AC Gelcaps

Do not use if the individual blister unit is open or torn.

- 1 gelcap relieves heartburn due to acid indigestion (Head Package insert before use).
- Pepcid AC prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Pepcid AC Gelcaps **FREE Pepcid AC Offer**
 (Mail-in offer. See details inside.)

0000000

Distributed by:
Johnson & Johnson • MERCK
 CONSUMER PHARMACEUTICALS CO.
 FORT WASHINGTON, PA 19034 USA

If You Have Questions Or Comments
 Please Call 1-800-755-4008

Registered trademark of Merck & Co., Inc.

FOR POSITION ONLY

000000 000000

NEW! GELCAPS

Pepcid AC

Famotidine Tablets 10mg/Acid Reducer®

Gelcaps

Relieves & Prevents Heartburn Due To Acid Indigestion

30 Gelcaps (Gelatin Coated, Capsule Shaped Tablets) Just One Per Dose

LOT AND EXP. DATE
 STAMPED HERE

Pepcid AC Gelcaps

Drug Facts (continued)

Other information
 • read the directions and warnings before use
 • keep the carton and package insert. They contain important information.
 • store at 25-30°C (77-86°F) • protect from moisture

Inactive ingredients benzyl alcohol, black iron oxide, butylparaben, castor oil, edetate calcium disodium, FD&C red #40, gelatin, hydroxypropyl methylcellulose, magnesium stearate, methylparaben, microcrystalline cellulose, pregelatinized corn starch, propylene glycol, propylparaben, sodium lauryl sulfate, sodium propionate, talc, titanium dioxide

Questions or comments? 1-800-755-4008



Do not use if the individual blister unit is open or torn.

WHAT YOU SHOULD KNOW ABOUT:



PACKAGE INSERT

A non-prescription stomach medicine

PEPCID AC Gelcaps contain a medicine, different from antacids, that doctors have prescribed for years to treat acid-related problems in millions of people.


- Product Benefits:**
- 1 gelcap relieves heartburn due to acid indigestion.
 - Pepcid AC prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.
 - It contains famotidine, a prescription-proven medicine.



Shown actual size

How to use PEPCID AC Gelcaps


Use PEPCID AC Gelcaps to relieve or prevent heartburn due to acid indigestion and sour stomach.



TO RELIEVE SYMPTOMS

Swallow 1 gelcap with a glass of water (do not chew).

PEPCID AC can be used up to twice daily (up to 2 gelcaps in 24 hours).



TO PREVENT SYMPTOMS

Swallow 1 gelcap with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn.

Excess acid: a burning problem

It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the wrong place (the esophagus), or too much acid, can cause burning pain and discomfort that interfere with everyday activities.

Heartburn— Caused by acid in the esophagus

A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position



Burning pain/ discomfort
Excess acid moves up into esophagus

Tips for Managing Heartburn


- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

Proven effective in clinical studies

In clinical studies, PEPCID AC film-coated tablets were significantly better in relieving and preventing heartburn than placebo tablets (tablets without the medicine). PEPCID AC Gelcaps contain the same active ingredient.

Percent of heartburn episodes completely relieved

Percent of patients with prevention or reduction of heartburn symptoms



*Time taken before eating a meal that is expected to cause symptoms.

Excellent safety record

The ingredient in PEPCID AC, famotidine, has been prescribed by doctors for years to treat millions of patients safely and effectively. The active ingredient in PEPCID AC has been taken safely with many frequently prescribed medications.

Know when to see your doctor

Heartburn and acid indigestion are common, but you should stop taking PEPCID AC and see your doctor promptly if:

- You have trouble swallowing or stomach pain continues. You may have a serious condition that may need different treatment.
- You need to take this product for more than 14 days.

Important: Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use with other acid reducers. This product should not be given to children under 12 years old, unless directed by a doctor. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a poison control center immediately.

Questions or comments?




Please call 1-800-755-4008 for more information and a guide of how to help your heartburn.

© Registered trademark of Merck & Co., Inc.
Distributed by:
Johnson & Johnson • MERCK
CONSUMER PHARMACEUTICALS CO.
FORT WASHINGTON, PA 19034 USA

00000000

COUPON





Pepcid
AC
Gelcaps

FREE
Pepcid AC
Offer
(Mail-in offer. See details inside.)

0000000

Drug Facts (continued)

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 gelcap with a glass of water
 - to **prevent** symptoms, swallow 1 gelcap with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 gelcaps in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton and package insert. They contain important information.
- store at 25°-30°C (77°-86°F)
- protect from moisture

Inactive ingredients

benzyl alcohol, black iron oxide, butylparaben, castor oil, edetate calcium disodium, FD&C red #40, gelatin, hydroxypropyl methylcellulose, magnesium stearate, methylparaben, microcrystalline cellulose, pregelatinized corn starch, propylene glycol, propylparaben, sodium lauryl sulfate, sodium propionate, talc, titanium dioxide

Questions or comments?
1-800-755-4008



Pepcid
AC
Famotidine Tablets 10mg/Acid Reducer



Gelcaps

Just One Per Dose

Relieves & Prevents Heartburn Due To Acid Indigestion

50 Gelcaps (Gelatin Coated, Capsule Shaped Tablets)

NEW!
Easy-Open Bottle

Pepcid
AC
Gelcaps

Relieves & Prevents Heartburn Due To Acid Indigestion

Distributed by:
Johnson & Johnson • MERCK
CONSUMER PHARMACEUTICALS CO.
FORT WASHINGTON, PA 19034 USA

If You Have Questions Or Comments
Please Call 1-800-755-4008

©Registered trademark of Merck & Co., Inc.



Pepcid
AC
Gelcaps

Do not use if carton is open or printed foil seal under bottle cap is open or torn.

PEPCID AC is Now Available in an Easy-to-Swallow Gelcap

- **1 gelcap** relieves heartburn due to acid indigestion (Read Package Insert before use).
- **PEPCID AC** prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Drug Facts

Active ingredient (in each gelcap) Purpose
Famotidine 10 mg.....Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble swallowing
- with other acid reducers

Stop use and ask a doctor if

- stomach pain continues
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

LOT AND EXP. DATE
STAMPED HERE

Do not use if printed foil seal under bottle cap is open or torn.

Active ingredient (in each gelcap) Famotidine 10 mg **Purpose** Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use • if you have trouble swallowing • with other acid reducers

Stop use and ask a doctor if • stomach pain continues • you need to take this product for more than 14 days

LOT AND EXP. DATE STAMPED HERE

Pepcid

AC

Famotidine Tablets 10mg/Acid Reducer®

Gelcaps

Relieves & Prevents Heartburn Due To Acid Indigestion

50 Gelcaps (Gelatin Coated, Capsule Shaped Tablets) **Just One Per Dose**

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** • adults and children 12 years and over: • to **relieve** symptoms, swallow 1 gelcap with a glass of water • to **prevent** symptoms, swallow 1 gelcap with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn • do not use more than 2 gelcaps in 24 hours • children under 12 years: ask a doctor

Other information • read the directions, warnings and accompanying product information before use • keep the carton and package insert. They contain important information. • store at 25°-30°C (77°-86°F) • protect from moisture

Dist. by: **Johnson & Johnson** • MERCK 0000000
CONSUMER PHARMACEUTICALS CO. ®Registered trademark
FORT WASHINGTON, PA 19004 USA of Merck & Co., Inc.

Do not use if printed foil seal under bottle cap is open or torn.

Active ingredient (in each gelcap) Famotidine 10 mg **Purpose** Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use • if you have trouble swallowing • with other acid reducers

Stop use and ask a doctor if • stomach pain continues • you need to take this product for more than 14 days

LOT AND EXP. DATE STAMPED HERE

Pepcid

AC

Famotidine Tablets 10mg/Acid Reducer®

Gelcaps

Relieves & Prevents Heartburn Due To Acid Indigestion

50 Gelcaps (Gelatin Coated, Capsule Shaped Tablets) **Just One Per Dose**

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** • adults and children 12 years and over: • to **relieve** symptoms, swallow 1 gelcap with a glass of water • to **prevent** symptoms, swallow 1 gelcap with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn • do not use more than 2 gelcaps in 24 hours • children under 12 years: ask a doctor

Other information • read the directions, warnings and accompanying product information before use • keep the carton and package insert. They contain important information. • store at 25°-30°C (77°-86°F) • protect from moisture

Dist. by: **Johnson & Johnson** • MERCK 0000000
CONSUMER PHARMACEUTICALS CO. ®Registered trademark
FORT WASHINGTON, PA 19004 USA of Merck & Co., Inc.

200%