

## ***Drug Facts***

### ***Active Ingredient (in each tablet)***

ranitidine hydrochloride 84 mg; equivalent to 75 mg ranitidine.

### ***Purpose***

**Acid reducer**

### ***Uses***

- relieves heartburn associated with acid indigestion and sour stomach.
- prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages.

### ***Warnings***

**Allergy Alert:** Do not use if you are allergic to Zantac (ranitidine hydrochloride) or other acid reducers

Do not use with other acid reducers

When using this product, do not take the maximum daily dose for more than 14 consecutive days, unless directed by your doctor.

Stop use and ask a doctor If you have difficulty swallowing or persistent abdominal pain. You may have a serious condition that may need different treatment.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

### ***Directions***

- for **relief** of symptoms swallow 1 tablet with a glass of water.
- to **prevent** of symptoms swallow 1 tablet with a glass of water 30 to 60 minutes **before** eating food or drinking beverages that cause heartburn.
- can be used up to twice daily (up to 2 tablets in 24 hours).
- this product should not be given to children under 12 years old unless directed by a doctor.

### ***Other Information***

- do not use if the blister unit is open or torn
- store at 20 - 25 C (68 - 77 F)
- Zantac 75 tablets are sodium and sugar free
- Avoid excessive heat or humidity

### ***Inactive Ingredients***

hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, synthetic red iron oxide, titanium dioxide and triacetin.

**Questions or comments?** Call 1-800-223-0182. Information is available 24 hours a day, 7 days a week.