


## NARMS

**Figure 15: Minimum Inhibitory Concentration of Gentamicin for *Enterococcus* in Chicken Breast (N=381 Isolates)**


**Breakpoints: Susceptible <500  $\mu\text{g}/\text{mL}$  Resistant  $\geq 500 \mu\text{g}/\text{mL}$**



## NARMS

**Figure 15: Minimum Inhibitory Concentration of Gentamicin for *Enterococcus* in Ground Turkey (N=387 Isolates)**


**Breakpoints: Susceptible <500  $\mu\text{g}/\text{mL}$  Resistant  $\geq 500 \mu\text{g}/\text{mL}$**



## NARMS

**Figure 15: Minimum Inhibitory Concentration of Gentamicin for *Enterococcus* in Ground Beef (N=383 Isolates)**

**Breakpoints: Susceptible < 500  $\mu\text{g/mL}$  Resistant  $\geq$  500  $\mu\text{g/mL}$**



## NARMS

**Figure 15: Minimum Inhibitory Concentration of Gentamicin for *Enterococcus* in Pork Chop (N=369 Isolates)**

**Breakpoints: Susceptible <500  $\mu\text{g}/\text{mL}$  Resistant  $\geq 500 \mu\text{g}/\text{mL}$**

