

Table 30. *Enterococcus* Species by Site and Meat Type, 2002

		Chicken Breast		Ground Turkey		Ground Beef		Pork Chop	
Site	Species	n	%	#	%	#	%	#	%
GA	<i>faecalis</i> (n=393)	84	21.4%	118	30.0%	85	21.6%	106	27.0%
	<i>faecium</i> (n=47)	27	57.4%	2	4.3%	7	14.9%	11	23.4%
	<i>hirae</i> (n=33)	7	21.2%		0.0%	24	72.7%	2	6.1%
	<i>avium</i> (n=2)	2	100.0%		0.0%		0.0%		0.0%
	<i>durans</i> (n=2)		0.0%		0.0%	2	100.0%		0.0%
	Total (n=477)	120	25.2%	120	25.2%	118	24.7%	119	24.9%
MD	<i>faecalis</i> (n=117)	10	8.5%	38	32.5%	31	26.5%	38	32.5%
	<i>faecium</i> (n=284)	105	37.0%	74	26.1%	56	19.7%	49	17.3%
	<i>hirae</i> (n=31)	1	3.2%	1	3.2%	20	64.5%	9	29.0%
	<i>avium</i> (n=1)	1	100.0%		0.0%		0.0%		0.0%
	<i>durans</i> (n=5)		0.0%		0.0%		0.0%	5	100.0%
	Total (n=438)	117	26.7%	113	25.8%	107	24.4%	101	23.1%
OR	<i>faecalis</i> (n=115)	21	18.3%	35	30.4%	22	19.1%	37	32.2%
	<i>faecium</i> (n=27)	17	63.0%	4	14.8%	4	14.8%	2	7.4%
	<i>hirae</i> (n=16)	2	12.5%		0.0%	14	87.5%		0.0%
	<i>gallinarum</i> (n=1)		0.0%	1	100.0%		0.0%		0.0%
	Total (n=159)	40	25.2%	40	25.2%	40	25.2%	39	24.5%
TN	<i>faecalis</i> (n=268)	19	7.1%	103	38.4%	72	26.9%	74	27.6%
	<i>faecium</i> (n=148)	82	55.4%	9	6.1%	26	17.6%	31	20.9%
	<i>hirae</i> (n=22)	2	9.1%	1	4.5%	18	81.8%	1	4.5%
	<i>avium</i> (n=1)		0.0%		0.0%	1	100.0%		0.0%
	<i>durans</i> (n=3)	1	33.3%		0.0%	1	33.3%	1	33.3%
	<i>gallinarum</i> (n=4)		0.0%	1	25.0%		0.0%	3	75.0%
	Total (n=446)	104	23.3%	114	25.6%	118	26.5%	110	24.7%