

Table 11. Antimicrobial Resistance among *Salmonella* Isolates by Meat Type,* 2002.

<i>Antimicrobial Agent</i>	<i>Chicken Breast (n=60)</i>	<i>Ground Turkey (n=74)</i>	<i>Ground Beef (n=9)</i>	<i>Pork Chop (n=10)</i>
Tetracycline	33.3%	55.4%	22.2%	70.0%
Streptomycin	28.3%	37.8%	22.2%	70.0%
Sulfamethoxazole	16.7%	20.3%	22.2%	70.0%
Ampicillin	16.7%	16.2%	22.2%	40.0%
Cephalothin	13.3%	14.9%	22.2%	20.0%
Gentamicin	10.0%	14.9%		30.0%
Amoxicillin/Clavulanic Acid	10.0%	12.2%	22.2%	20.0%
Kanamycin	6.7%	18.9%		10.0%
Cefoxitin	10.0%	8.1%	22.2%	20.0%
Ceftiofur	10.0%	8.1%	22.2%	20.0%
Chloramphenicol		1.4%	22.2%	40.0%
Nalidixic Acid		8.1%		
Trimethoprim/Sulfamethoxazole		1.4%		20.0%

* No resistance seen to Amikacin, Ciprofloxacin or Ceftriaxone for these isolates.