

Table 15. Antimicrobial Resistance among *Salmonella* by Site, Meat Type, and Antimicrobial Agent, 2002.

Site	Meat Type	Antimicrobial Agent												
		TET	STR	SMX	AMP	CEP	GEN	AMC	KAN	FOX	TIO	CHL	NAL	COT
CT	CB (n=17)	29.4%	11.8%	29.4%	23.5%	11.8%	5.9%	5.9%		5.9%	5.9%			
	GT (n=21)	52.4%	57.1%	47.6%	42.9%	38.1%	33.3%	33.3%	42.9%	19.1%	19.1%		19.1%	
	GB (n=5)	20.0%	20.0%	20.0%	20.0%	20.0%		20.0%		20.0%	20.0%	20.0%		
	PC (n=1)													
	Total (n=44)	38.6%	34.1%	36.4%	31.8%	25.0%	18.2%	20.5%	20.5%	13.6%	13.6%	2.3%	9.1%	0.0%
GA	CB (n=14)	28.6%	21.4%	7.1%	7.1%	7.1%	7.1%							
	GT (n=19)	52.6%	26.3%		5.3%	5.3%			5.3%					
	GB (n=2)													
	PC (n=2)	100.0%	50.0%	50.0%	50.0%				50.0%			50.0%		
	Total (n=37)	43.2%	24.3%	5.4%	8.1%	5.4%	2.7%	0.0%	5.4%	0.0%	0.0%	2.7%	0.0%	0.0%
MD	CB (n=8)	25.0%	25.0%		50.0%	50.0%		50.0%		50.0%	50.0%			
	GT (n=9)	33.3%	22.2%	22.2%	11.1%	11.1%	11.1%	11.1%	11.1%	11.1%	11.1%	11.1%	22.2%	11.1%
	GB (n=2)	50.0%	50.0%	50.0%	50.0%	50.0%		50.0%		50.0%	50.0%	50.0%		
	PC (n=6)	83.3%	100.0%	100.0%	50.0%	33.3%	50.0%	33.3%		33.3%	33.3%	50.0%		33.3%
	Total (n=25)	44.0%	44.0%	36.0%	36.0%	32.0%	16.0%	32.0%	4.0%	32.0%	32.0%	20.0%	8.0%	12.0%
MN	CB (n=4)	25.0%	25.0%		25.0%	25.0%		25.0%		25.0%	25.0%			
	GT (n=7)	28.6%	42.9%	14.3%	14.3%	14.3%	14.3%	14.3%		14.3%	14.3%			
	GB (n=0)													
	PC (n=0)													
	Total (n=11)	27.3%	36.4%	9.1%	18.2%	18.2%	9.1%	18.2%	0.0%	18.2%	18.2%	0.0%	0.0%	0.0%
OR	CB (n=4)	50.0%	75.0%											
	GT (n=2)	100.0%												
	GB (n=0)													
	PC (n=0)													
	Total (n=6)	66.7%	50.0%	0.0%	0.0%	0.0%								
TN	CB (n=13)	46.2%	46.2%	30.8%			30.8%		30.8%					
	GT (n=16)	81.3%	37.5%	12.5%			12.5%		18.8%					
	GB (n=0)													
	PC (n=1)													
	Total (n=30)	63.3%	40.0%	20.0%	0.0%	0.0%	20.0%	0.0%	23.3%	0.0%	0.0%	0.0%	0.0%	0.0%
Total %R (N=153)		45.8%	35.3%	22.2%	18.3%	15.0%	13.1%	12.4%	12.4%	10.5%	10.5%	4.6%	3.9%	2.0%