

NDA 13-684/S-092, 16-677/S-139, 16-678/S-100, 16-679/S-099, 16-682/S-100, 16-683/S-096, 16-687/S-097, 16-689/S-100, 16-692/S-091, 16-693/S-091, 16-695/S-093, 16-696/S-094, 16-697/S-093, 17-378/S-063, 17-385/S-055, 17-390/S-060, 17-438/S-059, 17-451/S-058, 17-484/S-062, 17-634/S-065, 17-648/S-065, 18-008/S-065, 18-016/S-057, 18-037/S-065, 18-840/S-029, 19-022/S-022, 19-047/S-024, 19-308/S-022, 19-367/S-022

Page 123

Baxter

PLASMA-LYTE R INJECTION (Multiple Electrolytes Injection, Type 2, USP) in AVIVA Plastic Container

Description

PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) is a sterile, nonpyrogenic isotonic solution in a single dose container for intravenous administration. Each 100 mL contains 640 mg of Sodium Acetate Trihydrate, USP ($C_2H_3NaO_2 \cdot 3H_2O$); 496 mg of Sodium Chloride, USP (NaCl); 89.6 mg of Sodium Lactate ($C_3H_5NaO_3$); 74.6 mg of Potassium Chloride, USP (KCl); 36.8 mg of Calcium Chloride, USP ($CaCl_2 \cdot 2H_2O$); and 30.5 mg of Magnesium Chloride, USP ($MgCl_2 \cdot 6H_2O$). It contains no antimicrobial agents. The pH is adjusted with hydrochloric acid. The pH is 5.5 (4.0 to 8.0).

PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) administered intravenously has value as a source of water, electrolytes, and calories. One liter has an ionic concentration of 140 mEq sodium, 10 mEq potassium, 5 mEq calcium, 3 mEq magnesium, 103 mEq chloride, 47 mEq acetate, and 8 mEq lactate. The osmolarity is 312 mOsmol/L (calc). Normal physiologic osmolarity range is approximately 280 to 310 mOsmol/L. Administration of substantially hypertonic solutions may cause vein damage. The caloric content is 11 kcal/L.

The flexible container is made with non-latex plastic materials specially designed for a wide range of parenteral drugs including those requiring delivery in containers made of polyolefins or polypropylene. For example, the AVIVA container system is compatible with and appropriate for use in the admixture and administration of paclitaxel. In addition, the AVIVA container system is compatible with and appropriate for use in the admixture and administration of all drugs deemed compatible with existing polyvinyl chloride container systems. The solution contact materials do not contain PVC, DEHP, or other plasticizers.

The suitability of the container materials has been established through biological evaluations, which have shown the container passes Class VI U.S. Pharmacopeia (USP) testing for plastic containers. These tests confirm the biological safety of the container system.

The flexible container is a closed system, and air is prefilled in the container to facilitate drainage. The container does not require entry of external air during administration.

The container has two ports: one is the administration outlet port for attachment of an intravenous administration set and the other port has a medication site for addition of supplemental medication (See Directions for Use). The primary function of the overwrap is to protect the container from the physical environment.

Clinical Pharmacology

NDA 13-684/S-092, 16-677/S-139, 16-678/S-100, 16-679/S-099, 16-682/S-100, 16-683/S-096, 16-687/S-097, 16-689/S-100, 16-692/S-091, 16-693/S-091, 16-695/S-093, 16-696/S-094, 16-697/S-093, 17-378/S-063, 17-385/S-055, 17-390/S-060, 17-438/S-059, 17-451/S-058, 17-484/S-062, 17-634/S-065, 17-648/S-065, 18-008/S-065, 18-016/S-057, 18-037/S-065, 18-840/S-029, 19-022/S-022, 19-047/S-024, 19-308/S-022, 19-367/S-022

Page 124

PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) has value as a source of water and electrolytes. It is capable of inducing diuresis depending on the clinical condition of the patient.

PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) produces a metabolic alkalinizing effect. Acetate and lactate ions are metabolized ultimately to carbon dioxide and water, which requires the consumption of hydrogen cations.

Indications and Usage

PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) is indicated as a source of water and electrolytes or as an alkalinizing agent.

Contraindications

None known

Warnings

PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) should be used with great care, if at all, in patients with congestive heart failure, severe renal insufficiency and in clinical states in which there exists edema with sodium retention.

PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) should be used with great care, if at all, in patients with hyperkalemia, severe renal failure and in conditions in which potassium retention is present.

PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) should be used with great care in patients with metabolic or respiratory alkalosis. The administration of lactate or acetate ions should be done with great care in those conditions in which there is an increased level or an impaired utilization of these ions, such as severe hepatic insufficiency.

PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) should not be administered simultaneously with blood through the same administration set because of the likelihood of coagulation.

The intravenous administration of PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) can cause fluid and/or solute overloading resulting in dilution of serum electrolyte concentrations, overhydration, congested states or pulmonary edema. The risk of dilutional states is inversely proportional to the electrolyte concentrations of the injection. The risk of solute overload causing congested states with peripheral and pulmonary edema is directly proportional to the electrolyte concentrations of the injection.

In patients with diminished renal function, administration of PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) may result in sodium or potassium retention. PLASMA-LYTE R

NDA 13-684/S-092, 16-677/S-139, 16-678/S-100, 16-679/S-099, 16-682/S-100, 16-683/S-096, 16-687/S-097, 16-689/S-100, 16-692/S-091, 16-693/S-091, 16-695/S-093, 16-696/S-094, 16-697/S-093, 17-378/S-063, 17-385/S-055, 17-390/S-060, 17-438/S-059, 17-451/S-058, 17-484/S-062, 17-634/S-065, 17-648/S-065, 18-008/S-065, 18-016/S-057, 18-037/S-065, 18-840/S-029, 19-022/S-022, 19-047/S-024, 19-308/S-022, 19-367/S-022

Page 125

Injection (Multiple Electrolytes Injection, Type 2, USP) is not for use in the treatment of lactic acidosis.

Precautions

General

Do not connect flexible plastic containers of intravenous solutions in series, i.e., do not piggyback connections. Such use could result in air embolism due to residual air being drawn from one container before administration of the fluid from a secondary container is completed.

Pressurizing intravenous solutions contained in flexible plastic containers to increase flow rates can result in air embolism if the residual air in the container is not fully evacuated prior to administration.

Use of a vented intravenous administration set with the vent in the open position could result in air embolism. Vented intravenous administration sets with the vent in the open position should not be used with flexible plastic containers.

PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) should be used with caution. Excess administration may result in metabolic alkalosis.

Laboratory Tests

Clinical evaluation and periodic laboratory determinations are necessary to monitor changes in fluid balance, electrolyte concentrations, and acid base balance during prolonged parenteral therapy or whenever the condition of the patient warrants such evaluation.

Drug Interactions

Caution must be exercised in the administration of PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) to patients receiving corticosteroids or corticotropin.

Studies have not been conducted to evaluate additional drug/drug or drug/food interactions with PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP).

Carcinogenesis, mutagenesis, impairment of fertility

Studies with PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) have not been performed to evaluate carcinogenic potential, mutagenic potential, or effects on fertility.

Pregnancy: Teratogenic Effects

Pregnancy Category C. Animal reproduction studies have not been conducted with PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP). It is also not known whether PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) should be given to a pregnant woman only if clearly needed.

Labor and Delivery

NDA 13-684/S-092, 16-677/S-139, 16-678/S-100, 16-679/S-099, 16-682/S-100, 16-683/S-096, 16-687/S-097, 16-689/S-100, 16-692/S-091, 16-693/S-091, 16-695/S-093, 16-696/S-094, 16-697/S-093, 17-378/S-063, 17-385/S-055, 17-390/S-060, 17-438/S-059, 17-451/S-058, 17-484/S-062, 17-634/S-065, 17-648/S-065, 18-008/S-065, 18-016/S-057, 18-037/S-065, 18-840/S-029, 19-022/S-022, 19-047/S-024, 19-308/S-022, 19-367/S-022

Page 126

Studies have not been conducted to evaluate the effects of PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) on labor and delivery. Caution should be exercised when administering this drug during labor and delivery.

Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) is administered to a nursing woman.

Pediatric Use

Safety and effectiveness of PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) in pediatric patients have not been established by adequate and well controlled trials, however, the use of electrolyte solutions in the pediatric population is referenced in the medical literature. The warnings, precautions and adverse reactions identified in the label copy should be observed in the pediatric population.

Geriatric Use

Clinical studies of PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

Adverse Reactions

Reactions which may occur because of the solution or the technique of administration include febrile response, infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection, extravasation and hypervolemia.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures and save the remainder of the fluid for examination if deemed necessary.

Dosage and Administration

As directed by a physician. Dosage is dependent upon the age, weight and clinical condition of the patient as well as laboratory determinations.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

NDA 13-684/S-092, 16-677/S-139, 16-678/S-100, 16-679/S-099, 16-682/S-100,
16-683/S-096, 16-687/S-097, 16-689/S-100, 16-692/S-091, 16-693/S-091,
16-695/S-093, 16-696/S-094, 16-697/S-093, 17-378/S-063, 17-385/S-055,
17-390/S-060, 17-438/S-059, 17-451/S-058, 17-484/S-062, 17-634/S-065,
17-648/S-065, 18-008/S-065, 18-016/S-057, 18-037/S-065, 18-840/S-029,
19-022/S-022, 19-047/S-024, 19-308/S-022, 19-367/S-022

Page 127

Do not administer unless solution is clear and seal is intact.

All injections in AVIVA plastic containers are intended for intravenous administration using sterile equipment.

Additives may be incompatible. Complete information is not available. Those additives known to be incompatible should not be used. Consult with pharmacist, if available. If, in the informed judgment of the physician, it is deemed advisable to introduce additives, use aseptic technique. Mix thoroughly when additives have been introduced. Do not store solutions containing additives.

How Supplied

PLASMA-LYTE R Injection (Multiple Electrolytes Injection, Type 2, USP) in AVIVA plastic containers is available as shown below:

Code	Size (mL)	NDC
6E2504	1000	NDC 0338-6320-04

Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat. It is recommended the product be stored at room temperature (25°C); brief exposure up to 40°C does not adversely affect the product.

Directions for Use of AVIVA Plastic Container

To Open

Tear overwrap down side at slit and remove solution container. Moisture and some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. Check for minute leaks by squeezing inner bag firmly. If leaks are found, discard solution as sterility may be impaired. If supplemental medication is desired, follow directions below.

Preparation for Administration

Caution: Do not use plastic containers in series connections.

Caution: Use only with a non-vented set or a vented set with the vent closed.

1. Suspend container from eyelet support.
2. Remove protector from outlet port at bottom of container.
3. Attach administration set. Refer to complete directions accompanying set.

To Add Medication

Additives may be incompatible.

NDA 13-684/S-092, 16-677/S-139, 16-678/S-100, 16-679/S-099, 16-682/S-100,
16-683/S-096, 16-687/S-097, 16-689/S-100, 16-692/S-091, 16-693/S-091,
16-695/S-093, 16-696/S-094, 16-697/S-093, 17-378/S-063, 17-385/S-055,
17-390/S-060, 17-438/S-059, 17-451/S-058, 17-484/S-062, 17-634/S-065,
17-648/S-065, 18-008/S-065, 18-016/S-057, 18-037/S-065, 18-840/S-029,
19-022/S-022, 19-047/S-024, 19-308/S-022, 19-367/S-022

Page 128

To add medication before solution administration

1. Prepare medication site.
2. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
3. Mix solution and medication thoroughly. For high density medication such as potassium chloride, squeeze ports while ports are upright and mix thoroughly.

To add medication during solution administration

1. Close clamp on the set.
2. Prepare medication site.
3. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
4. Remove container from IV pole and/or turn to an upright position.
5. Evacuate both ports by squeezing them while container is in the upright position.
6. Mix solution and medication thoroughly.
7. Return container to in use position and continue administration.

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Printed in USA

* Bar Code Position Only

XXXXXXXX

©Copyright XXXX Baxter Healthcare Corporation

All rights reserved.

X-XX-XX-XXX

Rev. August 2005

Baxter, PLASMA-LYTE, and AVIVA are trademarks of Baxter International Inc.