

NDC 59130-704-78

OXYTETRACYCLINE HCl SOLUBLE POWDER-343

A broad spectrum
ANTIBIOTIC

**For control and treatment of specific diseases in poultry, cattle,
sheep, swine and bees.**

This packet contains 102.4 grams of oxytetracycline HCl

This packet will make:

512 gallons (1,938 L) containing 200 mg oxytetracycline HCl per gallon

256 gallons (969 L) containing 400 mg oxytetracycline HCl per gallon

128 gallons (484.5 L) containing 800 mg oxytetracycline HCl per gallon

For oral use only in poultry, cattle, sheep, swine and bees.

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

NET WEIGHT: 135.5 g (4.78 oz)

ANADA 200-247, APPROVED BY FDA

AmTech®
Group, Inc.

200-247

F0151

INDICATIONS AND DIRECTIONS FOR USE

For the control of the following poultry diseases caused by organisms susceptible to oxytetracycline: Add the following amount to two gallons of stock solution when proportioner is set to meter at the rate of one ounce per gallon.

CHICKENS	DOSAGE	PACKETS/2 GALLONS STOCK SOLUTION
Infectious synovitis caused by <i>Mycoplasma synoviae</i>	200-400 mg/gal	1/4-1/2
Chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i>	400-800 mg/gal	1/2-1
Fowl cholera caused by <i>Pasteurella multocida</i>	400-800 mg/gal	1/2-1

TURKEYS

Hexamitiasis caused by <i>Hexamita meleagridis</i>	200-400 mg/gal	1/4-1/2
Infectious synovitis caused by <i>Mycoplasma synoviae</i>	400 mg/gal	1/2

Growing Turkeys—Complicating bacterial organisms associated with bluecomb (transmissible enteritis, body weight daily consumption (1 packet will treat 8,192 pounds of turkeys.)
coronaviral enteritis)

Medicate continuously at the first clinical signs of disease and continue for 7 to 14 consecutive days. If improvement is not noted within 24-48 hours, consult a poultry diagnostic laboratory or poultry pathologist to determine diagnosis and advice on dosage.

For the control and treatment of the following diseases caused by organisms susceptible to oxytetracycline:

SWINE

Bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis*

Bacterial pneumonia caused by *Pasteurella multocida*

For Breeding Swine: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by *Leptospira pomona*.

DOSAGE
Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days.

CALVES, BEEF CATTLE AND NON-LACTATING DAIRY CATTLE

Bacterial enteritis caused by *Escherichia coli*
Bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocida*.

Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days.

SHEEP

Bacterial enteritis caused by *Escherichia coli*
Bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocida*.

Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days.

This packet will treat 20,480 pounds of swine, cattle or sheep at 10 mg/pound.

HONEY BEES

American and European Foulbrood caused by *Bacillus larvae* susceptible to oxytetracycline.

200 mg/colony
Remove at least 6 weeks prior to main honey flow.

CAUTION: Use as sole source of oxytetracycline. Prepare fresh solutions every 24 hours.

Special Note: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate and the environmental temperature and humidity, each of which affects water consumption.

WARNING: Do not administer to turkeys, cattle or sheep within 5 days of slaughter. Zero-day slaughter withdrawal in swine. Do not administer to chickens or turkeys producing eggs for human consumption. Do not administer this product with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers.

A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

RECOMMENDED STORAGE: STORE BELOW 77°F (25°C)

FOR USE IN DRINKING WATER ONLY

NOT FOR USE IN LIQUID FEED SUPPLEMENTS

FOR ANIMAL USE ONLY. KEEP OUT OF REACH OF CHILDREN

Restricted Drug(s) (California). Not For Human Use. Use Only As Directed.

Manufactured by
Phoenix Scientific, Inc.
St. Joseph, MO 64503

800507
Rev. 9-02

Lot No.

Exp. Date



NDC 59130-704-27

OXYTETRACYCLINE HCl SOLUBLE POWDER-343

A broad spectrum
ANTIBIOTIC

**For control and treatment of specific diseases
in poultry, cattle, sheep, swine and bees.**

This packet contains 204.8 grams of oxytetracycline HCl

This packet will make:

1,024 gallons (3,876 L) containing 200 mg oxytetracycline HCl per gallon

512 gallons (1,938 L) containing 400 mg oxytetracycline HCl per gallon

256 gallons (969 L) containing 800 mg oxytetracycline HCl per gallon

For oral use only in poultry, cattle, sheep, swine and bees.

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

NET WEIGHT: 272.2 g (9.6 oz)

ANADA 200-247, APPROVED BY FDA

AmTech[®]
Group, Inc.

1797 BK

23444 amtech 3/30/00 133-2/00

INDICATIONS AND DIRECTIONS FOR USE

For the control of the following poultry diseases caused by organisms susceptible to oxytetracycline: Add the following amount to two gallons of stock solution when proportioner is set to meter at the rate of one ounce per gallon.

	DOSAGE	PACKETS/2 GALLONS STOCK SOLUTION
CHICKENS		
Infectious synovitis caused by <i>Mycoplasma synoviae</i>	200-400 mg/gal	1/2-1
Chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i>	400-800 mg/gal	1-2
Fowl cholera caused by <i>Pasteurella multocida</i>	400-800 mg/gal	1-2
TURKEYS		
Hexamitiasis caused by <i>Hexamita meleagridis</i>	200-400 mg/gal	1/2-1
Infectious synovitis caused by <i>Mycoplasma synoviae</i>	400 mg/gal	1
Growing Turkeys—Complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis)	25 mg/lb body weight daily	varies with age & water consumption (1 packet will treat 4,096 pounds of turkeys.)

Medicate continuously at the first clinical signs of disease and continue for 7 to 14 consecutive days. If improvement is not noted within 24-48 hours, consult a poultry diagnostic laboratory or poultry pathologist to determine diagnosis and advice on dosage.

For the control and treatment of the following diseases caused by organisms susceptible to oxytetracycline:

SWINE

Bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis*
 Bacterial pneumonia caused by *Pasteurella multocida*
 For Breeding Swine: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by *Leptospira pomona*.

DOSAGE
 Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days.

CALVES, BEEF CATTLE AND NON-LACTATING DAIRY CATTLE

Bacterial enteritis caused by *Escherichia coli*
 Bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocida*.

Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days.

SHEEP

Bacterial enteritis caused by *Escherichia coli*
 Bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocida*.

Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days.

This packet will treat 10,240 pounds of swine, cattle or sheep at 10 mg/pound.

HONEY BEES

American and European Foulbrood caused by *Bacillus larvae* susceptible to oxytetracycline.

200 mg/colony
 Remove at least 6 weeks prior to main honey flow.

CAUTION: Use as sole source of oxytetracycline. Prepare fresh solutions every 24 hours.

Special Note: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate and the environmental temperature and humidity, each of which affects water consumption.

WARNING: Do not administer to turkeys, cattle or sheep within 5 days of slaughter. Zero-day slaughter withdrawal in swine. Do not administer to chickens or turkeys producing eggs for human consumption. Do not administer this product with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. A withdrawal period has not been established for this product in prenatating calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.



Manufactured by
 Phoenix Scientific, Inc.
 St. Joseph, MO 64503

800507
 Iss. 8-02

RECOMMENDED STORAGE: STORE BELOW 77°F (25°C)
FOR USE IN DRINKING WATER ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS
FOR ANIMAL USE ONLY. KEEP OUT OF REACH OF CHILDREN
 Restricted Drug(s) (California). Not For Human Use. Use Only As Directed.



Lot No.
 Exp. Date

TO OPEN CUT HERE

5633

Terramycin-343[®]

(oxytetracycline HCl)
Soluble Powder

A broad-spectrum

antibiotic

for control and treatment of specific diseases
in poultry, cattle, swine, sheep, and bees.

This packet contains 204.8 grams of oxytetracycline HCl and will make:
1024 gallons (3876 L) containing
200 mg oxytetracycline HCl per gallon
512 gallons (1938 L) containing
400 mg oxytetracycline HCl per gallon
256 gallons (969 L) containing
800 mg oxytetracycline HCl per gallon

For oral use only

NADA #8-622, Approved by FDA

Net Weight: 9.55 oz (270.7 g)



For the control and treatment of the following diseases caused by organisms susceptible to oxytetracycline:

SWINE

Bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis*
Bacterial pneumonia caused by *Pasteurella multocida*
For Breeding Swine: Leptospirosis (reducing the incidence of abortions and shedding of leptospiral) caused by *Leptospira pomona*

CALVES, BEEF CATTLE AND NON-LACTATING DAIRY CATTLE

Bacterial enteritis caused by *Escherichia coli*
Bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocida*

SHEEP

Bacterial enteritis caused by *Escherichia coli*
Bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocida*

HONEY BEES

For control of American Foulbrood caused by *Bacillus larvae*

DOSEAGE

Administer in the drinking water at a level of 10 mg oxytetracycline HCl per lb of body weight daily. Administer up to 14 days.

Administer in the drinking water at a level of 10 mg oxytetracycline HCl per lb of body weight daily. Administer up to 14 days.

Administer in the drinking water at a level of 10 mg oxytetracycline HCl per lb of body weight daily. Administer up to 14 days.

200 mg/colony

The drug is administered in 3 applications of sugar syrup or 3 dustings at 4- to 5-day intervals. The drug should be fed early in the spring or fall and consumed by the bees before main honey flow begins to avoid contamination of production honey.

Remove at least 6 weeks prior to main honey flow.

CAUTION: Use as sole source of oxytetracycline. Prepare fresh solutions every 24 hours.

Special Note: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, and to offset the effect of water consumption.

992 01-4536-35-X1

00000

Flat: 11 1/2" (W) x 8 1/8" (H)
 Finished Packet: 5 3/4" (W) x 9" (H)

PMS 116

PMS 485

Black

01-4536-35-X1 Draft #1 2/23/99

INDICATIONS AND DIRECTIONS FOR USE

For the control of the following poultry diseases caused by organisms susceptible to oxytetracycline: Add the following amount to 2 gallons of stock solution when proportioner is set to meter at the rate of 1 ounce per gallon.

	DOSAGE	PACKS/2 GALLONS STOCK SOLUTION
CHICKENS Infectious synovitis caused by <i>Mycoplasma synoviae</i>	200-400 mg/gal	1/4-1/2
Chronic respiratory disease (CRD) and/or sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i>	400-800 mg/gal	1/2-1
Fowl cholera caused by <i>Pasteurella multocida</i>	400-800 mg/gal	1/2-1
TURKEYS Hexamitiasis caused by <i>Hexamita meleagridis</i>	200-400 mg/gal	1/4-1/2
Infectious synovitis caused by <i>Mycoplasma synoviae</i>	400 mg/gal	1/2
Growing Turkeys—Complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis)	25 mg/lb body weight daily	Varies with age & water consumption (1 packet will treat 8192 lb of turkeys)

Medicate continuously at the first clinical signs of disease and continue for 7-14 consecutive days. If improvement is not noted within 24-48 hours, consult a poultry diagnostic laboratory or poultry pathologist to determine diagnosis and advice on dosage.

WARNING: Do not administer to turkeys, cattle or sheep within 5 days of slaughter. Zero-day slaughter withdrawal in swine. Do not administer to chickens or turkeys producing eggs for human consumption. Do not administer this product with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers.

RECOMMENDED STORAGE: STORE BELOW 25°C (77°F)

FOR ANIMAL USE ONLY
 KEEP OUT OF REACH OF CHILDREN

Registered Brand (California) May Exp Human Use. Use Only As Directed.



Distributed by

Animal Health

Exton, PA 19341, USA
 Div. of Pfizer Inc.

TAKE TIME



OBSERVE LABEL DIRECTIONS