
Guidance for Industry

In Vivo Drug Metabolism/Drug Interaction Studies — Study Design, Data Analysis, and Recommendations for Dosing and Labeling

DRAFT GUIDANCE

This guidance document is being distributed for comment purposes only.

Comments and suggestions regarding this draft document should be submitted within 60 days of publication in the *Federal Register* of the notice announcing the availability of the draft guidance. Submit comments to Dockets Management Branch (HFA-305), Food and Drug Administration, 5630 Fishers Lane, rm. 1061, Rockville, MD 20852. All comments should be identified with the docket number listed in the notice of availability that publishes in the *Federal Register*. For questions regarding this draft document, contact Dr. Shiew-Mei Huang (CDER) at 301-594-5671; fax 301-594-2503; E-mail HuangSM@cder.fda.gov; or David Green (CBER) at 301-827-5349.

U.S. Department of Health and Human Services
Food and Drug Administration
Center for Drug Evaluation and Research (CDER)
Center for Biologics Evaluation and Research (CBER)
November 1998
Clin/Pharm #

J:\GUIDANC\2635DFT.WPD
10/28/98

98D-1001

GDL 1

Guidance for Industry

In Vivo Drug Metabolism/Drug Interaction Studies — Study Design, Data Analysis, and Recommendations for Dosing and Labeling

Additional copies are available from:
the Drug Information Branch (HFD-2 10),
Center for Drug Evaluation and Research (CDER),
5600 Fishers Lane, Rockville, MD 20857 (Tel) 301-827-4573
Internet at <http://www.fda.gov/cder/guidance/index.htm>

or
Office of Communication,
Training, and Manufacturers Assistance (HFM-40)
Center for Biologics Evaluation and Research (CBER)
1401 Rockville Pike, Rockville, MD 20852-1448
<http://www.fda.gov/cber/guidelines.htm>
(Fax) 888-CBERFAX or 301-827-3844
(Voice Information) 800-835-4709 or 301-827-1800

**U.S. Department of Health and Human Services
Food and Drug Administration
Center for Drug Evaluation and Research (CDER)
Center for Biologics Evaluation and Research (CBER)
November 1998
Clin/Pharm #**

DRAFT — NOT FOR IMPLEMENTATION

TABLE OF CONTENTS

I. INTRODUCTION	1
II. BACKGROUND	2
A. Metabolism	2
B. Metabolic Drug-Drug Interactions	2
III. GENERAL STRATEGIES	5
A. In Vitro Studies	5
B. Specific In Vivo Clinical Investigations	6
C. Population Pharmacokinetic Screens	6
IV. DESIGN OF IN VIVO METABOLIC DRUG-DRUG INTERACTION STUDIES	7
A. Study Design	7
B. Study Population	8
C. Choice of Substrate and Interacting Drugs	9
D. Route of Administration	10
E. Dose Selection	10
F. Endpoints	10
G. Sample Size and Statistical Considerations	11
V. LABELING	12
A. Drug Metabolism	12
B. Metabolic Drug-Drug Interaction Studies	12

DRAFT — NOT FOR IMPLEMENTATION

Guidance for Industry¹

**In Vivo Drug Metabolism/Drug Interaction Studies —
Study Design, Data Analysis, and Recommendations for Dosing and
Labeling**

I. INTRODUCTION

This guidance provides recommendations to sponsors/applicants of new drug applications (NDAs) and biologics license applications (BLAs) for therapeutic biologics (hereafter drugs) who intend to perform in vivo drug metabolism and metabolic drug-drug interaction studies. The guidance reflects the Agency's current view that the metabolism of an investigational new drug should be defined during drug development and that its interactions with other drugs should be explored as part of an adequate assessment of its safety and effectiveness. For metabolic drug-drug interactions, the approaches considered in the guidance are offered with the understanding that whether a particular study should be performed will vary depending on the drug in development and its intended clinical use. Furthermore, not every drug-drug interaction is metabolism-based, but may arise from changes in pharmacokinetics caused by absorption, binding, distribution, and excretion interactions. Although less well studied, drug-drug interactions may alter PK/PD relationships. These important areas are not considered in detail in this guidance.

Previous guidance from FDA on the use of in vitro approaches to study drug metabolism and metabolic drug-drug interactions is available in a guidance document entitled *Drug Metabolism/Drug Interaction Studies in the Drug Development Process: Studies In Vitro* (April 1997). The present guidance should be viewed as a companion to this earlier guidance. Discussion of metabolic and other types of drug-drug interactions is also provided in other guidances, including the International Conference on Harmonisation (ICH) *E8 General*

¹ This guidance has been prepared by the In Vivo Metabolic Drug-Drug Interaction Working Group in the Clinical Pharmacology Section of the Medical Policy Coordinating Committee in the Center for Drug Evaluation and Research, with input from the Center for Biologics Evaluation and Research at the Food and Drug Administration. This guidance document represents the Agency's current thinking on the subject of in vivo drug and metabolic drug-drug interactions. It does not create or confer any rights for or on any person and does not operate to bind FDA or the public. An alternative approach may be used if such approach satisfies the requirements of the applicable statute, regulations, or both.

DRAFT — NOT FOR IMPLEMENTATION

Considerations for Clinical Trials (December 1997), *E7 Studies in Support of Special Populations: Geriatrics* (August 1994), and *E3 Structure and Content of Clinical Study Reports* (July 1996), and the Agency guidances *Studying Drugs Likely to be Used in the Elderly* (November 1989) and *Study and Evaluation of Gender Differences in the Clinical Evaluation of Drugs* (July 1993).²

II. BACKGROUND

A. Metabolism

The desirable and undesirable effects of a drug arising from its concentrations at the sites of action are usually related either to the amount administered (dose) or to the resulting blood concentrations, which arise from its absorption, distribution, and metabolism and/or excretion. Elimination of a drug or its metabolizes occurs either by metabolism, usually by the liver, or by excretion, usually by the kidneys. Hepatic elimination occurs primarily by the cytochrome P450 family of enzymes located in the hepatic microsome but may also occur by non-P450 enzyme systems, such as N-acetyl and glucuronosyl transferases. P450 enzyme systems located in gut mucosa can also significantly affect the amount of drug absorbed into the systemic circulation.³ Many factors can alter hepatic and intestinal drug metabolism, including the presence or absence of disease and/or concomitant medications. While most of these factors are usually relatively stable over time, concomitant medications can alter metabolic routes of absorption and elimination abruptly and are of particular concern. The influence of concomitant medications on hepatic and intestinal metabolism becomes more complicated when a drug, including a prodrug, is metabolized to one or more active metabolizes. In this case, the safety and efficacy of the drug/prodrug are determined not only by exposure to the parent drug but by exposure to the active metabolizes, which in turn is related to their formation, distribution, and elimination.

B. Metabolic Drug-Drug Interactions

²See cover for information on how to obtain copies of these documents.

³No distinction is made in this document between the effects of concomitant drugs and/or alterations in metabolism on gastrointestinal absorption and hepatic elimination, although the pharmacokinetic effects of the two may be different.

DRAFT — NOT FOR IMPLEMENTATION

Many enzymatic metabolic routes of elimination, including most of those occurring via the P450 family of enzymes, can be inhibited or induced by concomitant drug treatment. Observed changes arising from metabolic drug-drug interactions can be substantial — an order of magnitude or more decrease or increase in the blood and tissue concentrations of a drug or metabolize — and can include formation of toxic metabolizes or increased exposure to a more toxic parent compound. Examples of substantially changed exposure associated with administration of another drug include (1) increased levels of terfenadine, cisapride, or astemizole with ketoconazole or erythromycin (inhibition of CYP3A4); (2) increased levels of simvastatin and its acid metabolize with mibefradil or itraconazole (inhibition of CYP3A4); (3) increased levels of desipramine with fluoxetine, paroxetine, or quinidine (inhibition of CYP2D6); and (4) decreased carbamazepine levels with rifampin (induction of CYP3A4). These large changes in exposure can alter the safety and efficacy profile of a drug and/or its active metabolizes in important ways. This is most obvious and expected for a drug with a narrow therapeutic range (NTR), but is also possible for non-NTR drugs as well (e.g., HMG COA reductase inhibitors and terfenadine). Depending on the extent and consequence of the interaction, the fact that a drug's metabolism can be significantly inhibited by other drugs and that the drug itself can inhibit the metabolism of other drugs can require important changes in either its dose or the doses of drugs with which it interacts, that is, on its labeled conditions of use. Rarely, metabolic drug-drug interactions may affect the ability of a drug to be safely marketed.

The following general concepts underlie the recommendations in this guidance:

- Adequate assessment of the safety and effectiveness of a drug includes a description of its metabolism and the contribution of metabolism to overall elimination.

Metabolic drug-drug interaction studies should explore whether an investigational agent is likely to affect importantly the metabolic elimination of drugs already in the marketplace and, conversely, whether drugs in the marketplace are likely to affect the metabolic elimination of the investigational drug.

Even drugs that are not substantially metabolized can have important effects on the metabolism of concomitant drugs. For this reason, metabolic drug-drug interactions should be explored even for an investigational compound that is not eliminated significantly by metabolism.

DRAFT — NOT FOR IMPLEMENTATION

- In some cases, metabolic drug-drug interaction studies cannot be informative unless metabolites and prodrugs have been identified and their pharmacological properties described.
- Identifying metabolic differences in patient groups based on genetic polymorphism, or on other readily identifiable factors, such as age, race, and gender, can aid in interpreting results.
- The impact of an investigational or approved interacting drug can be either to inhibit or induce metabolism.
- A specific objective of metabolic drug-drug interaction studies is to determine whether the interaction is sufficiently large to necessitate a dosage adjustment of the drug itself or the drugs it might be used with.
- In some instances, understanding how to adjust dosage in the presence of an interacting drug, or how to avoid such interactions, may allow marketing of a drug that would otherwise have been associated with an unacceptable level of toxicity. Sometimes a drug interaction may be used positively to increase levels or reduce elimination of another drug. Rarely, the degree of interaction caused by a drug, or the degree to which other drugs alter its metabolism, may be such that it cannot be marketed safely.
- The blood or plasma concentrations of the parent drug and/or its active metabolites (systemic exposure) provide an important link between drug dose (exposure) and desirable and/or undesirable drug effects. For this reason, the development of sensitive and specific assays for a drug and its metabolites is critical to the study of metabolism and drug-drug interactions.
- For drugs whose presystemic or systemic clearance occurs primarily by metabolism, differences arising from various sources, including administration of another drug, are an important source of inter-individual and intra-individual variability.
- Unlike relatively fixed influences on metabolism, such as hepatic function or genetic characteristics, metabolic drug-drug interactions can lead to abrupt changes in exposure. These effects can occur both at the time when a drug is initially prescribed and when it has been well-titrated and given at stable doses. Interactions can occur after even a single concomitant dose of an inhibitor.

DRAFT — NOT FOR IMPLEMENTATION

- The effects of a drug on the metabolism of other drugs and the effects of other drugs on a drug's metabolism should be assessed relatively early in drug development so that the clinical implications of interactions can be assessed as fully as possible in later studies.

III. GENERAL STRATEGIES

To the extent possible, drug development should follow a sequence where early in vitro and in vivo investigations can either fully address a question of interest or provide information to guide further studies. Optimally, a sequence of studies should be planned, moving from in vitro studies, to early exploratory studies, to later more definitive studies, employing special study designs and methodology where necessary and appropriate. In many cases, negative findings from early in vitro and early clinical studies can eliminate the need for later clinical investigations. Early investigations should explore whether a drug is eliminated primarily by excretion or metabolism, with identification of the principal metabolic routes in the latter case. Using suitable in vitro probes and careful selection of interacting drugs for early in vivo studies, the potential for drug-drug interactions can be studied early in the development process, with further study of observed interactions assessed later in the process as needed. In certain cases and with careful study designs and planning, these early studies may also provide information about dose, concentration, and response relationships in the general population, subpopulations, and individuals, which can be useful in interpreting the consequences of a metabolic drug-drug interaction.

A. In Vitro Studies

A complete understanding of the relationship between in vitro findings and in vivo results of metabolism/drug-drug interaction studies is still emerging. Nonetheless, in vitro studies can frequently serve as an adequate screening mechanism to rule out the importance of a metabolic pathway and drug-drug interactions that occur via this pathway so that subsequent in vivo testing is unnecessary. This opportunity should be based on appropriately validated experimental methods and rational selection of substrate/interacting drug concentrations. For example, if suitable in vitro studies indicate that CYP2D6 or 3A4 enzyme systems do not metabolize an investigational drug, then clinical studies to identify the impact of the CYP2D6 slow metabolize phenotype or to study the effect of CYP2D6 inhibitors or CYP3A4 inhibitors/inducers on the elimination of the investigational drug will not be needed. Similarly, if in vitro studies indicate that an investigational drug does not inhibit CYP2D6 or 3A4 metabolism, then in vivo drug-drug interaction studies of the investigational drug and concomitant medications eliminated by these pathways are not needed.

DRAFT — NOT FOR IMPLEMENTATION

In contrast, when positive findings arise in in vitro metabolic and/or drug-drug interaction studies, clinical studies are recommended because of the limited ability at present of in vitro findings to give a good quantitative estimate of the clinical importance of a metabolic pathway or interaction. Further evaluation of parameters such as the ratio between the unbound concentration of drug and the K_i (inhibition constant) for the interaction may lead to continued improvements in the ability of in vitro studies to predict in vivo results, but the overall experience to date is not large enough to allow reliable conclusions. Although in vitro studies can assess the presence or absence of inhibition, they have limited capability to identify induction. For this reason, in vivo studies remain the primary source of information about induction of metabolic pathways caused by concomitant medications.

B. Specific In Vivo Clinical Investigations

Appropriately designed pharmacokinetic and pharmacodynamic studies, usually performed in the early phases of drug development, can provide important information about metabolic routes of elimination, their contribution to overall elimination, and metabolic drug-drug interactions. Together with information from in vitro studies, these investigations can be a primary basis of labeling guidance and can often help avoid the need for further investigations. Further recommendations about these types of studies appear in Section IV of this guidance.

C. Population Pharmacokinetic Screens

Population pharmacokinetic analyses of data obtained from blood samples collected infrequently (sparse sampling) in clinical studies conducted in the later phase of drug development can be valuable in detecting unsuspected drug-drug interactions, in characterizing the clinical impact of known or newly identified interactions, and in guiding possible dosage modifications. Population pharmacokinetic data can also be valuable in confirming the absence of a pharmacokinetic drug-drug interaction when this is suggested by in vitro drug-drug interaction studies. The power of a sparse sampling strategy to detect drug-drug interactions is not yet well established, however, and it is unlikely that population analysis can be used to prove the absence of an interaction that is strongly suggested by information arising from in vitro or in vivo studies specifically designed to assess a drug-drug interaction. To be optimally informative, population pharmacokinetic studies should have carefully designed study and sample collections. A draft guidance for industry entitled *Population Pharmacokinetics* was published in September 1997 and currently is being finalized.⁴

⁴See cover for information on how to obtain a copy of this document.

DRAFT — NOT FOR IMPLEMENTATION

Iv. DESIGN OF IN VIVO METABOLIC DRUG-DRUG INTERACTION STUDIES

If in vitro studies and other information suggest a need for in vivo metabolic drug-drug interaction studies, the following general issues and approaches should be considered. In the following discussion, the term *substrate (S)* is used to indicate the drug studied to determine if its exposure is changed by another drug, which is termed the *interacting drug (I)*. Depending on the study objectives, the substrate and the interacting drug may be the investigational agents in the development or one or more approved products.

A. Study Design

In vivo metabolic drug-drug interaction studies generally are designed to compare substrate levels with and without the interacting drug. Because a specific study may consider a number of questions and clinical objectives, no one correct study design for studying drug-drug interactions can be defined. A study can use a randomized crossover (e.g., S followed by S+I, S+I followed by S), a one-sequence crossover (e.g., S always followed by S+I or the reverse), or a parallel design (S in one group of subjects and S +I in another). The following possible dose combinations for a substrate and interacting drug may also be used: single dose/single dose, single dose/multiple dose, multiple dose/single dose, and multiple dose/multiple dose. The selection of one of these or another study design depends upon a number of factors for both the substrate and interacting drug, including (1) acute or chronic use of the substrate and/or interacting drug; (2) safety considerations, including whether a drug is likely to be an NTR (narrow therapeutic range) or non-NTR drug; (3) pharmacokinetic and pharmacodynamic characteristics of the substrate and interacting drugs; and (4) the need to assess induction as well as inhibition. The inhibiting/inducing drugs and the substrates should be dosed so that the exposure of both drugs are relevant to their clinical use. The following considerations may be useful:

- Pharmacokinetic measures and/or parameters may be used to indicate clinically important routes of metabolism and drug-drug interactions. Subsequent interpretation of findings from these studies will be aided by good understanding of dose/concentration and concentration/response relationships for both desirable and undesirable drug effects in the general population, in subpopulations, and within individuals. In certain instances, reliance on endpoints other than pharmacokinetic measures/parameters may be useful.
- When both substrate and interacting drug are likely to be given chronically over an extended period of time, administration of the substrate to steady state with

DRAFT — NOT FOR IMPLEMENTATION

collection of blood samples over one or more dosing intervals could be followed by administration of the interacting drug, again with collection of blood for measurement of both the substrate and the interacting drug (as feasible) over the same intervals. This is an example of a one-sequence crossover design with or without replicate sample collection periods.

- The time at steady state before collection of endpoint observations depends on whether inhibition or induction is to be studied. Inducers can take several days or longer to exert their effects, while inhibitors generally exert their effects more rapidly. For this reason, a more extended period of time after attainment of steady state for the substrate or interacting drug may be necessary if induction is to be assessed.
- When attainment of steady state is important and either the substrate or interacting drugs and/or their metabolizes exhibit long half-lives, special approaches may be useful. These include use of a loading dose to achieve steady state conditions more rapidly and selection of a one-sequence crossover or a parallel design, rather than a randomized crossover study design.
- When a substrate and/or an interacting drug is to be studied at steady state, documentation that steady state has been attained is important both for each drug and its metabolizes of interest. This documentation can be accomplished by sampling over several days prior to the periods when samples are collected. This is important for both metabolizes and the parent drug, particularly when the half-life of the metabolize is longer than the parent, and is especially important if both parent drug and metabolizes are metabolic inhibitors or inducers.
- Studies can usually be open label (unblinded), unless pharmacodynamic endpoints (e.g., adverse events that are subject to bias) are part of the assessment of the interaction.

B. Study Population

Clinical drug-drug interaction studies may generally be performed using healthy volunteers or volunteers drawn from the general population, on the assumption that findings in this population should predict findings in the patient population for which the drug is intended. Safety considerations, however, may preclude the use of healthy subjects. In certain circumstances, subjects drawn from the general population and/or patients for whom the investigational drug is intended offer certain advantages, including the opportunity to study pharmacodynamic endpoints not present in healthy subjects and

DRAFT — NOT FOR IMPLEMENTATION

reduced reliance on extrapolation of findings from healthy subjects. In either patient or healthy/general population subject studies, performance of phenotype and genotype determinations to identify genetically determined metabolic polymorphisms is important in evaluating effects on enzymes with polymorphisms, notably CYP2D6 and CYP2C19.

C. Choice of Substrate and Interacting Drugs

1. Substrates for an Investigational Drug

In contrast to earlier approaches that focused mainly on a specific group of approved drugs (digoxin, hydrochlorothiazide) where co-administration was likely or the clinical consequences of an interaction were of concern, improved understanding of the metabolic basis of drug-drug interactions enables more general approaches to and conclusions from specific drug-drug interaction studies. In studying an investigational drug as the interacting drug, the choice of substrates (approved drugs) for initial in vivo studies depends on the P450 enzymes affected by the interacting drug. In testing inhibition, the substrate selected should generally be one whose metabolism by a specific enzyme is markedly inhibited by known inhibitors (i.e., a very sensitive substrate should be chosen) to assess the impact of the interacting investigational drug. Examples include (1) simvastatin or lovastatin for a CYP3A4 inhibitor; (2) theophylline for a CYP1A2 inhibitor; (3) warfarin for a CYP2C9 inhibitor; and (4) desipramine or imipramine for a CYP2D6 inhibitor. If the initial study is positive for inhibition, further studies of other substrates may be useful, representing a range of substrates based on the likelihood of co-administration. For example, possible substrates for further study of a CYP3A4 interacting investigational drug might include dihydropyridine calcium channel blockers and triazolobenzodiazepines, or for a CYP2D6 inhibiting investigational drug might include metoprolol. If the initial study is negative with the most sensitive substrates, it can be presumed that less sensitive substrates will also be unaffected.

2. Investigational Drug as Substrate

In testing an investigational drug for the possibility that its metabolism is inhibited or induced (i.e., as a substrate), selection of the interacting drugs should be based on in vitro or other metabolism studies identifying the enzyme systems that metabolize the drug. The choice of interacting drug should then be based on known, important inhibitors of the pathway under investigation. For example, if the investigational drug is shown to be metabolized by CYP3A4, the choice of inhibitor and inducer could be ketoconazole and rifampin, respectively, because of the substantial effects of these interacting drugs on CYP3A4 metabolism (i.e., they are the most sensitive in identifying an effect of interest). If the study results are negative, then absence of a clinically important drug-drug

DRAFT — NOT FOR IMPLEMENTATION

interaction for the metabolic pathway could be claimed. If the clinical study of the most potent specific inhibitor/inducer is positive and the sponsor wishes to claim lack of an interaction between the test drug and other less potent specific inhibitors, or give advice on dosage adjustment, further clinical studies would generally be recommended. Certain approved drugs are not optimal selections as the interacting drug. For example, cimetidine is not considered an optimal choice because its inhibition affects multiple pathways.

D. Route of Administration

The route of administration chosen for a metabolic drug-drug interaction study is important. For an investigational agent used as either an interacting drug or substrate, the route of administration should generally be the one planned for in product labeling. If only oral dosage forms will be marketed, studies with an intravenous formulation would not usually be necessary, although information from oral and intravenous dosings may be useful in discerning the relative contributions of alterations in absorption and/or presystemic clearance to the overall effect observed for a drug interaction. Sometimes certain routes of administration can reduce the utility of information from a study. For example, an intravenous study would generally not reveal an effect on intestinal CYP3A4 activity that markedly altered bioavailability. For an approved agent used either as a substrate or interacting drug, the route of administration will depend on available marketed formulations, which in most instances will be oral.

E. Dose Selection

For both a substrate (investigational drug or approved drug) and interacting drug (investigational drug or approved drug), testing should maximize the possibility of finding an interaction. For this reason, the maximum planned or approved dose and shortest dosing interval of the interacting drug should be used. Doses smaller than those to be used clinically may be needed for substrates on safety grounds.

F. Endpoints

1. Pharmacokinetic Endpoints

The following measures and parameters are recommended for assessment of the substrate: (1) exposure measures such as AUC, C_{max}, time to C_{max} (T_{max}), and others as appropriate; and (2) pharmacokinetic parameters such as clearance, volumes of distribution, and half lives. In some cases, these measures may be of interest for the inhibitor or inducer as well, notably where the study is assessing possible interactions

DRAFT — NOT FOR IMPLEMENTATION

between both study drugs. Additional measures may help in steady state studies (e.g., trough concentration (C_{min})) to demonstrate that dosing strategies were adequate to achieve steady state before and during the interaction. In certain instances, an understanding of the relationship between dose, blood levels and response may lead to a special interest in certain pharmacokinetic measures/parameters. For example, if a clinical outcome is most closely related to peak concentration (e.g., tachycardia with sympathomimetics), C_{max} or another early exposure measure might be most appropriate. Conversely, if the clinical outcome is related more to extent of absorption, AUC would be preferred. The frequency of sampling should be adequate to allow accurate determination of the relevant measures/parameters for the parent and metabolizes. For the substrate, whether the investigational drug or approved drug, determination of the pharmacokinetics of important active metabolizes is important. Because this guidance focuses on metabolic drug-drug interactions, protein binding determinations are considered unnecessary except for data interpretation.

2. Pharmacodynamic Endpoints

Pharmacokinetic measures are usually sufficient for metabolic drug-drug interaction studies, although pharmacodynamic measures can sometimes provide additional useful information. This may occur when a pharmacokinetic/pharmacodynamic relationship for the substrate endpoint(s) of interest is not established or when pharmacodynamic changes do not result solely from pharmacokinetic interactions (e.g., additive cardiovascular effect of quinidine and tricyclic antidepressants). When an approved drug is studied as a substrate, the impact of a given change in blood level (C_{max} , AUC) caused by an investigational interaction should be known from other interaction studies about the approved drug, although this may not always be the case for older drugs.

G. Sample Size and Statistical Considerations

For both investigational drugs and approved drugs, when used as substrates and/or interacting drugs in drug-drug interaction studies, the general objective of the analysis should be either to state that a certain type of drug-drug interaction is or is not likely and, if likely, an expected clinical impact. These statements should be based on the use of pharmacokinetic measures most relevant to understanding the relationship between dose (exposure) and therapeutic outcome. A confidence interval approach, as opposed to a significance test, is preferred, since metabolic drug-drug interactions are comparative in nature. Selection of the number of subjects for a given study will thus depend on several factors, including the criterion used for purposes of comparison (e.g., comparison of log-transformed pharmacokinetic measures/parameters), the width of the equivalence interval beyond which an adjustment in dose is considered important, and the confidence interval

DRAFT — NOT FOR IMPLEMENTATION

(usually set at 90%). For both investigational drugs and approved drugs used as substrates, this width will depend on an understanding of dose response and concentration response relationships in the population, subpopulations, and the individual. In general, the sponsor can recommend, prior to the conduct of the studies, specific *no effect* boundaries based on information available for both the investigational agent and the approved drugs used in the studies. In the absence of other information to determine a different equivalence interval, a standard interval of 80-125% can be employed for both the investigational drug and the approved drugs used in the interaction studies. For example, if the 90% confidence interval of the ratio of the pharmacokinetic measures/parameters with and without the interacting drug ((S+I)/S) falls completely within the 80- 125% default interval, the conclusion can be reached that no metabolic drug-drug interaction was present. This approach applies for both the investigational drug as substrate as well as the approved drug as substrate.

V. LABELING

A. Drug Metabolism

All information on the metabolic pathways and metabolizes and pharmacokinetic interaction should be included in the CLINICAL PHARMACOLOGY section of the labeling. The consequences of metabolism and interactions should be placed in PRECAUTIONS/WARNINGS, CONTRAINDICATIONS, and DOSAGE AND ADMINISTRATION sections, as appropriate.

B. Metabolic Drug-Drug Interaction Studies

Relevant in vitro and in vivo metabolic drug-drug interaction data describing the drug's effects on substrates and the effects of inhibitors and inducers on the drug should be presented in the DRUG-DRUG INTERACTIONS section of the labeling in the CLINICAL PHARMACOLOGY section, including both positive and important negative findings. The types of studies on which statements are based should be identified *briefly* in the labeling. If findings indicate a known or potential interaction of clinical significance or lack of an important interaction that might have been expected, these should be mentioned briefly in the clinical pharmacology interactions section and described more fully in the interaction section under PRECAUTIONS, with advice on how to adjust treatment placed in WARNINGS/PRECAUTIONS, DOSAGE AND ADMINISTRATION, and CONTRAINDICATIONS, as appropriate. In certain cases, information based on clinical studies not using the labeled drug under investigation can be described with an explanation that similar results maybe expected for the labeled

DRAFT — NOT FOR IMPLEMENTATION

drug. For example, a strong inhibitor of CYP3A4 does not need to be tested with all 3A4 substrates to warn against an interaction. Examples of appropriate labeling language are provided in italics below.

DRUG-DRUG INTERACTIONS, CLINICAL PHARMACOLOGY

- In vivo metabolic drug-drug interaction studies indicate little or no pharmacokinetic effect:

Data from a drug-drug interaction study involving @@ and (simvastatin) in _____ patients/healthy individuals indicate that the PK disposition of (simvastatin) is not altered when the drugs are co-administered. This indicates that (drug) does not inhibit CYP3A4 and will not alter the metabolism of drugs metabolized by this enzyme.

- In vivo metabolic drug-drug interaction studies indicate a clinically significant pharmacokinetic interaction:

The effect of (drug) on the pharmacokinetics of (simvastatin) has been studied in _____ patients/healthy subjects. The C_{max}, A_{UC}, half-life and clearances of (simvastatin) increased/decreased by _____% (90% Confidence Interval: to _____%) in the presence of (drug). This indicates that (drug) can inhibit the metabolism of drugs metabolized by CYP3A4 and can increase blood levels of such drugs. (See PRECAUTIONS, WARNINGS, DOSAGE AND ADMINISTRATION, or CONTRAINDICATIONS sections.)

Specific enzymes have been identified as metabolizing the test drug, but no in vivo or in vitro drug interaction studies have been conducted:

In vitro drug metabolism studies reveal that (drug) is a substrate of the CYP _____ enzyme. No in vitro or clinical drug interaction studies have been performed to evaluate interactions. However, based on the in vitro data, blood levels of (drug) are expected to increase in the presence of inhibitors of _____ such as _____, _____, or _____

- Neither in vivo nor in vitro drug-drug interaction studies have been conducted and there is no significant metabolism of the drug:

In vivo or in vitro drug-drug interaction studies have not been conducted. The drug interaction potential resulting in changes of PK of (drug) is expected to be low because approximately 90% of the recovered dose of _____ is excreted in the urine as unchanged drug. However, whether (drug) can inhibit or induce metabolic enzymes is not known.

DRAFT — NOT FOR IMPLEMENTATION

- In vitro interaction has been studied but no in vivo studies have been conducted to confirm or refute a finding:

In vitro interaction demonstrated:

In vitro drug interaction studies reveal that the metabolism of (drug) is by CYP 3A4 and can be inhibited by the CYP3A4 inhibitor ketoconazole. No clinical studies have been performed to evaluate this finding. Based on the in vitro findings, it is likely that ketoconazole, grapefruit juice, and other 3A4 inhibitors will lead to substantial increase of (drug) blood levels. Refer to PRECAUTIONS, as appropriate.

In vitro interaction not demonstrated:

In vitro drug interaction studies reveal no inhibition of the metabolism of (drug) by the CYP 3A4 inhibitor ketoconazole. No clinical studies have been performed to evaluate this finding. However, based on the in vitro findings, a metabolic interaction with ketoconazole, grapefruit juice, and other 3A4 inhibitors is not anticipated. Refer to PRECAUTIONS, as appropriate.

PRECAUTIONS and/or WARNINGS

- An interacting drug causes increased levels of the substrate but the need for both drugs may continue with appropriate dosage adjustment. Results of the studies are described in CLINICAL PHARMACOLOGY, DRUG-DRUG INTERACTIONS, PRECAUTIONS and/or WARNINGS and may state:

*Drug_____/class of drug causes **significant** increases in concentrations of_____ when co-administered so that dose of_____ must be adjusted (see DOSAGE AND ADMINISTRATION). If there is an important interaction, information for patients should point this out also.*

- An interacting drug causes increased risk because of increased levels of the substrate and the interacting drug should not be used with the substrate. After describing the interaction in the CLINICAL PHARMACOLOGY section, there should be a CONTRAINDICATIONS section and possibly a boxed warning if the risk is serious.

*Drug_____/class of drug can cause **significant** increases in concentrations of drug_____ when co-administered. The two drugs should not be used together.*

DRAFT — NOT FOR IMPLEMENTATION

DOSAGE AND ADMINISTRATION

- An interacting drug causes increased risk because of increased levels of the substrate but the need for both drugs may continue with suitable monitoring:

*Drug_____/class of drug leads to **significant** increases in blood levels of_____ by _____%. The dose of_____ should be decreased by _____% when the patient is also taking _____*

CONTRAINDICATIONS

- An interacting drug causes increased risk because of increased levels of the substrate and should be discontinued:

Drug_____/class of drug leads to significant increases in blood levels of_____, with potentially serious adverse events. Administration of_____ to patients on drug_____/class of drug is contraindicated.

DRAFT — NOT FOR IMPLEMENTATION

COMIS # 2635

Drafted: Shiew-Mei Huang

Reviewed: SO'Malley

Proofed: NDerr: 10/19/98

SO'Malley
10/29/98