

PATIENT PRESCRIPTION INFORMATION

05-16-2001

PHIMS. U

027A

ATENOLOL 50MG TABLET MYL
MYLAN
TAKE 1 TABLET EVERY DAY

IF YOU HAVE ANY QUESTIONS ABOUT YOUR MEDICATION, PLEASE CONTACT YOUR PHARMACIST:

BETA BLOCKERS - ORAL

WARNING: If you have angina or have had heart problems, do not suddenly stop using this medication without first consulting your doctor. If your doctor decides you should no longer use this medication, you must stop this medication gradually according to your doctor's instructions.

USES: This medication is used for chest pain (angina), high blood pressure and irregular heartbeats.

HOW TO TAKE THIS MEDICATION: Take this medication exactly as prescribed. Try to take it at the same time(s) each day. Do not suddenly stop taking this medication without consulting your doctor. Some conditions may become worse when the drug is suddenly stopped.

SIDE EFFECTS: You may experience dizziness, lightheadedness, drowsiness, and blurred vision as your body adjusts to the medication. Use caution engaging in activities requiring alertness. Because beta-blockers reduce blood circulation to the extremities, your hands and feet may be more susceptible to the cold. Dress warm. Inform your doctor if you develop: easy bruising or bleeding, swollen hands or feet, confusion, depression, a sore throat. In the unlikely event you have an allergic reaction to this drug, seek medical attention immediately. Symptoms of an allergic reaction include: rash, itching, swelling, dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS: Before taking this drug, tell your doctor if you have a history of: heart disease, kidney disease, liver disease, asthma, bronchitis, emphysema, any other lung disease, diabetes, overactive thyroid gland, any drug allergies. Before having surgery, tell the doctor or dentist in charge that you are taking a beta-blocker. This drug should be used only if clearly needed during pregnancy. Discuss the risks and benefits with your doctor. This drug is excreted into breast milk and though no problems have been noted, the possibility for adverse effects to the infant exists. Consult your doctor before breast-feeding.

DRUG INTERACTIONS: Tell your doctor of all prescription and nonprescription drugs you may use, especially of: diuretics, cold preparations and nasal decongestants, reserpine, other heart or high blood pressure medications. Do not start or stop any medicine without doctor or pharmacist approval.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include usually fast or slow heartbeat, dizziness, slow or shallow breathing, seizures, unconsciousness, weakness, or fatigue.

NOTES: Your doctor may want you to take your pulse each day while you take this medication. Learn how to monitor your pulse.

MISSED DOSE: If you miss a dose, take as soon as remembered but not if it is within 4 hours of next dose (8 hours for penbutolol, atenolol, nadolol). If it is, skip the missed dose and resume your usual dosing schedule. Do not "double-up" the dose to catch up.

STORAGE: Store at room temperature between 59 and 86 degrees F (between 15 and 30 degrees C) away from moisture and sunlight. Do not store in the bathroom.



For faster refills, phone in 24 hours in advance

Keep Out of Reach of Childrer

047 A

QUESTIONS? ASK YOUR

PHARMACIST.

RX:

ATENOLOL 50MG TABLET

Directions: TAKE 1 TABLET EVERY DAY

IMPORTANT NOTE: THE FOLLOWING INFORMATION IS INTENDED TO SUPPLEMENT, NOT SUBSTITUTE FOR, THE EXPERTISE AND JUDGMENT OF YOUR PHYSICIAN, PHARMACIST OR OTHER HEALTHCARE PROFESSIONAL.

IT SHOULD NOT BE CONSTRUED TO INDICATE THAT USE OF THE DRUG IS SAFE, APPROPRIATE, OR EFFECTIVE FOR YOU.

CONSULT YOUR HEALTHCARE PROFESSIONAL BEFORE USING THIS DRUG.

BETA BLOCKERS - ORAL

USES: This medication is used for chest pain (angina), high blood pressure and irregular heartbeats.

HOW TO TAKE THIS MEDICATION: Take this medication exactly as prescribed. Try to take it at the same time(s) each day. Do not suddenly stop taking this medication without consulting your doctor. Some conditions may become worse when the drug is suddenly stopped.

SIDE EFFECTS: You may experience dizziness, lightheadedness, drowsiness, and blurred vision as your body adjusts to the medication. Use caution engaging in activities requiring alertness. Because beta-blockers reduce blood circulation to the extremities, your hands and feet may be more susceptible to the cold. Dress warm.

Inform your doctor if you develop: easy bruising or bleeding, swollen hands or feet, confusion, depression, a sore throat. In the unlikely event you have an allergic reaction to this drug, seek medical attention immediately. Symptoms of an

allergic reaction include: rash, itching, swelling, dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

Prescription Refill: Visit our online pharmacy at

Refills by Phone™: Call the number on your prescription bottle and follow the prompts.

SHEET 1 of 1

ORDER NO.

PRECAUTIONS: Before taking this drug, tell your doctor if

you have a history of: heart disease, kidney disease, liver disease, asthma, bronchitis, emphysema, any other lung disease, diabetes, overactive thyroid gland, any drug allergies. Before having surgery, tell the doctor or dentist in charge that you are taking a beta-blocker. This drug should be used only if clearly needed during pregnancy. Discuss the risks and benefits with your doctor. This drug is excreted into breast milk and though no problems have been noted, the possibility for adverse effects to the infant exists. Consult your doctor before breast-feeding.

DRUG INTERACTIONS: Tell your doctor of all prescription and nonprescription drugs you may use, especially of: diuretics, cold preparations and nasal decongestants, reserpine, other heart or high blood pressure medications. Do not start or stop any medicine without doctor or pharmacist approval.

NOTES: Your doctor may want you to take your pulse each day while you take this medication. Learn how to monitor your pulse.

MISSED DOSE: If you miss a dose, take as soon as remembered but not if it is within 4 hours of next dose (8 hours for penbutolol, atenolol, nadolol). If it is, skip the missed dose and resume your usual dosing schedule. Do not "double-up" the dose to catch up.

STORAGE: Store at room temperature between 59 and 86 degrees F (between 15 and 30 degrees C) away from moisture and sunlight. Do not store in the bathroom.

The information in this leaflet may be used as an educational aid. This information does not cover all possible uses, actions, precautions, side effects, or interactions of this medicine. This information is not intended as medical advice for individual problems.

Patient:

Medication: **ATENOLOL 50MG TABLET**

128A

USES:

This medication is used for chest pain (angina), high blood pressure and irregular heartbeats.

HOW TO TAKE THIS MEDICATION:

Take this medication exactly as prescribed. Try to take it at the same time(s) each day. Do not suddenly stop taking this medication without consulting your doctor. Some conditions may become worse when the drug is suddenly stopped.

SIDE EFFECTS:

You may experience dizziness, lightheadedness, drowsiness, and blurred vision as your body adjusts to the medication. Use caution engaging in activities requiring alertness. Because beta-blockers reduce blood circulation to the extremities, your hands and feet may be more susceptible to the cold. Dress warm. Inform your doctor if you develop easy bruising or bleeding, swollen hands or feet, confusion, depression, a sore throat. In the unlikely event you have an allergic reaction to this drug, seek medical attention immediately. Symptoms of an allergic reaction include: rash, itching, swelling, dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS:

Before taking this drug, tell your doctor if you have a history of: heart disease, kidney disease, liver disease, asthma, bronchitis, emphysema, any other lung disease, diabetes, overactive thyroid gland, any drug allergies. Before having surgery, tell the doctor or dentist in charge that you are taking a beta-blocker. This drug should be used only if clearly needed during pregnancy. Discuss the risks and benefits with your doctor. This drug is excreted into breast milk and though no problems have been noted, the possibility for adverse effects to the infant exists. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Tell your doctor of all prescription and nonprescription drugs you may use, especially of: diuretics, cold preparations and nasal decongestants, reserpine, other heart or high blood pressure medications. Do not start or stop any medicine without doctor or pharmacist approval.

MISSED DOSE:

If you miss a dose, take as soon as remembered but not if it is within 4 hours of next dose (8 hours for penbutolol, atenolol, nadolol). If it is, skip the missed dose and resume your usual dosing schedule. Do not "double-up" the dose to catch up.

STORAGE:

Store at room temperature between 59 and 86 degrees F (between 15 and 30 degrees C) away from moisture and sunlight. Do not store in the bathroom.

This information is an educational service and does not address all possible uses, actions, precautions, interactions, or side effects of this medicine. Consult your pharmacist or physician with a list of medications you are currently taking if you have any questions about adverse drug reactions.

Pharmacy Pharmacy

Directions: Take 1 tablet every day

Drug: ATENOLOL 50MG TAB
Rx#: Date: 03/28/01

002A

COMMON NAME: ATENOLOL (a-TEN-oh-lole)

COMMON USES: This medicine is a beta blocker used to treat high blood pressure and angina pectoris (chest pain). It is also used after a heart attack. This medicine may also be used to treat other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking mibefradil. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking amiodarone, disopyramide, flecainide, indomethacin, quinidine, or verapamil. Inform your doctor of any other medical conditions, allergies, pregnancy, or breast-feeding. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have a history of heart block. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine may be taken on an empty stomach or with food. **STORE THIS MEDICINE** at room temperature, away from heat and light. **IF YOU MISS A DOSE OF THIS MEDICINE**, take it if you remember the same day. If you do not remember until the next day, skip the missed dose. Do not take 2 doses at once.

CAUTIONS: **DO NOT STOP TAKING THIS MEDICINE** without checking with your doctor. **BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY**, tell the doctor or dentist that you are using this medicine. **THIS MEDICINE MAY CAUSE** drowsiness or dizziness. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. **IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK** your blood pressure and heart rate regularly, be sure to do so. **BEFORE TAKING ANY NEW MEDICINE**, either prescription or over-the-counter, check with your doctor or pharmacist. This includes any medicines used to treat colds or congestion. **FOR WOMEN: THIS MEDICINE HAS BEEN SHOWN TO CAUSE HARM** to the human fetus. **IF YOU PLAN ON BECOMING PREGNANT**, discuss with your doctor the benefits and risks of using this medicine during pregnancy. **THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING** while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby. **FOR PATIENTS WITH DIABETES MELLITUS:** this medicine may mask signs of low blood sugar such as a rapid heart rate. Check blood glucose levels regularly.

POSSIBLE SIDE EFFECTS: **SIDE EFFECTS**, that may go away during treatment, include mild drowsiness; lightheadedness or dizziness; or unusual tiredness or weakness. If they continue or are bothersome, check with your doctor. **CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE** if you experience difficulty breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include slowed heart rate, dizziness, weakness, and difficulty breathing.

ADDITIONAL INFORMATION: **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME**, obtain refills before your supply runs out.

The information in this monograph is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This information is generalized and is not intended as specific medical advice. If you have questions about the medicines you are taking or would like more information, check with your doctor, pharmacist, or nurse.

Rx#: _____
Date: 04/14/2001

If you have an
contact _____

**IF YOU HAVE CONCERNS ABOUT TAKING THE
MEDICATION BELOW. PLEASE CONTACT YOUR
PHARMACIST OR PHYSICIAN IMMEDIATELY.**

Directions: TAKE ONE TABLET BY MOUTH DAILY.

ATENOLOL 50MG TAB GENE

007A

ATENOLOL (a-TEN-oh-lole)

COMMON USES: This medicine is a beta blocker used to treat high blood pressure and angina pectoris (chest pain). It is also used after a heart attack. This medicine may also be used to treat other conditions as determined by your doctor.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine may be taken on an empty stomach or with food. **STORE THIS MEDICINE** at room temperature, away from heat and light. **IF YOU MISS A DOSE OF THIS MEDICINE**, take it if you remember the same day. If you do not remember until the next day, skip the missed dose. Do not take 2 doses at once.

CAUTIONS: **DO NOT STOP TAKING THIS MEDICINE** without checking with your doctor. **BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY**, tell the doctor or dentist that you are using this medicine. **THIS MEDICINE MAY CAUSE** drowsiness or dizziness. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. **IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK** your blood pressure and heart rate regularly, be sure to do so. **BEFORE TAKING ANY NEW MEDICINE**, either prescription or over-the-counter, check with your doctor or pharmacist. This includes any medicines used to treat colds or congestion. **FOR WOMEN: THIS MEDICINE HAS BEEN SHOWN TO CAUSE HARM** to the human fetus. **IF YOU PLAN ON BECOMING PREGNANT**, discuss with your doctor the benefits and risks of using this medicine during pregnancy. **THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING** while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby. **FOR PATIENTS WITH DIABETES MELLITUS:** this medicine may mask signs of low blood sugar such as a rapid heart rate. Check blood glucose levels regularly.

POSSIBLE SIDE EFFECTS: **SIDE EFFECTS**, that may go away during treatment, include mild drowsiness; lightheadedness or dizziness; or unusual tiredness or weakness. If they continue or are bothersome, check with your doctor. **CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE** if you experience difficulty breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist.

BEFORE USING THIS MEDICINE: Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking mibefradil. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking amiodarone, disopyramide, flecainide, indomethacin, quinidine, or verapamil. Inform your doctor of any other medical conditions, allergies, pregnancy, or breast-feeding. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have a history of heart block. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include slowed heart rate, dizziness, weakness, and difficulty breathing.

***Promptly report unusual symptoms/effects to Dr**
***Inform Dr/Dentist prior to any type of surgery.**

04/14/2001 RAO/RAO
ATENOLOL 50MG TAB GENE QTY 15

**TAKE ONE TABLET BY MOUTH
DAILY.**

ATENOLOL 50MG TAB GENE
TAKE ONE TABLET BY MOUTH ONE TIME DAILY

Prepared For:
DOB:
Refills: 3
Rx#

Prescriber:
Pharmacist:
Date: 05/29/2001

249A

GENERIC NAME: ATENOLOL (a-TEN-oh-lole)

COMMON USES: This medicine is a beta blocker used to treat high blood pressure and angina pectoris (chest pain). It is also used after a heart attack. This medicine may also be used to treat other conditions as determined by your doctor.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine may be taken on an empty stomach or with food. STORE THIS MEDICINE at room temperature, away from heat and light. IF YOU MISS A DOSE OF THIS MEDICINE, take it if you remember the same day. If you do not remember until the next day, skip the missed dose. Do not take 2 doses at once.

CAUTIONS: DO NOT STOP TAKING THIS MEDICINE without checking with your doctor. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. THIS MEDICINE MAY CAUSE drowsiness or dizziness. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK your blood pressure and heart rate regularly, be sure to do so. BEFORE TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. This includes any medicines used to treat colds or congestion. FOR WOMEN: THIS MEDICINE HAS BEEN SHOWN TO CAUSE HARM to the human fetus. IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby. FOR PATIENTS WITH DIABETES MELLITUS: this medicine may mask signs of low blood sugar such as a rapid heart rate. Check blood glucose levels regularly.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS, that may go away during treatment, include mild drowsiness; lightheadedness or dizziness; or unusual tiredness or weakness. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience difficulty breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist.

The information in this monograph is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This information is generalized and is not intended as specific medical advice. If you have questions about the medicines you are taking or would like more information, check with your doctor, pharmacist, or nurse.

PRESCRIPTION INFORMATION

GUEST:

DATE: 04/04/01

MEDICATION: ATENOLOL 50MG TAB

PRESCRIPTION:

DIRECTIONS: TAKE ONE TABLET BY MOUTH ONE TIME DAILY

176A

GENERIC NAME: ATENOLOL (a-TEN-oh-lole)

COMMON USES: This medicine is a beta blocker used to treat high blood pressure and angina pectoris (chest pain). It is also used after a heart attack. This medicine may also be used to treat other conditions as determined by your doctor.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine may be taken on an empty stomach or with food. STORE THIS MEDICINE at room temperature, away from heat and light. IF YOU MISS A DOSE OF THIS MEDICINE, take it if you remember the same day. If you do not remember until the next day, skip the missed dose. Do not take 2 doses at once.


CAUTIONS: DO NOT STOP TAKING THIS MEDICINE without checking with your doctor. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. THIS MEDICINE MAY CAUSE drowsiness or dizziness. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK your blood pressure and heart rate regularly, be sure to do so. BEFORE TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. This includes any medicines used to treat colds or congestion. FOR WOMEN: THIS MEDICINE HAS BEEN SHOWN TO CAUSE HARM to the human fetus. IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby. FOR PATIENTS WITH DIABETES MELLITUS: this medicine may mask signs of low blood sugar such as a rapid heart rate. Check blood glucose levels regularly.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS, that may go away during treatment, include mild drowsiness; lightheadedness or dizziness; or unusual tiredness or weakness. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience difficulty breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist.

The information in this monograph is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This information is generalized and is not intended as specific medical advice. If you have questions about the medicines you are taking or would like more information, check with your doctor, pharmacist, or nurse.

PATIENT
MEDICATION **ATENOLOL 50MG TABLETS**
DIRECTIONS **TAKE ONE TABLET BY MOUTH DAILY**

036A



Our records show that you have reported no allergies:

Our records show that you have reported no special health conditions:

GENERIC NAME:
ATENOLOL (a-TEN-oh-lole)

COMMON USES:
This medicine is a beta blocker used to treat high blood pressure and angina pectoris (chest pain). It is also used after a heart attack. This medicine may also be used to treat other conditions as determined by your doctor.

HOW TO USE THIS MEDICINE:
Follow the directions for using this medicine provided by your doctor. This medicine may be taken on an empty stomach or with food. **STORE THIS MEDICINE** at room temperature, away from heat and light. **IF YOU MISS A DOSE OF THIS MEDICINE**, take it if you remember the same day. If you do not remember until the next day, skip the missed dose. Do not take 2 doses at once.

CAUTIONS:
DO NOT STOP TAKING THIS MEDICINE without checking with your doctor. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. THIS MEDICINE MAY CAUSE drowsiness or dizziness. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK your blood pressure and heart rate regularly, be sure to do so. BEFORE TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. This includes any medicines used to treat colds or congestion. FOR WOMEN: THIS MEDICINE HAS BEEN SHOWN TO CAUSE HARM to the human fetus. IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby. FOR PATIENTS WITH DIABETES MELLITUS: this medicine may mask signs of low blood sugar such as a rapid heart rate. Check blood glucose levels regularly.

POSSIBLE SIDE EFFECTS:
SIDE EFFECTS, that may go away during treatment, include mild drowsiness; lightheadedness or dizziness; or unusual tiredness or weakness. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience difficulty breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist.

YOU CAN NOW USE THE INTERNET TO REQUEST REFILLS AT

KEEP OUT OF REACH OF CHILDREN - STORE IN SAFETY CONTAINERS OR SECURE AREA

Our Focus Is Pharmacy. Our Focus Is You.

Your Partner in Health Care.

3 Convenient Ways to Refill Your Prescription:

1. TOUCH-TONE: Use our automated phone service.
2. ON-LINE: Go to _____ for store pickup or direct mail.
3. AUTO PREFILL: Ask your pharmacist how your prescription can be refilled automatically.

NON-PATENT FUNDING

ATENOLOL TABLETS 50MG

RX#

05/14/01

377A

PATIENT COUNSELING MESSAGES

Follow directions. Do not stop without Dr approval
May cause drowsiness/dizziness. Drive with caution
Notify your Dr if you intend to become pregnant
Check with Dr. before taking any other medicine
Promptly report unusual symptoms/effects to Dr
Inform Dr/Dentist prior to any type of surgery.

THE INFORMATION ABOVE IS PROVIDED IN ORDER FOR YOUR MEDICATION TO BE MOST EFFECTIVE. PLEASE READ AND FOLLOW THESE INSTRUCTIONS WHILE USING THIS MEDICATION

IF YOU HAVE QUESTIONS PLEASE CALL THE PHARMACIST

PATIENT COUNSELING INFORMATION

CONSULT YOUR PHARMACIST OR PHYSICIAN WITH A LIST OF MEDICATIONS YOU ARE CURRENTLY TAKING IF YOU HAVE ANY QUESTIONS ABOUT ADVERSE DRUG REACTIONS.

156A

04/25/2001

RX#

ATENOLOL 50MG TAB*(MYL)

WHY AM I TAKING THIS DRUG?

To treat heart and/or blood pressure problems.

HOW SHOULD I TAKE IT?

Follow MD instructions. Do not miss doses and do not suddenly stop taking this without MD OK. Tell MD of other drugs you use/illnesses you have/allergies/if pregnant.

ARE THERE ANY SIDE EFFECTS

Very unlikely but report cold hands/feet, swollen hands/feet, mental changes, bruising, bleeding, weakness, trouble breathing.

HOW DO I STORE THIS?

Store at room temperature away from moisture and sunlight. Do not store in the bathroom.

IF I SHOULD MISS A DOSE?


Take missed dose as soon as remembered but not if it is within 4 hours of next dose (8 hours for penbutolol, atenolol, nadolol).

WHAT ABOUT GENERICS?

Generic alternatives are available for many drugs. They are safe, effective and can save you money. Ask your pharmacist for details.

TAKE 1 TABLET EVERY DAY

OTC DRUGS MAY AGGRAVATE CONDITION. READ LABELS. IF A WARNING SEEN, CALL DR. TAKE OR USE THIS EXACTLY AS DIRECTED. DO NOT SKIP DOSES OR DISCONTINUE.



This information is an educational service and does not address all possible uses, actions, precautions, interactions, or side effects of this medicine. If you desire any additional prescription counseling, please ask your pharmacist or your doctor. M600

PATIENT INFORMATION

RX

NDC:

038A

Prepared For:

Medication Name: ATENOLOL TAB 50MG

This medicine is used to treat hypertension (high blood pressure), angina, irregular heart beat, and other uses as determined by your doctor.

- Take this medicine at the same time each day.
- Avoid appetite suppressants and medicines for cough, colds, allergies.
- Do not interrupt or stop taking this medicine without talking to your doctor; he will slowly decrease your dose.
- Use caution when driving and performing tasks that require alertness while your body adjusts to this medicine.
- Be sure to follow your doctor's instructions on follow-up visits.

All medicine should be kept in its original containers away from children. If you need more information about this medicine, ask your doctor or pharmacist.

How to take this medicine:

Take this medicine as instructed by your doctor. It is best to take this medicine at the same time each day. Your doctor or pharmacist should be consulted before prescription or nonprescription cough, cold, allergy or appetite suppressants are taken at the same time. If you forget to take a dose, take it as soon as possible unless it is within 8 hours of the next scheduled dose. Do not double your dose unless directed by your doctor. Store this medicine at room temperature in a tightly closed container.

Side Effects and Warnings:

Side effects are possible with any medicine, but they are usually not severe enough to cause the patient to interrupt or stop taking the medicine. Dizziness, fatigue, slow heart or pulse rate, shortness of breath, diarrhea, rash or itching may occur. Call your doctor if these symptoms do not go away or become severe, or if you have any other bothersome side effects. Sudden changes in posture should be avoided. Use caution when driving and performing tasks that require alertness.

ATENOLOL TAB 50MG

Take 1 tablet(s) by mouth once daily

Follow directions. Do not stop without Dr approval
May cause drowsiness/dizziness. Drive with caution
Notify your Dr if you intend to become pregnant
Check with Dr. before taking any other medicine
Promptly report unusual symptoms/effects to Dr
Inform Dr/Dentist prior to any type of surgery.

A Message From your Pharmacist

ATENOLOL 50MG

Directions:

TAKE ONE (1) TABLET EACH DAY

Information:

- Take with or without food.
- Report rash or allergic reaction.
- MAY CAUSE DIZZINESS

042A

TRY OUR NEW FLAVORS
GOLDEN FARM CANDIES
BEST SUGAR FREE CANDY EVER

Rx

05/09/01

~~101~~
044A

ATENOLOL 50MG.-IPR
Best to take at the same
time daily. Do not double
dose if missed. Do not use
more or less often than
doctor said. Do not skip
any doses. May make you
sleepy; use caution
driving. Do not stop
medicine without calling
doctor. Store at room
temperature away from
sunlight. Tell doctor your

230

- ✓ Best to take at the same time each day to maintain blood levels
- ✓ Do not double the next dose if a dose is missed
- ✓ Use exactly as directed by doctor
- ✓ Must be compliant with therapy

- ✓ May cause drowsiness; Caution operating machinery or driving
- ✓ Check with doctor before discontinuing
- ✓ Store at room temperature away from heat and sunlight
- ✓ Physician must know patient's complete medical history

ATENOLOL 50MG GENEVA PHA
CASH/CP
Your pharmacist is:

04/12/2001
00 Rf16
15 TABS
Price: 6.99

PATIENT COUNSELING

For: ATENOLOL

50MG

MAY CAUSE DROWSINESS
TAKE WITH OR WITHOUT FOOD

083A

323A

04/23/01

ATENOLOL 50MG TABLET

..TAKE W/AFTER FOOD OR MILK
..DO NOT STOP MED. ABRUPTLY

┌ ..PLEASE REMEMBER-SOME DR.*
└ ..**OFFICES REQUIRE 24 HRS*
 ..*** NOTICE ON REFILLS ***

313A

RX # *****
04/18/01

ATENOLOL 50MG TABLET

..DO NOT STOP MED. ABRUPTLY
..DO NOT STOP MED. ABRUPTLY

..
..

398A

***** PATIENT INFORMATION LEAFLET *****

PATIENT:) DATE: April 10, 2001

L

RX # RFLS LEFT: 0 QTY: 15 RPH:
DR. PHONE#

INSTRUCTIONS: TAKE 1 TABLET BY MOUTH ONCE DAILY

DRUG NAME: ATENOLOL 50MG TAB
GENERIC NAME: ATENOLOL (a-TEN-oh-lole)

COMMON USES: This medicine is a beta blocker used to treat high blood pressure and angina pectoris (chest pain). It is also used after a heart attack. This medicine may also be used to treat other conditions as determined by your doctor.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine may be taken on an empty stomach or with food. STORE THIS MEDICINE at room temperature, away from heat and light. IF YOU MISS A DOSE OF THIS MEDICINE, take it if you remember the same day. If you do not remember until the next day, skip the missed dose. Do not take 2 doses at once.

CAUTIONS: DO NOT STOP TAKING THIS MEDICINE without checking with your doctor. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. THIS MEDICINE MAY CAUSE drowsiness or dizziness. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. IF YOUR DOCTOR HAS INSTRUCTED YOU TO CHECK your blood pressure and heart rate regularly, be sure to do so. BEFORE TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. This includes any medicines used to treat colds or congestion. FOR WOMEN: THIS MEDICINE HAS BEEN SHOWN TO CAUSE HARM to the human fetus. IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby. FOR PATIENTS WITH DIABETES MELLITUS: this medicine may mask signs of low blood sugar such as a rapid heart rate. Check blood glucose levels regularly.

398A

***** PATIENT INFORMATION LEAFLET *****

PATIENT: .

DATE: April 10, 2001

RX #

RFLS LEFT: 0

QTY: 15 RPH:
DR. PHONE#

INSTRUCTIONS: TAKE 1 TABLET BY MOUTH ONCE DAILY

DRUG NAME: ATENOLOL 50MG TAB

POSSIBLE SIDE EFFECTS: SIDE EFFECTS, that may go away during treatment, include mild drowsiness; lightheadedness or dizziness; or unusual tiredness or weakness. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience difficulty breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist.

The information in this monograph is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This information is generalized and is not intended as specific medical advice. If you have questions about the medicines you are taking or would like more information, check with your doctor, pharmacist, or nurse.