MUTAGENICITY TEST ON

COMBINED ETHANOL EXTRACTS OF FIVE MAMMARY IMPLANT COMPONENTS

IN THE CHO/HGPRT FORWARD MUTATION ASSAY

FINAL REPORT

VII. CONCLUSIONS

The test material, combined ethanol extracts of five mammary implant components, is considered negative for inducing forward mutations at the HGPRT locus in Chinese hamster ovary cells under both the S9 metabolic activation and nonactivation conditions of the assay.

VIII. REFERENCES

1. Hsie, A.W., Briner, P.A., Mitchell, T.J., and Gosslee, D.G.: Dose-response relationship for ethyl methanesulfonate-induced mutations at the hypoxanthine-guanine phosphoribosyl transferase locus in Chinese hamster ovary cells. Somat. Cell Genet., 1(3):247-261, 1975.

2. Hsie, A.W., Casciano, D.A., Couch, D.B., Krahn, D.F., O'Neill, J.P., and Whitfield, B.L.: The Use of Chinese Hamster Ovary Cells to Quantify Specific Locus Mutation and to Determine Mutagenicity of Chemicals. Mutation Res., 86:193-214, 1981.

3. Kao, F.T. and Puck, T.T.: Genetics of Somatic Mammalian Cells, VII. Induction and Isolation of Nutritional Mutants in Chinese Hamster Cells. Proc. Natl. Acad. Sci. U.S., 60(4):1275-1281, 1968.

4. Kastenbaum, M.A. and Bowman, K.O.: Tables for determining the statistical significance of mutation frequencies. Mutation Res., 9:527-549, 1970.

5. Myhr, B.C. and DiPaolo, J.A.: Mutagenesis of Chinese hamster cells *in vitro* by combination treatments with methyl methanesulfonate and N-acetoxy-2-acetylaminofluorene. Cancer Res., 38:2539-2543, 1978.