Dosage and Administration

General: Each dose should be taken with a full glass of water unless contraindicated. Doses may need to be individualized depending on indication.

Indications	Recommended Daily Dose	Duration of Therapy
Vascular Indications:		
Ischemic Strokes and TIA	50-325 milligrams (mg) daily	Indefinitely
Suspected Acute MI	160-162.5 mg taken as soon as infarction is suspected; then once daily	For 30 days post infarction (after 30 days consider further treatment based on indication for previous MI)
Prevention of Recurrent MI	75-325 mg daily	Indefinitely
Unstable Angina Pectoris	75-325 mg daily	Indefinitely
Chronic Stable Angina Pectoris	75-325 mg daily	Indefinitely
Revascularization Procedures in Select Patients:		
CABG	325 mg daily starting 6 hrs. postprocedure	1 year
PTCA	325 mg 2 hours presurgery Maintenance therapy: 160-325 mg daily	Indefinitely
Carotid Endarterectomy	80 mg daily to 650 mg twice a day started presurgery	Indefinitely
Rheumatologic Disease Indications:		
Rheumatoid Arthritis	Initial dose 3 g daily. Target plasma salicylate levels 150-300 micrograms/milliliter (μg/mL)	As indicated
Juvenile Rheumatoid Arthritis	Initial dose 90-130 mg/kilograms/day. Target plasma salicylate levels 150-300 μg/mL	As indicated
Spondyloarthropathies	Up to 4 grams (g) daily	As indicated
Osteoarthritis	Up to 3 g daily	As indicated
Arthritis and Pleurisy of SLE	Initial dose 3 g daily. Target plasma salicylate levels 150-300 μg/mL	As indicated