

**ATTACHMENT 5**  
**DRAFT GENERIC LABELING**

00040

Distributor NDC # to be printed here

Distributor NDC # used for barcode #

NDC XXXXX-XXX-XX	<b>CIII</b>	<b>PHARMACIST:</b> Dispense in a tight, light-resistant container with a child-resistant closure.	Manufactured for: XXXXXX
<b>Hydrocodone Bitartrate and Acetaminophen Oral Solution</b>	<b>USUAL DOSAGE:</b> See package insert for complete dosage recommendations.	Manufactured by: XXXXXX	 non-varnish area
<b>5 mg/325 mg per 5 mL</b>	Keep this and all medication out of reach of the children.	PSLXXXXXX ISS. XX/XX	
This product contains: Per 5 mL Hydrocodone Bitartrate, USP...5 mg Acetaminophen, USP.....325 mg Alcohol.....7 %	<b>Store at controlled room temperature, 15°-30°C (59°-86°F).</b>		
<b>1 Pint (473 mL)</b>	<b>Rx only</b>		

Manufacturer and Distributor Name and their locations to be printed here along with the actual PSL # and ISS date

Lot #: Expiration Date: (To be printed here)

**HYDROCODONE BITARTRATE AND ACETAMINOPHEN  
ORAL SOLUTION**

**5 mg/325 mg per 5 mL CIII**

R<sub>x</sub> only


ISS XX/XX

PSLXXXXXX

**Description**

Hydrocodone Bitartrate and Acetaminophen Oral Solution is supplied in liquid form for oral administration.


Hydrocodone bitartrate is an opioid analgesic and antitussive and occurs as fine, white crystals or as a crystalline powder. It is affected by light. The chemical name is 4,5a-Epoxy-3-methoxy-17-methylmorphinan-6-one tartrate (1:1) hydrate (2:5). It has the following structural formula:



$C_{18}H_{21}NO_3 \cdot C_4H_6O_6 \cdot 2\frac{1}{2}H_2O$

M.W. = 494.49

Acetaminophen, 4'-Hydroxyacetanilide, a slightly bitter, white, odorless, crystalline powder, is a non-opiate, non-salicylate analgesic and antipyretic. It has the following structural formula:



$C_8H_9NO_2$

M.W. 151.16

$C_8H_9NO_2$

M.W. =151.16

This product contains:

	Per 5 mL
Hydrocodone Bitartrate, USP.....	5 mg
Acetaminophen, USP.....	325 mg
Alcohol.....	7 %

In addition, the liquid contains the following inactive ingredients: citric acid anhydrous, ethyl maltol, glycerin, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, sorbital solution, sucrose, with D&C Yellow #10 and FD&C Yellow #6 as coloring and natural and artificial flavoring.

00042

## CLINICAL PHARMACOLOGY

Hydrocodone is a semisynthetic narcotic analgesic and antitussive with multiple actions qualitatively similar to those of codeine. Most of these involve the central nervous system and smooth muscle. The precise mechanism of actions of hydrocodone and other opiates is not known, although it is believed to relate to the existence of opiate receptors in the central nervous system. In addition to analgesia, narcotics may produce drowsiness, changes in mood and mental clouding.

The analgesic action of acetaminophen involves peripheral influences, but the specific mechanism is as yet undetermined. Antipyretic activity is mediated through hypothalamic heat regulating centers. Acetaminophen inhibits prostaglandin synthetase. Therapeutic doses of acetaminophen have negligible effects on the cardiovascular or respiratory systems; however, toxic doses may cause circulatory failure and rapid, shallow breathing.

### Pharmacokinetics:

The behavior of the individual components is described below.

Hydrocodone: Following a 10 mg oral dose of hydrocodone administered to five adult male subjects, the mean peak concentration was  $23.6 \pm 5.2$  ng/mL. Maximum serum levels were achieved at  $1.3 \pm 0.3$  hours and the half-life was determined to be  $3.8 \pm 0.3$  hours. Hydrocodone exhibits a complex pattern of metabolism including O-demethylation, N-demethylation and 6-ketoreduction to the corresponding 6- $\alpha$  and 6- $\beta$  hydroxymetabolites. See **OVERDOSAGE** for toxicity information.

Acetaminophen: Acetaminophen is rapidly absorbed from the gastrointestinal tract and is distributed throughout most body tissues. The plasma half-life is 1.25 to 3 hours, but may be increased by liver damage and following overdosage. Elimination of acetaminophen is principally by liver metabolism (conjugation) and subsequent renal excretion of metabolites. Approximately 85% of an oral dose appears in the urine within 24 hours of administration, most as the glucuronide conjugate, with small amounts of their conjugates and unchanged drug. See **OVERDOSAGE** for toxicity information.

## INDICATION AND USAGE

Hydrocodone Bitartrate and Acetaminophen Oral Solution is indicated for the relief of moderate to moderately severe pain.

## CONTRADICTIONS

Hydrocodone Bitartrate and Acetaminophen Oral Solution should not be administered to patients who have previously exhibited hypersensitivity to hydrocodone and acetaminophen.

Patients known to be hypersensitive to other opioids may exhibit cross-sensitivity to hydrocodone.

## WARNINGS

**Respiratory Depression:** At high doses or in sensitive patients, hydrocodone may produce dose-related respiratory depression by acting directly on the brain stem respiratory center. Hydrocodone also affects the center that controls respiratory rhythm, and may produce irregular and periodic breathing.

**Head Injury and Increased Intracranial Pressure:** The respiratory depressant effects of narcotics and their capacity to elevate cerebrospinal fluid pressure may be markedly exaggerated in the presence of head injury, other intracranial lesions or a pre-existing increase in intracranial

pressure. Furthermore, narcotics produce adverse reactions which may obscure the clinical course of patients with head injuries.

**Acute Abdominal Conditions:** The administration of narcotics may obscure the diagnosis or clinical course of patients with acute abdominal conditions.

### **PRECAUTIONS**

**General: Special Risk Patients:** As with any narcotics analgesic agent, Hydrocodone Bitartrate and Acetaminophen Oral Solution should be used with caution in elderly or debilitated patients, and those with severe impairment of hepatic or renal function, hypothyroidism, Addison's disease, prostatic hypertrophy or urethral stricture. The usual precautions should be observed and the possibility of respiratory depression should be kept in mind.

**Cough Reflex:** Hydrocodone suppresses the cough reflex; as with all narcotics, caution should be exercised when Hydrocodone Bitartrate and Acetaminophen Oral Solution are used postoperatively and in patients with pulmonary disease.

**Information for Patients:** Hydrocodone, like all narcotics, may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery; patients should be cautioned accordingly.

Alcohol and other CNS depressants may produce an additive CNS depression, when taken with this combination product, and should be avoided.

Hydrocodone may be habit-forming. Patients should take the drug only for as long as it is prescribed, in the amounts prescribed, and no more frequently than prescribed.

**Laboratory Tests:** In patients with severe hepatic or renal disease, effects of therapy should be monitored with serial liver and/or renal function tests.

**Drug Interactions:** Patients receiving other narcotics, antihistamines, antipsychotics, antianxiety agents, or other CNS depressants (including alcohol) concomitantly with Hydrocodone Bitartrate and Acetaminophen Oral Solution may exhibit an additive CNS depression. When combined therapy is contemplated, the dose of one or both agents should be reduced.

The use of MAO inhibitors or tricyclic antidepressants with hydrocodone preparations may increase the effect of either the antidepressant or hydrocodone.

**Drug/Laboratory Test Interactions:** Acetaminophen may produce false-positive test results for urinary 5-hydroxyindoleacetic acid.

**Carcinogenesis, Mutagenesis, Impairment of Fertility:** No adequate studies have been conducted in animals to determine whether hydrocodone or acetaminophen have a potential for carcinogenesis, mutagenesis, or impairment of fertility.

**Pregnancy: Teratogenic Effects: *Pregnancy Category C:*** There are no adequate and well-controlled studies in pregnant women. Hydrocodone Bitartrate and Acetaminophen Oral Solution should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

**Nonteratogenic Effects:** Babies born to mothers who have been taking opioids regularly prior to delivery will be physically dependent. The withdrawal signs include irritability and excessive crying, tremors, hyperactive reflexes, increased respiratory rate, increased stools, sneezing, yawning, vomiting, and fever. The intensity of the syndrome does not always correlate with the

duration of maternal opioid use or dose. There is no consensus on the best method of managing withdrawal.

**Labor and Delivery:** As with all narcotics, administration of this product to the mother shortly before delivery may result in some degree of respiratory depression in the newborn, especially if higher doses are used.

**Nursing Mothers:** Acetaminophen is excreted in breast milk in small amounts, but the significance of its effects on nursing infants is not known. It is not known whether hydrocodone is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from hydrocodone and acetaminophen, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

**Pediatric Use:** Safety and effectiveness in pediatric patients have not been established.

**Geriatric Use:** Clinical studies of hydrocodone bitartrate 5 mg and acetaminophen 500 mg did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

#### **ADVERSE REACTIONS**

The most frequently reported adverse reactions are light-headedness, dizziness, sedation, nausea and vomiting. These effects seem to be more prominent in ambulatory than in nonambulatory patients, and some of these adverse reactions may be alleviated if the patient lies down.

Other adverse reactions include:

**Central Nervous System:** Drowsiness, mental clouding, lethargy, impairment of mental and physical performance, anxiety, fear, dysphoria, psychic dependence, mood changes.

**Gastrointestinal System:** Prolonged administration of Hydrocodone Bitartrate and Acetaminophen Oral Solution may produce constipation.

**Genitourinary System:** Ureteral spasm, spasm of vesical sphincters and urinary retention have been reported with opiates.

**Respiratory Depression:** Hydrocodone bitartrate may produce dose-related respiratory depression by acting directly on the brain stem respiratory centers (see **OVERDOSAGE**).

**Special Senses:** Very rare cases of hearing impairment or loss have been reported in patients predominantly receiving very high doses of hydrocodone/acetaminophen for long periods of time.

**Dermatological:** Skin rash, pruritus.

The following adverse drug events may be borne in mind as potential effects of acetaminophen: allergic reactions, rash, thrombocytopenia, agranulocytosis. Potential effects of high dosage are listed in the **OVERDOSAGE** section.

## **DRUG ABUSE AND DEPENDENCE**

**Controlled Substance:** Hydrocodone Bitartrate and Acetaminophen Oral Solution is classified as a Schedule III controlled substance.

**Abuse and Dependence:** Psychic dependence, physical dependence, and tolerance may develop upon repeated administration of narcotics; therefore, this product should be prescribed and administered with caution. However, psychic dependence is unlikely to develop when Hydrocodone Bitartrate and Acetaminophen Oral Solution is used for a short time for the treatment of pain.

Physical dependence, the condition in which continued administration of the drug is required to prevent the appearance of a withdrawal syndrome, assumes clinically significant proportions only after several weeks of continued narcotics use, although some mild degree of physical dependence may develop after a few days of narcotics therapy. Tolerance, in which increasingly large doses are required in order to produce the same degree of analgesia, is manifested initially by a shortened duration of analgesic effect, and subsequently by decreases in the intensity of analgesia. The rate of development of tolerance varies among patients.

## **OVERDOSAGE**

Following an acute overdosage, toxicity may result from hydrocodone or acetaminophen.

**Signs and Symptoms:** Hydrocodone: Serious overdose with hydrocodone is characterized by respiratory depression (a decrease in respiratory rate and/or tidal volume, Cheyne-Stokes respiration, cyanosis), extreme somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, and sometimes bradycardia and hypotension. In severe overdosage, apnea, circulatory collapse, cardiac arrest and death may occur.

Acetaminophen: In acetaminophen overdosage: dose-dependent, potentially fatal hepatic necrosis is the most serious adverse effect. Renal tubular necrosis, hypoglycemic coma, and thrombocytopenia may also occur.

Early symptoms following a potentially hepatotoxic overdose may include: nausea, vomiting, diaphoresis and general malaise. Clinical and laboratory evidence of hepatic toxicity may not be apparent until 48 to 72 hours post-ingestion.

In adults, hepatic toxicity has rarely been reported with acute overdose of less than 10 grams, or fatalities with less than 15 grams.

**Treatment:** A single or multiple overdose with hydrocodone and acetaminophen is a potentially lethal polydrug overdose, and consultation with a regional poison control center is recommended.

Immediate treatment includes support of cardiorespiratory function and measures to reduce drug absorption. Vomiting should be induced mechanically, or with syrup of ipecac, if the patient is alert (adequate pharyngeal and laryngeal reflexes). Oral activated charcoal (1 g/kg) should follow gastric emptying. The first dose should be accompanied by an appropriate cathartic. If repeated doses are used, the cathartic might be included with alternate doses as required. Hypotension is usually hypovolemic and should respond to fluids. Vasopressors and other supportive measures should be employed as indicated. A cuffed endotracheal tube should be inserted before gastric lavage of the unconscious patient and, when necessary, to provide assisted respiration.

Meticulous attention should be given to maintaining adequate pulmonary ventilation. In severe cases of intoxication, peritoneal dialysis, or preferably hemodialysis may be considered. If hypoprothrombinemia occurs due to acetaminophen overdose, vitamin K should be administered intravenously.

Naloxone, a narcotics antagonist, can reverse respiratory depression and coma associated with opioid overdose. Naloxone hydrochloride 0.4 mg to 2 mg is given parenterally. Since the duration of action of hydrocodone may exceed that of naloxone, the patient should be kept under continuous surveillance and repeated doses of the antagonist should be administered as needed to maintain adequate respiration. A narcotics antagonist should not be administered in the absence of clinically significant respiratory or cardiovascular depression.

If the dose of acetaminophen may have exceeded 140 mg/kg, acetylcysteine should be administered as early as possible. Serum acetaminophen levels should be obtained, since levels four or more hours following ingestion help predict acetaminophen toxicity. Do not await acetaminophen assay results before initiating treatment. Hepatic enzymes should be obtained initially, and repeated at 24-hour intervals.

Methemoglobinemia over 30% should be treated with methylene blue by slow intravenous administration.

The toxic dose for adults for acetaminophen is 10 grams.

#### **DOSAGE AND ADMINISTRATION**

Dosage should be adjusted according to severity of pain and response of the patient. However, it should be kept in mind that tolerance to hydrocodone can develop with continued use and that the incidence of untoward effects is dose related.

The usual adult dosage is one to two teaspoonfuls every four to six hours as needed for pain. The total daily dosage for adults should not exceed 6 teaspoonfuls.

#### **HOW SUPPLIED**

Hydrocodone Bitartrate and Acetaminophen Oral Solution is a liquid containing hydrocodone bitartrate (5 mg) and acetaminophen (325 mg) per 5 mL, with 7 % alcohol. It is supplied in containers of 1 pint (473 mL) NDC (XXXXXX-XXX-XX).

**Storage:** Store at controlled room temperature, 15°-30°C (59°-86° F).

Dispense in a tight, light-resistance container with child-resistance closure.

A Schedule CIII Narcotic.

Manufactured for  
XXXXXX

Manufactured by  
XXXXXX

ISS XX/XX

PSLXXXXXX