

EXHIBIT-8

Loratadine Syrup

Comparative Container Labeling – 4 fl oz (OTC)

Main Panel

<i>RLD - Claritin Hives Relief Fruit Flavored Syrup</i>	<i>Proposed Product – Loratadine Syrup</i>	<i>Remarks</i>
<p>SAFETY SEALED: Do not use if Schering Plough HealthCare imprinted bottle wrap torn or missing.</p> <p><i>Original Prescription Strength</i></p> <p>NEW!</p> <p style="text-align: center;"><i>Non-Drowsy *</i> Claritin® Hives Relief <i>Loratadine 5 mg/5 mL</i> <i>Antihistamine</i> Relief of Itching Due to Hives Fruit-Flavored Syrup</p> <p>24 hour * When taken as directed See Drug Facts Panel</p> <p style="text-align: right;">4 fl oz (120 mL)</p>	<p style="text-align: center;">Silarx Logo NDC 54838-538-40 <i>Original Prescription Strength</i></p> <p style="text-align: center;"><i>Non-Drowsy *</i> Loratadine Syrup Hives Relief <i>Loratadine 5 mg/5 mL</i> <i>Antihistamine</i> Relief of Itching Due to Hive Fruit Flavored Syrup</p> <p>24 hour * When taken as directed See Drug Facts Panel</p> <p style="text-align: center;">120 mL (4 FL OZ)</p>	<p>1</p> <p>2</p> <p>3</p> <p>4</p>

Remarks:

1. Moved statement of "Safety Sealed ..." on side panel
2. NDC number
3. Deleted "New"
4. Product name changed to Loratadine Syrup

Loratadine Syrup
Comparative Container Labeling – 4 fl oz (OTC)

Side Panels

<i>RLD - Claritin Hives Relief Fruit Flavored Syrup</i>	<i>Proposed Product – Loratadine Syrup</i>	<i>Remarks</i>
<p>Active ingredient (in each 5 mL) Purpose Loratadine 5 mg Antihistamine</p> <p>Use relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.</p> <p>Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. When using this product do not take more than directed. Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away See carton for full warnings.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ adults and children 6 years and over: 2 teaspoonfuls daily ■ do not take more than 2 teaspoonfuls in 24 hours <p>Other information</p> <ul style="list-style-type: none"> ■ Store between 2° and 25°C (36° and 77°F) <p>Questions or comments? 1-800-CLARITIN (1-800-252-7484) or www.claritin.com</p>	<p>Active ingredient (in each 5 mL) Purpose Loratadine 5 mg Antihistamine</p> <p>Use relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.</p> <p>Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. When using this product do not take more than directed. Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away See carton for full warnings.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ adults and children 6 years and over: 2 teaspoonfuls daily ■ do not take more than 2 teaspoonfuls in 24 hours <p>Other information</p> <ul style="list-style-type: none"> ■ SAFETY SEALED: Do not use if imprinted seal around cap of bottle is broken or missing. ■ Store between 2° and 25°C (36° and 77°F) 	<p>5</p> <p>6</p> <p>7</p>
<p>©2003 Distributed by Schering-Plough HealthCare Products, Inc. Memphis, TN 38151 USA. All right reserved. US Patent No. 6,132,758</p> <p>Lot Exp</p>	<p>Manufactured by: Silarx Pharmaceuticals, Inc. Spring Valley, NY 10977 USA ISS:01/2005</p> <p>Control # Exp. Date:</p>	

Remarks:

5. Added statement of "Safety Sealed"
6. Deleted "Questions or comments"
7. Manufacturer's name changed to Silarx, and deleted other information

Loratadine Syrup
Hives Relief

Comparative Carton Labeling

Loratadine Syrup

Comparative Carton Labeling – 4 fl oz (OTC)

Front Panel

<i>RLD - Claritin Hives Relief Fruit Flavored Syrup</i>	<i>Proposed Product – Loratadine Syrup</i>	<i>Remarks</i>
<p style="text-align: center;">NDC 0000-0000-0</p> <p style="text-align: center;"><i>Original Prescription Strength</i></p> <p>NEW!</p> <p style="text-align: center;"><i>Non-Drowsy *</i> Claritin® Hives Relief <i>Relief of Itching Due to Hives</i> <i>Loratadine 5 mg/5 mL</i> <i>Antihistamine</i> Fruit Flavored Syrup</p> <p>24 hour * When taken as directed See Drug Facts Panel</p> <p style="text-align: right;">4 fl oz 120 mL</p>	<p style="text-align: center;">Silarx Logo NDC 54838-538-40</p> <p style="text-align: center;"><i>Original Prescription Strength</i></p> <p style="text-align: center;"><i>Non-Drowsy *</i> Loratadine Syrup Hives Relief <i>Relief of Itching Due to Hives</i> <i>Loratadine 5 mg/5 mL</i> <i>Antihistamine</i> Fruit Flavored Syrup</p> <p>24 hour * When taken as directed See Drug Facts Panel</p> <p style="text-align: center;">120 mL (4 FL OZ)</p>	<p>1</p> <p>2</p> <p>3</p>

Remarks:

1. NDC number
2. Deleted "New"
3. Product name changed to Loratadine Syrup

Loratadine Syrup
Comparative Carton Labeling – 4 fl oz (OTC)

Right Side Panel

<i>RLD - Claritin Hives Relief Fruit Flavored Syrup</i>	<i>Proposed Product – Loratadine Syrup</i>	<i>Remarks</i>
Drug Facts	Drug Facts	
<i>Active ingredient (in each 5 mL) Purpose</i> Loratadine 5 mg Antihistamine	<i>Active ingredient (in each 5 mL) Purpose</i> Loratadine 5 mg Antihistamine	
Uses <ul style="list-style-type: none"> ▪ relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring. 	Uses <ul style="list-style-type: none"> ▪ relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring. 	
Warnings Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms: <ul style="list-style-type: none"> ▪ trouble swallowing ▪ dizziness or loss of consciousness ▪ swelling of tongue ▪ swelling in or around mouth ▪ trouble speaking ▪ drooling ▪ wheezing or problems breathing These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately . Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for “anaphylaxis” or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.	Warnings Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms: <ul style="list-style-type: none"> ▪ trouble swallowing ▪ dizziness or loss of consciousness ▪ swelling of tongue ▪ swelling in or around mouth ▪ trouble speaking ▪ drooling ▪ wheezing or problems breathing These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately . Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for “anaphylaxis” or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.	
Do not use <ul style="list-style-type: none"> ▪ to prevent hives from any known cause such as: <ul style="list-style-type: none"> ▪ foods ▪ insects stings ▪ medicines ▪ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause. <ul style="list-style-type: none"> ▪ if you have ever had an allergic reaction to this product (loratadine) or any of its ingredients 	Do not use <ul style="list-style-type: none"> ▪ to prevent hives from any known cause such as: <ul style="list-style-type: none"> ▪ foods ▪ insects stings ▪ medicines ▪ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause. <ul style="list-style-type: none"> ▪ if you have ever had an allergic reaction to this product (loratadine) or any of its ingredients 	

Loratadine Syrup

Comparative Carton Labeling – 4 fl oz (OTC)

Left Side Panel

<i>RLD - Claritin Hives Relief Fruit Flavored Syrup</i>	<i>Proposed Product – Loratadine Syrup</i>	<i>Remarks</i>												
Drug Facts (continued)	Drug Facts (continued)													
Ask a doctor before use if you have <ul style="list-style-type: none"> ■ liver or kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch 	Ask a doctor before use if you have <ul style="list-style-type: none"> ■ liver or kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch 													
When using this product do not take more than directed. Taking more than directed may cause drowsiness.	When using this product do not take more than directed. Taking more than directed may cause drowsiness.													
Stop use and ask a doctor if <ul style="list-style-type: none"> ■ an allergic reaction to this product occurs. Get medical help right away ■ symptoms don't improve after 3 days of treatment ■ the hives have lasted more than 6 weeks 	Stop use and ask a doctor if <ul style="list-style-type: none"> ■ an allergic reaction to this product occurs. Get medical help right away ■ symptoms don't improve after 3 days of treatment ■ the hives have lasted more than 6 weeks 													
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.													
Directions <table border="0" style="width: 100%;"> <tr> <td style="width: 50%;">adults and children 6 years and over</td> <td style="width: 50%;">2 teaspoonfuls daily; do not take more than 2 teaspoonfuls in 24 hours</td> </tr> <tr> <td>children under 6 years of age</td> <td>ask a doctor</td> </tr> <tr> <td>consumers with liver or kidney disease</td> <td>ask a doctor</td> </tr> </table>	adults and children 6 years and over	2 teaspoonfuls daily; do not take more than 2 teaspoonfuls in 24 hours	children under 6 years of age	ask a doctor	consumers with liver or kidney disease	ask a doctor	Directions <table border="0" style="width: 100%;"> <tr> <td style="width: 50%;">adults and children 6 years and over</td> <td style="width: 50%;">2 teaspoonfuls daily; do not take more than 2 teaspoonfuls in 24 hours</td> </tr> <tr> <td>children under 6 years of age</td> <td>ask a doctor</td> </tr> <tr> <td>consumers with liver or kidney disease</td> <td>ask a doctor</td> </tr> </table>	adults and children 6 years and over	2 teaspoonfuls daily; do not take more than 2 teaspoonfuls in 24 hours	children under 6 years of age	ask a doctor	consumers with liver or kidney disease	ask a doctor	
adults and children 6 years and over	2 teaspoonfuls daily; do not take more than 2 teaspoonfuls in 24 hours													
children under 6 years of age	ask a doctor													
consumers with liver or kidney disease	ask a doctor													
adults and children 6 years and over	2 teaspoonfuls daily; do not take more than 2 teaspoonfuls in 24 hours													
children under 6 years of age	ask a doctor													
consumers with liver or kidney disease	ask a doctor													
Other information <ul style="list-style-type: none"> ■ Safety sealed: do not use if bottle wrap imprinted with Schering-Plough HealthCare is open or torn ■ Store between 2° and 25°C (36° and 77°F) 	Other information <ul style="list-style-type: none"> ■ SAFETY SEALED. Do not use if imprinted seal around cap is broken or missing ■ Store between 2° and 25°C (36° and 77°F) 	4												
Inactive ingredients citric acid, edetate disodium, flavor, glycerin, propylene glycol, sodium benzoate, sugar, water.	Inactive ingredients citric acid, edetate disodium, flavor, glycerin, propylene glycol, sodium benzoate, sugar, water.													
Questions or comments? 1-800-CLARITIN (1-800-252-7484) or www.claritin.com		5												

Remarks:

4. Revised the statement of "Safety Sealed: Do not use ..."
5. Deleted questions/comments

Loratadine Syrup

Comparative Carton Labeling – 4 fl oz (OTC)

Back Panel

<i>RLD - Claritin Hives Relief Fruit Flavored Syrup</i>	<i>Proposed Product – Loratadine Syrup</i>	<i>Remarks</i>
<p>NDC 0000-0000-0</p> <p><i>Original Prescription Strength</i></p> <p>NEW!</p> <p><i>Non-Drowsy *</i></p> <p>Claritin®</p> <p>Hives Relief</p> <p><i>Relief of Itching Due to Hives</i></p> <p><i>Loratadine 5 mg/5 mL</i></p> <p><i>Antihistamine</i></p> <p>Fruit Flavored Syrup</p> <p>24 hour</p> <p>* When taken as directed See Drug Facts Panel</p> <p style="text-align: right;">4 fl oz 120 mL</p>	<p>Silarx Logo</p> <p>NDC 54838-538-40</p> <p><i>Original Prescription Strength</i></p> <p><i>Non-Drowsy *</i></p> <p>Loratadine Syrup</p> <p>Hives Relief</p> <p><i>Relief of Itching Due to Hive</i></p> <p><i>Loratadine 5 mg/5 mL</i></p> <p><i>Antihistamine</i></p> <p>Fruit Flavored Syrup</p> <p>24 hour</p> <p>* When taken as directed See Drug Facts Panel</p> <p style="text-align: center;">120 mL (4 FL OZ)</p>	<p>8</p> <p>9</p> <p>10</p>

Remarks:

- 8. NDC number
- 9. Deleted "New"
- 10. Product name changed to Loratadine Syrup