

**QUALIFIED HEALTH CLAIM PETITION**

**100% WHEY PROTEIN PARTIALLY HYDROLYZED  
in Infant Formula and  
REDUCING THE RISK OF ALLERGY IN INFANTS**

**REFERENCE LIST FOR ALL SECTIONS**

Acheson LS, Wiesner GL, Zyzanski SJ, Goodwin MA, Stange KC. Family history-taking in community family practice: implications for genetic screening. *Genet Med.* 2000;2:180-5.

American Academy of Allergy, Asthma and Immunology. Allergy statistics. Available from: [http://www.aaaai.org/media/resources/media\\_kit/allergy\\_statistics.stm](http://www.aaaai.org/media/resources/media_kit/allergy_statistics.stm). Accessed October 17, 2004.

American Academy of Allergy, Asthma and Immunology. Asthma statistics. Available from: [http://www.aaaai.org/media/resources/media\\_kit/asthma\\_statistics.stm](http://www.aaaai.org/media/resources/media_kit/asthma_statistics.stm). Accessed October 17, 2004.

American Academy of Allergy, Asthma and Immunology. Allergic conditions: Allergy. Available from: [http://www.aaaai.org/patients/allergic\\_conditions/allergy.stm](http://www.aaaai.org/patients/allergic_conditions/allergy.stm). Accessed April 21, 2005.

American Academy of Pediatrics. Committee on Nutrition. Hypoallergenic infant formulas. *Pediatrics* 2000;106:346-9.

American Academy of Pediatrics. Pediatric Nutrition Handbook, 5<sup>th</sup> edition. Editor: Kleinman RE. Food Sensitivity, pages 593-608. American Academy of Pediatrics, 2004.

Barberi I, Salpietro DC, Catalioto G, Fulia F. Clinical trial of a new hypoallergenic formula in risk infants. 2<sup>nd</sup> World Congress of Perinatal Medicine, Rome and Florence 1993 (September).

Baumgartner M, Brown CA, Exl BM, Secretin MC, van't Hof M, Haschke F. Controlled trials investigating the use of one partially hydrolyzed whey formula for dietary prevention of atopic manifestations until 60 months of age: an overview using meta-analytical techniques. *Nutr Res* 1998;18:1425-42.

Becker A, Watson W, Ferguson A, Dimich-Ward H, Chan-Yeung M. The Canadian asthma primary prevention study: outcomes at 2 years of age. *J Allergy Clin Immunol* 2004;113:650-6.

- Bender BG. Psychologic dysfunction associated with atopic dermatitis. *Immunol Allergy Clin North Am* 2002;22:43-53.
- Bergmann RL, Edenharter G, Bergmann KE, Guggenmoos-Holzmann I, Forster J, Bauer CP, Wahn V, Zepp F, Wahn U. Predictability of early atopy by cord blood-IgE and parental history. *Clin Exp Allergy* 1997;27:752-60.
- Bergmann RL, Bergmann KE, Wahn U. Can we predict atopic disease using perinatal risk factors? *Clin Exp Allergy* 1998;28:905-7.
- Boguniewicz M, Eichenfield L, Hultsch T. Current management of atopic dermatitis and interruption of the atopic march. *J Allergy Clin Immunol* 2003;112:S140-50.
- Bousquet J, Kjellman NIM. Childhood allergy. Predictive value of tests in childhood allergy. *J Allergy Clin Immunol* 1986;78:1019-22.
- The Center for Disease Control and Prevention. Asthma's impact on children and adolescents. Available from: <http://www.cdc.gov/nceh/airpollution/asthma/children.htm>
- Chamlin SL, Frieden IJ, Williams ML, Chren MM. Effects of atopic dermatitis on young American children and their families. *Pediatrics* 2004;114:607-11.
- Chan YH, Shek LPC, Aw M, Quak SH, Lee BW. Use of hypoallergenic formula in the prevention of atopic disease among Asian children. *J Paediatr Child Health* 2002;38:84-88.
- Chan-Yeung M, Manfreda J, Dimich-Ward H, Ferguson A, Watson W, Becker A. A randomized controlled study on the effectiveness of a multifaceted intervention program in the primary prevention of asthma in high-risk infants. *Arch Pediatr Adolesc Med* 2000;154:657-63.
- Chandra RK, Singh G, Shridhara B. Effect of feeding whey hydrolysate, soy and conventional cow milk formulas on incidence of atopic disease in high risk infants. *Ann Allergy* 1989;63:102-6.
- Chandra RK, Hamed A. Cumulative incidence of atopic disorders in high risk infants fed whey hydrolysate, soy, and conventional cow milk formulas. *Ann Allergy* 1991;67:129-32.
- Chandra RK, Hamed A, Prasad C, Singh GK. Cumulative incidence of allergic disorders in high risk infants fed whey hydrolysate, soy and conventional cow's milk formulas. *Am J Clin Nutr* 1992;56:758 (Abstract).
- Chandra, RK. Five-year follow-up of high-risk infants with family history of allergy who were exclusively breast-fed or fed partial whey hydrolysate, soy, and conventional cow's milk formulas. *J Pediatr Gastroenterol Nutr* 1997;24:380-8.

Chandra RK. Breast feeding, hydrolysate formulas and delayed introduction of selected foods in the prevention of food hypersensitivity and allergic disease. *Nutr Res* 2002;21:125-35.

Chirico G, Gasparoni A, Ciardelli L, De Amici M, Colombo A, Rondini G. Immunogenicity and antigenicity of a partially hydrolyzed cow's milk infant formula. *Allergy* 1997;52:82-8.

D'Agata A, Betta P, Sciacca P, Morano C, Practico G, Curreri R, Quattrocchi O, Sciacca F. The role of preventing diet in newborn at risk of atopy. Follow-up results. *Med Surg Ped* 1996;18:469-72.

Dahl RE, Bernhisel-Broadbent J, Seanlon-Holdford S, Sampson HA, Lupo M. Sleep disturbances in children with atopic dermatitis. *Arch Pediatr Adolesc Med* 1995;149:856-60.

De Seta L, Siani P, Cirillo G, Di Gruttola M, Cimaduomo L, Coletta S. Prevention of allergic disease by an hypoallergenic formula: preliminary results at the 24 months follow-up. *Med Surg Ped* 1994;16:251-4.

Exl BM, Deland U, Wall M, Preysch U, Secretin MC, Shmerling DH. Zug-Frauenfeld nutritional survey ("Zuff Study"): allergen-reduced nutrition in a normal infant population and its health-related effects: results at the age of six months. *Nutr Res* 1998;18:1443-62.

Exl BM, Deland U, Secretin MC, Preysch U, Wall M, Shmerling DH. Improved general health status in an unselected infant population following an allergen-reduced dietary intervention programme: the ZUFF-STUDY-PROGRAMME. Part II: infant growth and health status to age 6 months. *Eur J Nutr* 2000;39:145-56.

Exl BM. A review of recent developments in the use of moderately hydrolyzed whey formulae in infant nutrition. *Nutr Res* 2001;21:355-79.

Fritsché R, Pahud JJ, Pecquet S, Pfeifer A. Induction of systemic immunologic tolerance to  $\beta$ -lactoglobulin by oral administration of a whey protein hydrolysate. *J Allergy Clin Immunol* 1997;100:266-73.

Fritsché R, Bonzon M. Determination of cow milk formula allergenicity in the rat model by in vitro mast cell triggering and in vivo IgE induction. *Int Arch Allergy and Appl Immunol* 1990;93:289-93.

Gowland MH. Food allergen avoidance-the patient's viewpoint. *Allergy* 2001;36:Suppl 67:117-20.

Halken S, Jacobsen HP, Host A, Holmenlund D. The effect of hypo-allergenic formulas in infants at risk of allergic disease. *Eur J Clin Nutr* 1995;49:S77-83.

Halken S, Host A. The lessons of noninterventional and interventional prospective studies on the development of atopic disease during childhood. *Allergy* 2000;55:793-802.

Halken S, Skamstrup Hansen K, Jacobsen HP, Estmann A, Engberg Faelling A, Hansen LG, Kier SR, Lassen K, Lintrup M, Mortensen S, Kaas Ibsen K, Østerballe O, Høst A. Comparison of a partially hydrolyzed infant formula with two extensively hydrolyzed formulas for allergy prevention: A prospective, randomized study. *Pediatr Allergy Immunol* 2000;11:149-61.

Han YS, Park HY, Ahn KM, Lee JS, Choi HM, Lee SI. Short-term effect of partially hydrolyzed formula on the prevention of development of atopic dermatitis in infants at high risk. *J Korean Med Sci* 2003;18:547-51.

Hartman CT, Fredericks GL, Katz ES, Brown CA. Prevalence of symptomatic formula intolerance and allergy in a general infant population. *J Allergy Clin Immunol* 1994;210 (Abstract).

Iikura Y, Akimoto K, Ebisawa M, Onda T, Akazawa A, Saito H, Kimura T, Ishizawa K, Koya N. Effect of hydrolyzed whey protein formula for babies on development of allergic symptoms during infancy. *Nestlé Nutrition Workshop Intestinal Immunology and Food Allergy*, Raven Press 1995;34:269-75.

Kemp AS. Cost of illness of atopic dermatitis in children: A societal perspective. *Pharmacoeconomics*. 2003;21:105-13.

Kjellman NI. Atopic disease in seven-year-old children. Incidence in relation to family history. *Acta Paediatr Scand* 1977;66:465-71.

Lam BCC, Yeung CY. The effect of breast milk, infant formula and hypoallergenic formula on incidence of atopic manifestation in high risk infants. 6th Annual Scientific Meeting of Hong Kong College of Physicians and Hong Kong College of Paediatricians, Hong Kong 1992 (October).

Lam BCC, Yeung CY. The effect of breast milk, infant formula and hypoallergenic formula on incidence of atopic manifestation. Internal report, Nestec SA 1992.

Lapidus CS, Honig PJ. Atopic dermatitis. *Pediatr Rev* 1994;15:327-32.

Laughter D, Istvan JA, Tofte SJ, Hanifin JM. The prevalence of atopic dermatitis in Oregon schoolchildren. *J Am Acad Dermatol* 2000;43:649-55.

Lewis-Jones MS, Finlay AY. The Children's Dermatology Life Quality Index (CDLQI): initial validation and practical use. *Br J Dermatol* 1995;132:942-9.

Marini A, Agosti M, Motta G, Mosca F. Effects of a dietary and environmental prevention programme on the incidence of allergic symptoms in high atopic risk infants: three years follow-up. *Acta Paediatr Suppl* 1996;414:1-21.

Martinez Valverde A. Study on the assessment of serum IgE and specific IgE to Alpha lactalbumin, Beta lactoglobulin, and casein in infants fed hypoallergenic formula and other forms of feeding. Thesis, Universidad de Malaga, Spain, 1993.

MedlinePlus. Medical Encyclopedia. Available from:  
<http://www.nlm.nih.gov/medlineplus/ency/article/000812.htm>. Accessed April 21, 2005.

Moore MM, Rifas-Shiman SL, Rich-Edwards JW, Kleinman KP, Camargo CA, Gold DR, Weis SC, Gillman MW. Perinatal predictors of atopic dermatitis occurring in the first six months of life. *Pediatrics* 2004;113:468-74.

Nentwich I, Michkova E, Nevoral J, Urbanek R, Szepfalusi Z. Cow's milk-specific cellular and humoral immune responses and atopy skin symptoms in infants from atopic families fed a partially (pHF) or extensively (eHF) hydrolyzed infant formula. *Allergy* 2001;56:1144-56.

Oldaeus G, Anjou K, Bjorksten B, Moran JR, Kjellman NI. Extensively and partially hydrolysed infant formulas for allergy prophylaxis. *Arch Dis Child*. 1997;77:4-10.

Osborn DA, Sinn J. Formulas containing hydrolysed protein for prevention of allergy and food intolerance in infants (Cochrane Review). In: *The Cochrane Library*, Issue 4, 2003. Chichester, UK: John Wiley & Sons, Ltd.

Porch MC, Shahane AD, Leiva LE, Elston RC, Sorensen RU. Influence of breast milk, soy or two hydrolyzed formulas on the development or allergic manifestations in infants at risk. *Nutr Res* 1998;18:1413-24.

Ram FSF, Ducharme FM, Scarlett J. Cow's milk protein avoidance and development of childhood wheeze in children with a family history of atopy (Cochrane Review) In: *The Cochrane Library*, Issue 2, 2003. Oxford: Update Software.

Roberts G, Patel N, Levi-Schaffer F, Habibi P, Lack G. Food allergy as a risk factor for life-threatening asthma in childhood: A case-controlled study. *J Allergy Clin Immunol*. 2003;112:168-74.

Schmidt E, Eden-Köhler J, Tonkaboni F, Tölle J. Alimentary allergy prevention in infants with familial increased allergic risk: The effect of different feeding regimens in the first 6 months on atopic manifestations during the first year of life. A large-scale feeding trial. Nestlé Nutrition Workshop Intestinal Immunology and Food Allergy, Raven Press 1995;34:231-48.

Schoetzau A, Gehring U, Wichmann HE. Prospective cohort studies using hydrolysed formulas for allergy prevention in atopy-prone newborns: a systematic review. *Eur J Pediatr* 2001;160:323-32.

Sicherer SH, Sampson HA. Food hypersensitivity and atopic dermatitis: Pathophysiology, epidemiology, diagnosis, and management. *J Allergy Clin Immunol* 1999;104:S114-22.

Sicherer SH, Noone SA, Munoz-Furlong A. The impact of childhood food allergy on quality of life. *Ann Allergy Asthma Immunol* 2001;87:461-4.

Silva Rey AL, Garcia G, Nogales A. Preventive effect of a partially hydrolysed cow milk infant formula: clinical and biological 24 month follow-up study. Thesis, Spain, 1996.

Spergel JM, Paller AS. Atopic dermatitis and the atopic march. *J Allergy Clin Immunol* 2003;112:S118-27.

Spergel JM. Atopic march: link to upper airways. *Curr Opin Allergy Clin Immunol* 2005;5:17-21.

Su JC, Kemp AS, Varigos GA, Nolan TM. Atopic eczema: its impact on the family and financial cost. *Arch Dis Child* 1997;76:159-62.

Szajewska H, Mrukowicz JZ, Stoinska B, Prochowska A. Extensively and partially hydrolysed preterm formulas in the prevention of allergic diseases in preterm infants: a randomized, double-blind trial. *Acta Paediatr.* 2004;93:1159-65.

Tsai Y, Chou C, Hsieh K. The effect of hypoallergenic formula on the occurrence of allergic diseases in high risk infants. *Acta Paed Sin* 1991;32:137-44.

Vandenplas Y, Deneyer M, Sacre L, Loeb H. Preliminary data on a field study with a new hypo-allergenic formula. *Eur J Pediatr* 1988;148:274-7.

Vandenplas Y, Malfroot A, Dab I. Short-term prevention of cow's milk protein allergy in infants. *Immunology & Allergy Practice* 1989;11:430-7.

Vandenplas Y, Hauser B, Van den Borre C, Sacre L, Dab I. Effect of a whey hydrolysate prophylaxis of atopic disease. *Ann Allergy* 1992;68:419-24.

Vandenplas Y. Atopy at 3 years in high-risk infants fed whey hydrolysate or conventional formula. *Lancet* 1992;339:1118. Letter to the Editor.

Vandenplas Y, Hauser B, Van den Borre C, Clybouw C, Mahler T, Hachimi-Idrissi S, Deraeve L, Malfroot A, Dab I. The long-term effect of a partial whey hydrolysate formula on the prophylaxis of atopic disease. *Eur J Pediatr* 1995;154:488-94.

Von Berg A, Koletzko S, Grübl A, Filipiak-Pittroff B, Wichmann H, Bauer CP, Reinhardt D, Berdel D, German Infant Nutritional-Intervention Study Group. The effect of hydrolyzed cow's milk formula for allergy prevention in the first year of life: The German Infant Nutritional Intervention Study, a randomized double-blind trial. *J Allergy Clin Immunol* 2003;111:533-40.

Wadonda-Kabondo N, Sterne JAC, Golding J, Kennedy CTC, Archer CB, Dunnill MGS, and the Alspac Study Team. A prospective study of the prevalence and incidence of atopic dermatitis in children aged 0-42 months. *Br J Derm* 2003;149:1023-8.

Wahn U, von Mutius E. Childhood risk factors for atopy and the importance of early intervention. *J Allergy Clin Immunol* 2001;107:567-74.

Weiss KB, Sullivan SD. Socio-economic burden of asthma, allergy, and other atopic illnesses. *Pediatr Allergy Immunol* 1994;5:7-12.

Willems R, Duchateau J, Magrez P, Denis R, Casimir G. Influence of hypoallergenic milk formula on the incidence of early allergic manifestations in infants predisposed to atopic diseases. *Ann Allergy* 1993;71:147-50.

World Allergy Organization Project Report and Guidelines. Editors: Johansson SGO, Haahtella T. *Prevention of Allergy and Allergic Asthma*. Pages XVI-XVIII. Karger Publishers, 2004.

Yoon PW, Scheuner MT, Peterson-Oehlke KL, Gwinn M, Faucett A, Khoury MJ. Can family history be used as a tool for public health and preventive medicine? *Genet Med*. 2002;4:304-10.

Zeiger RS. Use of elemental formulas: hydrolysates in allergy prevention. In: *Physiologic/Immunologic Response to Dietary Nutrients: Role of Elemental and Hydrolysate Formulas in Management of the Pediatric Patient, Report of the 107<sup>th</sup> Ross Conference on Pediatric Research*. Columbus, OH: Ross Products Division, Abbott Laboratories Inc; 1998;108:26.

Zeiger RS. Dietary aspects of food allergy prevention in infants and children. *J Pediatr Gastroenterol Nutr* 2000;39:S77-86.

Zeiger RS. Food allergen avoidance in the prevention of food allergy in infants and children. *Pediatrics* 2003;111:1662-71.