B

Drug Facts

Active Ingredients (in each powder) — Purpose

Aspirin 650 mg — Pain reliever

Sodium 33.3 mg — Pain reliever aid

Salicylamide 195 mg — Pain reliever aid

Uses - temporarily relieves mild to moderate pain and fever due to:
- headaches
- minor joint pain
- colds
- muscular aches
- temporary reduces fever

Warnings

Reye's Syndrome: Children and teenagers should not use this drug for chicken pox or flu symptoms before a doctor is asked about Reye's Syndrome, a rare but serious illness reported to be associated with Aspirin.

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Do not use if you have ever had an allergic reaction to any other pain relieve/fever reducer

Ask a doctor before use if you have
- a bleeding problem
- asthma
- diabetes
- stomach problems that last or come back such as heartburn, upset stomach or pain

Ask a doctor or pharmacist before use if you are taking a prescription drug for:
- gout
- diabetes
- gastritis
- anticoagulation (blood thinners)

When using this product the risk of Reye's syndrome, a rare but serious illness, may be reduced by not taking other products containing aspirin, ibuprofen, or alcohol, because too much aspirin may cause bleeding in stomach or intestines, dizziness, and unconsciousness, and gastrointestinal ulcers.

Stop use and ask a doctor if
- fever lasts more than 3 days
- pain gets worse or lasts more than 10 days
- new symptoms occur
- redness or swelling is present
- bleeding in stool or loss of hearing occurs

If pregnant or nursing, consult a doctor before taking this product. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- do not exceed 4 powders in 24 hours
- adults and children 12 and over: place 1 powder on tongue, every 2 - 4 hours, followed with liquid. May also prepare by dissolving powder in water or other liquid.
- children under 12: ask a doctor

Other Information
- each powder contains: potassium 32.9 mg

Inactive Ingredients: dicalcium phosphate, magnesium sulfate anhydrous, lactic acid, lactose, potassium chloride

Bad Dog Company, Inc.
Memphis, TN 38113

Title: 7.1 pt.
Headings: 7 pt.
Sub-Headings: 8 pt.
Body: 6 pt.
Leading: 6 pt.
Font: Helvetica Condensed
Bullets: 5 pt.
em's between bullets: 2

Project Number: BC-015
File Name: BC 2's-Columns-Mod-Var Run-on
Revision: 0
Date: 11.02.00
Designer: lam

CREATIVE SERVICES GRAPHICS