C

GLUE FLAP

UNFOLDED GLUE FLAP WITH CURRENT COPY

LOT & EXPIRATION

DO NOT USE IF SEALED ON ORIGINAL OR IMPROPERLY SEALED OR TAMPERED

ORIGINAL FORMULA

BC®

FAST PAIN RELIEF

PAIN RELIEVER • FEVER REDUCER

for the pain associated with the

HEADACHES

temporary relief of minor

BODY ACHES • FEVER

2 ANALGESIC POWDERS

Black Drug Company, Inc. Memphis, TN 38110

UNFOLDED GLUE FLAP WITH CURRENT COPY

FOLDED GLUE FLAP WITHOUT CURRENT COPY REPLACED BY FDA COPY

Drug Facts

Active ingredients (in each powder) — **Purpose**

Aspirin 850 mg ..... Pain reliever

Caffeine 33.3 mg ..... Pain reliever aid

Salicylamide 195 mg ..... Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:
  - headaches
  - minor arthritis pain
  - colds
  - muscular aches
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers should not use this drug for chicken pox or flu symptoms before a doctor is asked about Reye's syndrome, a rare but serious illness reported to be associated with aspirin.

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain reliever/fever reducer. Aspirin may cause a stomach bleeding.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • shock • asthma (wheezing)

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if you have

- a bleeding problem
- asthma
- ulcers
- stomach problems that last or come back such as heartburn, upset stomach or pain

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- diabetes
- anticoagulation (blood thinning)
- arthritis
- gout

When taking this product limit the use of caffeine containing drugs, foods, or drinks, because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heartbeat.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- new symptoms occur
- fever lasts more than 3 days
- redness or swelling is present
- ringing in ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed 4 powders in 24 hours
- adults and children 12 and over: place 1 powder on tongue every 3 - 4 hours, followed with liquid. May mix powder into glass of water or other liquid.
- children under 12: ask a doctor

Other information

- each powder contains: potassium 92.9 mg

Inactive ingredients disodium sulfosuccinate, fumaric acid, lactose, potassium chloride

Title:	8.1 pt.		
Headings:	8 pt.		
Sub-Headings:	6 pt.		
Body:	6 pt.	Project Number:	BC-015
Leading:	8.5 pt.	File Name:	BC 2's-Columns-Stan-Hor
Font:	Helvetica Condensed	Revision:	0
Bullets:	5 pt.	Date:	10.16.00
em's between bullets:	2	Designer:	tam

Black | PMS 186 | PMS 300

CREATIVE SERVICES GRAPHICS