

2.5"

2 3/8" (2.375)

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions	<ul style="list-style-type: none"> ■ do not take more than directed
adults and children 12 years and over	<ul style="list-style-type: none"> ■ take 2 caplets every 4-6 hours as needed ■ do not take more than 8 caplets in 24 hours
children under 12 years	<ul style="list-style-type: none"> ■ do not use this adult product in children under 12 years of age; this will provide more than the recommended dose (overdose) and could cause serious health problems.

3

2

Drug Facts (continued)

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other product containing acetaminophen

Ask a doctor before use if you have

- heart disease ■ high blood pressure
- thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland

When using this product

- do not exceed recommended dosage

Stop use and ask a doctor if

- new symptoms occur ■ redness or swelling is present
- pain gets worse or lasts for more than 7 days
- fever gets worse or lasts for more than 3 days
- you get nervous, dizzy or sleepless

OUTSIDE PAGES

Maximum Strength
TYLENOL[®]
Pain Reliever
SINUS
Nasal Decongestant

2 Caplets
NON-DROWSY

Drug Facts

Active ingredient (in each caplet)	Purpose
Acetaminophen 500 mg.....	Pain reliever
Pseudoephedrine HCl 30 mg.....	Nasal decongestant

Uses temporarily relieves:

- sinus pain
- headache
- nasal and sinus congestion

Warnings

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.

▼ Open here to view complete product information ▼

1

4

Drug Facts (continued)

Other information

- do not use if pouch is opened
- store at room temperature

Inactive ingredients

carnauba wax, cellulose, corn starch, D&C Yellow #10, FD&C Blue #1, FD&C Red #40, hydroxypropyl methylcellulose, iron oxide, magnesium stearate, polyethylene glycol, polysorbate 80, sodium starch glycolate, titanium dioxide

Questions or comments?
call toll-free 1-877-TYLENOL (1-877-895-3665)

McNeil
MCNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA
©MCN-PPC, Inc. '01
Visit us at www.tylenol.com XXXXXXX

Labeling Format Information:

Fonts: Helvetica bold, black and black italic.

Drug Facts:	8.5 pt	Leading:	6.5 pt
Header:	8 pt	Bullets:	5 pt
Subheader:	6 pt	Barlines:	2.5 pt
Body Text:	6 pt	Hairlines:	.5 pt
Drug Facts (continued):	8 pt		

Horizontal Scale: 70-75%

Average Kerning: 0