

Draft Guidance on Amoxicillin

This draft guidance, once finalized, will represent the Food and Drug Administration's (FDA's) current thinking on this topic. It does not create or confer any rights for or on any person and does not operate to bind FDA or the public. You can use an alternative approach if the approach satisfies the requirements of the applicable statutes and regulations. If you want to discuss an alternative approach, contact the Office of Generic Drugs.

Active ingredient: Amoxicillin

Form/Route: Oral Suspension/Oral

Recommended studies: 2 studies

1. Type of study: Fasting
Design: Single-dose, two-way crossover in vivo
Strength: 400 mg/ 5 mL
Subjects: Healthy males and nonpregnant females, general population.

2. Type of study: Fed
Design: Single-dose, two-way crossover in vivo
Strength: 400 mg/ 5 mL
Subjects: Healthy males and nonpregnant females, general population.

Analytes to measure (in appropriate biological fluid): Amoxicillin in plasma

Bioequivalence based on (90% CI): Amoxicillin

Waiver request of in vivo testing: 200 mg/5 mL based on (i) acceptable bioequivalence studies on the 400 mg/5 mL strength, (ii) acceptable in vitro dissolution testing of all strengths, and (iii) proportional similarity of the formulations across all strengths.

Dissolution test method and sampling times:

Please note that a **Dissolution Methods Database** is available to the public at the OGD website at <http://www.accessdata.fda.gov/scripts/cder/dissolution/>. Please find the dissolution information for this product at this website. Please conduct comparative dissolution testing on 12 dosage units each of all strengths of the test and reference products. Specifications will be determined upon review of the application.