UNPRINTED AREA FOR DATE & LOT NUMBER - UNVARNISHED

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS Title	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE AFTER BULLETED SECTION	2 sms
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

100%



200%

See box for important warnings and save for future use.

Uses temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold: runny nose, sneezing, itchy, watery eyes, itching of the nose and throat, nasal congestion, headache, fever, minor aches and pains, and sinus pressure

Warnings **Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing), shock

Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.

Do not use if the child has ever had an allergic reaction to any pain reliever/fever reducer, nasal decongestant, or antihistamine. **Stop use and ask a doctor if** an allergic reaction occurs. Seek medical help right away. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Children's
Advil®
Allergy Sinus
Suspension
Pain Reliever/Fever Reducer
Nasal Decongestant/Antihistamine

Active ingredients
(in each 5 mL teaspoon) *Purposes*
Chlorpheniramine maleate 1 mg.....Antihistamine
Ibuprofen 100 mg.....Pain reliever/Fever reducer
Pseudoephedrine HCl 15 mg.....Nasal decongestant

Berry Flavored Liquid
Alcohol Free 1 FL OZ (30 mL)

Do Not Use if breakable ring on bottle cap is separated or missing.

Directions ■ **do not give more than directed**
■ **shake well before using**
■ find right dose on chart below. If possible, use weight to dose; otherwise use age.
■ If needed, repeat dose every **6 hours** while symptoms persist
■ do not use more than **4 times a day**
■ measure only with dosing cup provided

Dosing Chart

Weight (lb)	Age (yrs)	Dose (teaspoons)
under 48	under 6	ask a doctor
48 – 95	6 – 11	2 teaspoons

Wyeth Consumer Healthcare
Madison, NJ 07940 U.S.A. U0183-10-1 FDA05

128 CODE AREA- SUBSET C 10 MIL DENSITY WITH 4 OR 6 DIGITS-UNVARNISHED

UNPRINTED AREA FOR DATE & LOT NUMBER - UNVARNISHED

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS Title	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEAD & BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE AFTER BULLETED SECTION	2 sms
• BAR LINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

3/16" X 3/16" DATA MATRIX CODE (01832001), PRINTED IN BLACK OR DARK BLUE ON UNPRINTED WHITE BACKGROUND WITH A MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES.

100%

UNVARNISHED UNPRINTED AREA FOR LOT & EXP.

150%

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE AFTER BULLETED SECTION	2 ems
• BOX LINE	.5 pt
• SMALLEST ®, TM etc.	4.5 pt

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS Title	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 sims
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

100%



150%

See box for important warnings and save for future use.

Active ingredients (in each 5 mL teaspoon) *Purposes*
Chlorphenirramine maleate 1 mg.....Antihistamine
Ibuprofen 100 mg.....Pain reliever/Fever reducer
Pseudoephedrine HCl 15 mg.....Nasal decongestant

Uses temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold: runny nose, sneezing, itchy, watery eyes, itching of the nose and throat, nasal congestion, headache, fever, minor aches and pains, and sinus pressure

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.
Do not use if the child has ever had an allergic reaction to any pain reliever/fever reducer, nasal decongestant, or antihistamine
Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Packaged with Tamper-Evident bottle cap.
Do Not Use if breakable ring is separated or missing.

Children's
Advil®
Allergy Sinus
Suspension
Pain Reliever/Fever Reducer
Nasal Decongestant/Antihistamine

Berry Flavored Liquid
Alcohol Free 8 FL OZ (240 mL)

Directions
■ do not give more than directed
■ shake well before using
■ find right dose on chart below. If possible, use weight to dose; otherwise use age.
■ If needed, repeat dose every 6 hours while symptoms persist
■ do not use more than 4 times a day
■ measure only with dosing cup provided

Dosing Chart

Weight (lb)	Age (yrs)	Dose (teaspoons)
under 48	under 6	ask a doctor
48 – 95	6 – 11	2 teaspoons

Other information
■ each 5 mL teaspoon contains: sodium 10 mg
■ store at 20-25°C (68-77°F)

Questions or comments? call weekdays from 9AM to 5PM EST at 1-800-88-ADVIL or ask your pharmacist, doctor or healthcare professional

Wyeth Consumer Healthcare,
Madison, NJ 07940 U.S.A.
www.advil.com U0183-30-1.FDA05

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE AFTER BULLETED SECTION	2 ems
• BOX LINE	.5 pt
• SMALLEST ®, TM etc.	5.75 pt