00000000

Motrin Cold & Sinus

contains Ibuprofen 200 mg/Pseudoephedrine HCl 30 mg

▶ Sinus Headache ▶ Sinus Pressure ▶ Nasal Congestion

NON-DROWSY FORMULA

Motrin Cold & Sinus

contains Ibuprofen 200 mg/ Pseudoephedrine HCl 30 mg

EXP:

Drug Facts

Active Ingredients (in each caplet) — **Purpose**

Ibuprofen 200 mg ..... Pain reliever/fever reducer

Pseudoephedrine HCl 30 mg ..... Nasal decongestant

Uses temporarily relieves these symptoms associated with the common cold, sinusitis, or flu:

■ headache ■ nasal congestion ■ fever ■ minor body aches and pains ▶

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take Ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding. ▲

XXXXXXX

Drug Facts (continued)

Do not use if you

■ have ever had an allergic reaction to any other pain relievers/fever reducer

■ are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

■ heart disease ■ high blood pressure

■ thyroid disease ■ diabetes

■ trouble urinating due to an enlarged prostate gland

■ had serious side effects from taking any pain relievers/fever reducers

Ask a doctor or pharmacist before use if you are

■ taking any other product that contains ibuprofen or pseudoephedrine

■ taking any other pain relievers/fever reducer or nasal decongestant

■ under a doctor's care for any continuing medical condition

■ taking other drugs on a regular basis

When using this product

■ **do not use more than directed**

■ give with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ an allergic reaction occurs. Seek medical help right away.

■ you get nervous, dizzy or sleepless

■ nasal congestion lasts for more than 7 days

■ fever lasts for more than 3 days ▶

Drug Facts (continued)

■ symptoms continue or get worse

■ new or unexpected symptoms occur

■ stomach pain occurs with use of this product or if even mild symptoms persist

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use this product during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years of age and older	■ take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used. ■ do not use more than 6 caplets in any 24-hour period unless directed by a doctor ■ the smallest effective dose should be used
children under 12 years of age	consult a doctor

Other Information

■ **do not use if blister unit is broken or open**

■ store at 20 - 25°C (68 - 77°F)

■ avoid excessive heat above 40°C (104°F)

■ read all warnings and directions before use. Keep carton.

■ see end panel for lot number and expiration date ▲

00000000

00000000

MCNEIL – PPC, INC., FORT WASHINGTON, PA 19034 USA ©MCN-PPC, Inc. '03 Visit us at www.motrin.com

Questions or comments? Call toll-free 1-800-962-5357

Inactive Ingredients

carnauba wax, cellulose, corn starch, FD&C Red #40, hypromellose, silicon dioxide, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide, triacetin

0 0000-0000-0 9

CTN MOTRIN COLD & SINUS CPLT 20
3004315 A-1
MAY 13, 2003

Labeling Format Information:

Fonts: Helvetica regular, bold and bold italic.

Drug Facts:	8.15 pt	**Leading:**	6 pt
Header:	7 pt	**Bullets:**	5 pt
Subheader:	6 pt	**Barlines:**	2.5 pt
Body Text:	6 pt	**Hairlines:**	0.5 pt
Drug Facts (continued):	7 pt		

Horizontal Scale: 78-80% Average Kerning: -20

Motrin
Cold & Sinus
McNeil
Div. of McNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
LOT XXXXXX EXP 00/00
See Carton for Complete Usage Information.
0001018

TEAR OR CUT AT NOTCH
TEAR OR CUT AT NOTCH

Motrin
Cold & Sinus
McNeil
Div. of McNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
LOT XXXXXX EXP 00/00
See Carton for Complete Usage Information.
0001018

TEAR OR CUT AT NOTCH
TEAR OR CUT AT NOTCH

Motrin
Cold & Sinus
McNeil
Div. of McNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
LOT XXXXXX EXP 00/00
See Carton for Complete Usage Information.
0001018

TEAR OR CUT AT NOTCH
TEAR OR CUT AT NOTCH

Motrin
Cold & Sinus
McNeil
Div. of McNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
LOT XXXXXX EXP 00/00
See Carton for Complete Usage Information.
0001018

TEAR OR CUT AT NOTCH
TEAR OR CUT AT NOTCH

Motrin
Cold & Sinus
McNeil
Div. of McNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
LOT XXXXXX EXP 00/00
See Carton for Complete Usage Information.
0001018

TEAR OR CUT AT NOTCH
TEAR OR CUT AT NOTCH

Motrin
Cold & Sinus
McNeil
Div. of McNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
LOT XXXXXX EXP 00/00
See Carton for Complete Usage Information.
0001018

Motrin
Cold & Sinus
McNeil
Div. of McNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
LOT XXXXXX EXP 00/00
See Carton for Complete Usage Information.
0001018

Motrin
Cold & Sinus
McNeil
Div. of McNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
LOT XXXXXX EXP 00/00
See Carton for Complete Usage Information.
0001018

Motrin
Cold & Sinus
McNeil
Div. of McNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
LOT XXXXXX EXP 00/00
See Carton for Complete Usage Information.
0001018

Motrin
Cold & Sinus
McNeil
Div. of McNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
LOT XXXXXX EXP 00/00
See Carton for Complete Usage Information.
0001018