

LACHMAN CONSULTANT SERVICES, INC.
Westbury, NY 11590

ATTACHMENT C

IV

**PROPOXYPHENE HYDROCHLORIDE,
ACETAMINOPHEN AND CAFFEINE CAPSULES**
(65 mg/ 389 mg/ 32.4 mg)


Rx only

Code 000000

Rev. 03/2002

DESCRIPTION


Propoxyphene Hydrochloride, USP is an odorless, white crystalline powder with a bitter taste. It is freely soluble in water. Chemically, it is (2S,3R)-(+)-4-(Dimethylamino)-3-methyl-1,2-diphenyl-2-butanol propionate (ester) hydrochloride, which can be represented by the accompanying structural formula.



$C_{22}H_{29}NO_2 \cdot HCl$

M.W. = 375.93


Acetaminophen (4'-hydroxyacetanilide), a slightly bitter, white, odorless, crystalline powder, is a non-opiate, non-salicylate analgesic and antipyretic. It has the following structural formula:



$C_8H_9NO_2$

MW = 151.16

Caffeine (1,3,7-trimethylxanthine), a bitter, white powder or white-glistening needles, is a central nervous system stimulant. It has the following structural formula:



$C_8H_{10}N_4O_2$

MW = 194.19

Each capsule of propoxyphene hydrochloride, acetaminophen and caffeine contains 65 mg (172.9 μ mol) propoxyphene hydrochloride, 389 mg (2,159 μ mol) acetaminophen and 32.4 mg (166.8 μ mol) caffeine.

Each capsule also contains the following inactive ingredients: to be determined.

CLINICAL PHARMACOLOGY

Propoxyphene is a centrally acting narcotic analgesic agent. Equimolar doses of propoxyphene hydrochloride or napsylate provide similar plasma concentrations. Following administration of 65, 130, or 195 mg of propoxyphene hydrochloride, the bioavailability of propoxyphene is equivalent to that of 100, 200, or 300 mg respectively of propoxyphene napsylate. Peak plasma concentrations of propoxyphene are reached in 2 to 2 1/2 hours. After a 65-mg oral dose of propoxyphene hydrochloride, peak plasma levels of 0.05 to 0.1 µg/mL are achieved.

Repeated doses of propoxyphene at 6-hour intervals lead to increasing plasma concentrations, with a plateau after the ninth dose at 48 hours.

Propoxyphene is metabolized in the liver to yield norpropoxyphene. Propoxyphene has a half-life of 6 to 12 hours, whereas that of norpropoxyphene is 30 to 36 hours.

Norpropoxyphene has substantially less central-nervous-system-depressant effect than propoxyphene but a greater local anesthetic effect, which is similar to that of amitriptyline and antiarrhythmic agents, such as lidocaine and quinidine.

In animal studies in which propoxyphene and norpropoxyphene were continuously infused in large amounts, intracardiac conduction time (PR and QRS intervals) was prolonged. Any intracardiac conduction delay attributable to high concentrations of norpropoxyphene may be of relatively long duration.

ACTIONS

Propoxyphene is a mild narcotic analgesic structurally related to methadone. The potency of propoxyphene hydrochloride is from two thirds to equal that of codeine.

The combination of propoxyphene with a mixture of acetaminophen and caffeine produces greater analgesia than that produced by either propoxyphene or acetaminophen and caffeine administered alone.

INDICATION

This product is indicated for the relief of mild to moderate pain, either when pain is present alone or when it is accompanied by fever.

CONTRAINDICATION

Hypersensitivity to propoxyphene, acetaminophen, or caffeine.

WARNINGS

- Do not prescribe propoxyphene for patients who are suicidal or addiction-prone.
- Prescribe propoxyphene with caution for patients taking tranquilizers or antidepressant drugs and patients who use alcohol in excess.
- Tell your patients not to exceed the recommended dose and to limit their intake of alcohol.

Propoxyphene products in excessive doses, either alone or in combination with other CNS depressants, including alcohol, are a major cause of drug-related deaths. Fatalities within the first hour of overdosage are not uncommon. In a survey of deaths due to overdosage conducted in 1975, in approximately 20% of the fatal cases, death occurred within the first hour (5% occurred within 15 minutes). Propoxyphene should not be taken in doses higher than those recommended by the physician. The judicious prescribing of propoxyphene is essential to the safe use of this drug. With patients who are depressed or suicidal, consideration should be given to the use of non-narcotic analgesics. Patients should be cautioned about the concomitant use of propoxyphene products and alcohol because of potentially serious CNS-additive effects of these agents. Because of its added depressant effects, propoxyphene should be prescribed with caution for those patients whose medical condition requires the concomitant administration of sedatives, tranquilizers, muscle relaxants, antidepressants, or other CNS-depressant drugs. Patients should be advised of the additive depressant effects of these combinations.

Many of the propoxyphene-related deaths have occurred in patients with previous histories of emotional disturbances or suicidal ideation or attempts as well as histories of misuse of tranquilizers, alcohol, and other CNS-active drugs. Some deaths have occurred as a consequence of the accidental ingestion of excessive quantities of propoxyphene alone or in combination with other drugs. Patients taking propoxyphene should be warned not to exceed the dosage recommended by the physician.

Drug Dependence – Propoxyphene, when taken in higher-than-recommended doses over long periods of time, can produce drug dependence characterized by psychic dependence and, less frequently, physical dependence and tolerance. Propoxyphene will only partially suppress the withdrawal syndrome in individuals physically dependent on morphine or other narcotics. The abuse liability of propoxyphene is qualitatively similar to that of codeine although quantitatively less, and propoxyphene should be prescribed with the same degree of caution appropriate to the use of codeine.

Usage in Ambulatory Patients – Propoxyphene may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a car or operating machinery. The patient should be cautioned accordingly.

PRECAUTIONS

General – Propoxyphene should be administered with caution to patients with hepatic or renal impairment since higher serum concentrations or delayed elimination may occur.

Drug Interactions – The CNS-depressant effect of propoxyphene is additive with that of other CNS depressants, including alcohol.

As is the case with medicinal agents, propoxyphene may slow the metabolism of a concomitantly administered drug. Should this occur, the higher serum concentrations of that drug may result in increased pharmacologic or adverse effects of that drug. Such occurrences have been reported when propoxyphene was administered to patients on antidepressants, anticonvulsants, or warfarin-like drugs. Severe neurologic signs, including coma, have occurred with concurrent use of carbamazepine.

Usage in Pregnancy – Safe use in pregnancy has not been established relative to possible adverse effects on fetal development. Instances of withdrawal symptoms in the neonate have been reported following usage during pregnancy. Therefore, propoxyphene should not be used in pregnant women unless, in the judgment of the physician, the potential benefits outweigh the possible hazards.

Usage in Nursing Mothers – Low levels of propoxyphene have been detected in human milk. In postpartum studies involving nursing mothers who were given propoxyphene, no adverse effects were noted in infants receiving mother's milk.

Usage in Pediatric Patients – Safety and effectiveness in pediatric patients have not been established.

Usage in the Elderly – The rate of propoxyphene metabolism may be reduced in some patients. Increased dosing interval should be considered.

A Patient Information Sheet is available for this product. See text following "How Supplied" section below.

ADVERSE REACTIONS

In a survey conducted in hospitalized patients, less than 1% of patients taking propoxyphene hydrochloride at recommended doses experienced side effects. The most frequently reported were dizziness, sedation, nausea, and vomiting. Some of these adverse reactions may be alleviated if the patient lies down.

Other adverse reactions include constipation, abdominal pain, skin rashes, lightheadedness, headache, weakness, euphoria, dysphoria, hallucinations, and minor visual disturbances.

Propoxyphene therapy has been associated with abnormal liver function tests and, more rarely, with instances of reversible jaundice (including cholestatic jaundice).

Renal papillary necrosis may result from chronic acetaminophen use, particularly when the dosage is greater than recommended and when combined with aspirin.

Subacute painful myopathy has occurred following chronic propoxyphene overdosage.

DOSAGE AND ADMINISTRATION

This product is given orally. The usual dosage is 65 mg propoxyphene hydrochloride, 369 mg acetaminophen, and 32.4 mg caffeine every 4 hours as needed for pain.

The maximum recommended dose of propoxyphene hydrochloride is 390 mg/day.

Consideration should be given to a reduced total daily dosage in patients with hepatic or renal impairment.

MANAGEMENT OF OVERDOSAGE

In all cases of suspected overdosage, call your regional Poison Control Center to obtain the most up-to-date information about the treatment of overdose. This recommendation is made because, in general, information regarding the treatment of overdosage may change more rapidly than do package inserts.

Initial consideration should be given to the management of the CNS effects of propoxyphene overdosage. Resuscitative measures should be initiated promptly.

Symptoms of Propoxyphene Overdosage – The manifestations of acute overdosage with propoxyphene are those of narcotic overdosage. The patient is usually somnolent but may be stuporous or comatose and convulsing. Respiratory depression is characteristic. The ventilatory rate and/or tidal volume is decreased, which results in cyanosis and hypoxia. Pupils, initially pinpoint, may become dilated as hypoxia increases. Cheyne-Stokes respiration and apnea may occur. Blood pressure and heart rate are usually normal initially, but blood pressure falls and cardiac performance deteriorates, which ultimately results in pulmonary edema and circulatory collapse, unless the respiratory depression is corrected and adequate ventilation is restored promptly. Cardiac arrhythmias and conduction delay may be present. A combined respiratory-metabolic acidosis occurs owing to retained CO₂ (hypercapnia) and

to lactic acid formed during anaerobic glycolysis. Acidosis may be severe if large amounts of salicylates have also been ingested. Death may occur.

Treatment of Propoxyphene Overdosage – Attention should be directed first to establishing a patent airway and to restoring ventilation. Mechanically assisted ventilation, with or without oxygen, may be required, and positive pressure respiration may be desirable if pulmonary edema is present. The narcotic antagonist naloxone will markedly reduce the degree of respiratory depression, and 0.4 to 2 mg should be administered promptly, preferably intravenously. If the desired degree of counteraction with improvement in respiratory functions is not obtained, naloxone should be repeated at 2- to 3-minute intervals. The duration of action of the antagonist may be brief. If no response is observed after 10 mg of naloxone have been administered, the diagnosis of propoxyphene toxicity should be questioned. Naloxone may also be administered by continuous intravenous infusion.

Treatment of Propoxyphene Overdosage in Pediatric Patients – The usual initial dose of naloxone in pediatric patients is 0.01 mg/kg body weight given intravenously. If this dose does not result in the desired degree of clinical improvement, a subsequent increased dose of 0.1 mg/kg body weight may be administered. If an IV route of administration is not available, naloxone may be administered IM or subcutaneously in divided doses. If necessary, naloxone can be diluted with Sterile Water for injection.

Blood gases, pH, and electrolytes should be monitored in order that acidosis and any electrolyte disturbance present may be corrected promptly. Acidosis, hypoxia, and generalized CNS depression predispose to the development of cardiac arrhythmias. Ventricular fibrillation or cardiac arrest may occur and necessitate the full complement of cardiopulmonary resuscitation (CPR) measures. Respiratory acidosis rapidly subsides as ventilation is restored and hypercapnia eliminated, but lactic acidosis may require intravenous bicarbonate for prompt correction.

Electrocardiographic monitoring is essential. Prompt correction of hypoxia, acidosis, and electrolyte disturbance (when present) will help prevent these cardiac complications and will increase the effectiveness of agents administered to restore normal cardiac function.

In addition to the use of a narcotic antagonist, the patient may require careful titration with an anticonvulsant to control convulsions. Analeptic drugs (for example, caffeine or amphetamine) should not be used because of their tendency to precipitate convulsions.

General supportive measures, in addition to oxygen, include, when necessary, intravenous fluids, vasopressor-inotropic compounds, and, when infection is likely, anti-infective agents. Gastric lavage may be useful and activated charcoal can adsorb a significant amount of ingested propoxyphene. Dialysis is of little value in poisoning due to propoxyphene. Efforts should be made to determine whether other agents, such as alcohol, barbiturates, tranquilizers, or other CNS depressants, were also ingested, since these increase CNS depression as well as cause specific toxic effects.

Symptoms of Acetaminophen Overdosage: Shortly after oral ingestion of an overdose of acetaminophen and for the next 24 hours, anorexia, nausea, vomiting, diaphoresis, general malaise and abdominal pain have been noted. The patient may then present no symptoms, but evidence of liver dysfunction may become apparent up to 72 hours after ingestion, with elevated serum transaminase and lactic dehydrogenase levels, an increase in serum bilirubin concentrations, and a prolonged prothrombin time. Death from hepatic failure may result 3 to 7 days after overdose.

Acute renal failure may accompany the hepatic dysfunction and has been noted in patients who do not exhibit signs of fulminant hepatic failure. Typically, renal impairment is more apparent 6 to 9 days after ingestion of the overdose.

Treatment of Acetaminophen Overdosage: Acetaminophen in massive overdosage may cause hepatic toxicity in some patients. In all cases of suspected overdose, immediately call your regional poison center or the Rocky Mountain Poison Center's toll-free number (800) 525-6115 for assistance in diagnosis and for directions in the use of N-acetylcysteine as an antidote.

In adults, hepatic toxicity has rarely been reported with acute overdoses of less than 10 g and fatalities with less than 15 g. Importantly, young children seem to be more resistant than adults to the hepatotoxic effect of an acetaminophen overdose. Despite this, the measures outlined below should be initiated in any adult or pediatric patient suspected of having ingested an acetaminophen overdose.

Because clinical and laboratory evidence of hepatic toxicity may not be apparent until 48 to 72 hours postingestion, liver function studies should be obtained initially and repeated at 24-hour intervals.

Consider emptying the stomach promptly by lavage or by induction of emesis with syrup of ipecac. Patients' estimates of the quantity of a drug ingested are notoriously unreliable. Therefore, if an acetaminophen overdose is suspected, a serum acetaminophen assay should be obtained as early as possible, but no sooner than 4 hours following ingestion. The antidote, N-acetylcysteine, should be administered as early as possible, and within 16 hours of the overdose ingestion for optimal results. Following recovery, there are no residual, structural, or functional hepatic abnormalities.

HOW SUPPLIED

Propoxyphene Hydrochloride, Acetaminophen and Caffeine Capsules,
65 mg/ 389 mg/ 32.4 mg

NDC# 00000-000-00

Dosage Form: Capsule

Shape, Color, and Scoring: To be determined.

Packaging: To be determined.

STORAGE: Store at controlled room temperature 15° to 30°C (59° to 86°F) [See USP].

PHARMACIST: Dispense in a tight, light-resistant container with a child-resistant closure.

Patient Information Sheet YOUR PRESCRIPTION FOR A PROPOXYPHENE PRODUCT



Summary

Products containing propoxyphene are used to relieve pain.

LIMIT YOUR INTAKE OF ALCOHOL WHILE TAKING THIS DRUG. Make sure your doctor knows if you are taking tranquilizers, sleep aids, antidepressants, antihistamines, or any other drugs that make you sleepy. Combining propoxyphene with alcohol or these drugs in excessive doses is dangerous.

Use care while driving a car or using machines until you see how the drug affects you because propoxyphene can make you sleepy. Do not take more of the drug than your doctor prescribed. Dependence has occurred when patients have taken propoxyphene for a long period of time at doses greater than recommended.

The rest of this leaflet gives you more information about propoxyphene. Please read it and keep it for future use.

Uses of Propoxyphene

Products containing Propoxyphene are used for the relief of mild to moderate pain. Products that contain Propoxyphene plus acetaminophen or caffeine are prescribed for the relief of pain or pain associated with fever.

Before Taking Propoxyphene

Make sure your doctor knows if you have ever had an allergic reaction to propoxyphene, acetaminophen, or caffeine.

The effect of propoxyphene in pediatric patients under 12 has not been studied. Therefore, use of the drug in this age group is not recommended.

How to Take Propoxyphene

Follow your doctor's directions exactly. Do not increase the amount you take without your doctor's approval. If you miss a dose of the drug, do not take twice as much the next time.

Pregnancy

Do not take propoxyphene during pregnancy unless your doctor knows you are pregnant and specifically recommends its use. Cases of temporary dependence in the newborn have occurred when the mother has taken propoxyphene consistently in the weeks before delivery. As a general principle, no drug should be taken during pregnancy unless it is clearly necessary.

General Cautions

Heavy use of alcohol with propoxyphene is hazardous and may lead to overdose symptoms (see "Overdose" below). THEREFORE, LIMIT YOUR INTAKE OF ALCOHOL WHILE TAKING PROPOXYPHENE.

Combinations of excessive doses of propoxyphene, alcohol, and tranquilizers are dangerous. Make sure your doctor knows if you are taking tranquilizers, sleep aids, antidepressant drugs, antihistamines, or any other drugs that make you sleepy. The use of these drugs with propoxyphene increases their sedative effects and may lead to overdose symptoms, including death (see "Overdose" below).

Propoxyphene may cause drowsiness or impair your mental and/or physical abilities; therefore, use caution when driving a vehicle or operating dangerous machinery. DO NOT perform any hazardous task until you have seen your response to this drug.

Propoxyphene may increase the concentration in the body of medications such as anticoagulants ("blood thinners"), antidepressants, or drugs used for epilepsy. The result may be excessive or adverse effects of these medications. Make sure your doctor knows if you are taking any of these medications.

Dependence

You can become dependent on propoxyphene if you take it in higher than recommended doses over a long period of time. Dependence is a feeling of need for the drug and a feeling that you cannot perform normally without it.

Overdose

An overdose of Propoxyphene, alone or in combination with other drugs, including alcohol, may cause weakness, difficulty in breathing, confusion, anxiety, and more severe drowsiness and dizziness. Extreme overdose may lead to unconsciousness and death.

If the propoxyphene product contains acetaminophen, the overdose symptoms include nausea, vomiting, lack of appetite, and abdominal pain. Liver damage may occur.

In any suspected overdose situation, contact your doctor or nearest hospital emergency room. **GET EMERGENCY HELP IMMEDIATELY.**

KEEP THIS DRUG AND ALL DRUGS OUT OF THE REACH OF THE PEDIATRIC POPULATION.

Possible Side Effects

When propoxyphene is taken as directed, side effects are infrequent. Among those reported are drowsiness, dizziness, nausea, and vomiting. If these effects occur, it may help if you lie down and rest.

Less frequently reported side effects are constipation, abdominal pain, skin rashes, lightheadedness, headache, weakness, hallucinations, minor visual disturbances, and feelings of elation or discomfort. If side effects occur and concern you, contact your doctor.

Other Information

The safe and effective use of propoxyphene depends on your taking it exactly as directed. This drug has been prescribed specifically for you and your present condition. Do not give this drug to others who may have similar symptoms. Do not use it for any other reason.

If you would like more information about propoxyphene, ask your doctor or pharmacist. They have a more technical leaflet (professional labeling) you may read.

**Manufactured by:
Manufacturer**

Code 000000

Rev. 03/2002

Ex USA Airbill
Express

FedEx
Tracking
Number

8312 2318 3820

Form
ID No.

0215

SPH13

Recipient's Copy

is portion can be removed for Recipient's records.

1/27/02 FedEx Tracking Number 831223183820

R. POLLUCK Phone

LACHMAN CONSULTANT SERVICE INC

1600 STEWART AVE STE 604

WESTBURY State NY ZIP 11590

Additional Billing Reference

DOCKETS MANAGEMENT BRANCH Phone

FOOD & DRUG ADMINISTRATION

5630 FISHERS LANE

Room 1061

ROCKVILLE State MD ZIP 20852



0199324048

4a Express Package Service

FedEx Priority Overnight
Next business morning

FedEx Standard Overnight
Next business afternoon

Packages up to 150 lbs.

FedEx First Overnight
Earliest next business morning
delivery to select locations

FedEx 2Day
Second business day

FedEx Express Saver
Third business day

NEW FedEx Extra Hours
Later drop-off with next business
afternoon delivery to select locations

4b Express Freight Service

FedEx 1Day Freight*
Next business day

FedEx 2Day Freight
Second business day

FedEx 3Day Freight
Third business day

Packages over 150 lbs.

5 Packaging

FedEx Envelope*

FedEx Pak*
Includes FedEx Small Pak, FedEx
Large Pak, and FedEx Sturdy Pak

Other Pkg.
Includes FedEx Box, FedEx
Tube, and customer pkg.

6 Special Handling

SATURDAY Delivery
Available only for FedEx Priority
Overnight and FedEx 2Day
to select ZIP codes

HOLD Weekday
at FedEx Location
Not available for
FedEx First Overnight

HOLD Saturday
at FedEx Location
Available only for
FedEx Priority Overnight
and FedEx 2Day to
select locations

Does this shipment contain dangerous goods?
One box must be checked.

No

Yes
As per attached
Shipper's Declaration

Yes
Shipper's Declaration
not required

Dry Ice
Dry Ice, 9, UN 1845 x kg

Dangerous Goods (incl. Dry Ice) cannot be shipped in FedEx packaging or with
FedEx Extra Hours service.

Cargo Aircraft Only

7 Payment Bill to:

Sender
Acct. No. in Section
1 will be billed

Recipient

Third Party

Credit Card

Obtain Recip.
Acct. No.

Cash/Check



Total Packages

Total Weight

Total Charges

*Our liability is limited to \$100 unless you declare a higher value. See the FedEx Service Guide for details.

8 Release Signature

Sign to authorize delivery without obtaining signature.

By signing you authorize us to deliver this shipment without obtaining a signature
and agree to indemnify and hold us harmless from any resulting claims.
Questions? Visit our Web site at fedex.com
or call 1 800 Go FedEx® 800.463.3339.
SRS • Rev. Date 7/01 • Part #1572525 • ©1994-2001 FedEx • PRINTED IN U.S.A.

447