

OXYTETRACYCLINE
TYPE C
BLUE BIRD
TURKEY FEED A
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For turkeys for control of hexamitiasis caused by *Hexamita meleagridis* susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT

Oxytetracycline 100 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min)..... %
Lysine (Min) %
Methionine (Min)..... %
Crude Fat (Min) %
Crude Fiber (Max) %
Calcium (Min)..... %
Calcium (Max) %
Phosphorus (Min)..... %
Salt, if added (Min)..... %
Salt, if added (Max) %
Total Sodium (Min)* %
Total Sodium (Max)* %

*Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed 100 g oxytetracycline per ton of feed continuously for 7-14 days.

**WARNING: ZERO-DAY WITHDRAWAL. DO NOT ADMINISTER TO TURKEYS
PRODUCING EGGS FOR HUMAN CONSUMPTION.**

_____ lbs (_____ kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000