

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Menomune® – A/C/Y/W-135 vaccine safely and effectively. See full prescribing information for Menomune® – A/C/Y/W-135 vaccine.

Menomune® – A/C/Y/W-135, Meningococcal Polysaccharide Vaccine, Groups A, C, Y and W-135 Combined
Solution for subcutaneous injection

Initial U.S. Approval: 1981

RECENT MAJOR CHANGES

Warnings and Precautions, Latex (5.1) [4/2016]
The latex warning has been removed for the diluent vial.

INDICATIONS AND USAGE

- Menomune – A/C/Y/W-135 vaccine is a vaccine indicated for active immunization against invasive meningococcal disease caused by *Neisseria meningitidis* serogroups A, C, Y, and W-135. Menomune – A/C/Y/W-135 vaccine is approved for use in persons 2 years of age and older. Menomune – A/C/Y/W-135 vaccine does not prevent *N meningitidis* serogroup B disease. (1)

DOSAGE AND ADMINISTRATION

- Single 0.5 mL subcutaneous injection. (2)

DOSAGE FORMS AND STRENGTHS

- Menomune – A/C/Y/W-135 vaccine is supplied as a single dose or multidose (10 dose) vial of lyophilized vaccine, with corresponding single dose or multidose vial of diluent. After reconstitution of the lyophilized vaccine with the diluent, each dose consists of a 0.5 mL solution for injection. (3)

CONTRAINDICATIONS

- Do not administer to anyone with a history of a severe allergic reaction (e.g., anaphylaxis) to Menomune – A/C/Y/W-135 vaccine or any component of the vaccine. (4)

WARNINGS AND PRECAUTIONS

- The stopper to the vial of lyophilized vaccine contains natural rubber latex which may cause allergic reactions. (5.1)

ADVERSE REACTIONS

- Common (>10%) adverse events in children 2 to 10 years of age were injection site pain, drowsiness, irritability, and diarrhea. (6)
- Common (>10%) adverse events in persons 11 to 55 years of age were pain, redness, and induration at the injection site, headache, fatigue, malaise, arthralgia, and diarrhea. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Sanofi Pasteur Inc. at 1-800-822-2463 (1-800-VACCINE) or VAERS at 1-800-822-7967 or <http://vaers.hhs.gov>.

DRUG INTERACTIONS

- Do not mix Menomune – A/C/Y/W-135 vaccine with other vaccines in the same syringe or vial. (7)
- Immunosuppressive therapies may reduce the immune response to Menomune – A/C/Y/W-135 vaccine. (7)
- No safety and immunogenicity data are available on the concomitant administration of Menomune – A/C/Y/W-135 vaccine with other US licensed vaccines. (7)

USE IN SPECIFIC POPULATIONS

- Pregnancy: No human or animal data. Use only if clearly needed. (8.1)
- Geriatrics: Not known whether persons aged 65 years and older respond differently from younger persons. (8.5)

See 17 for Patient Counseling Information.

Revised: []

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Administration
- 2.2 Primary Immunization
- 2.3 Revaccination

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

- 4.1 Hypersensitivity

5 WARNINGS AND PRECAUTIONS

- 5.1 Latex
- 5.2 Management of Acute Allergic Reactions
- 5.3 Moderate or Severe Acute Illness
- 5.4 Limitations of Vaccine Effectiveness
- 5.5 Altered Immunocompetence

6 ADVERSE REACTIONS

- 6.1 Data from Clinical Studies
- 6.2 Data from Post-Marketing Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action

13 NON-CLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, mutagenesis, impairment of fertility

14 CLINICAL STUDIES

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

1 **FULL PRESCRIBING INFORMATION:**

2 **1. INDICATIONS AND USAGE**

3 Menomune® – A/C/Y/W-135, Meningococcal Polysaccharide Vaccine, Groups A, C, Y and W-
4 135 Combined, is indicated for active immunization for the prevention of invasive
5 meningococcal disease caused by *Neisseria meningitidis* serogroups A, C, Y, and W-135.

6 Menomune – A/C/Y/W-135 vaccine is approved for use in persons 2 years of age and older.

7 Menomune – A/C/Y/W-135 vaccine does not prevent *N meningitidis* serogroup B disease.

8

9 **2. DOSAGE AND ADMINISTRATION**

10 **2.1. Administration**

11 The package contains a vial of lyophilized vaccine and a vial of diluent. The lyophilized vaccine
12 should be a white or off-white color to a light beige color. The diluent used for reconstitution is a
13 clear liquid.

14

15 After removing the "flip-off" caps, cleanse the vaccine and diluent vial stoppers with a suitable
16 germicide. Do not remove the vial stoppers or metal seals holding them in place. Using a suitable
17 sterile needle and syringe and aseptic technique, withdraw the supplied diluent (0.6 mL for single
18 dose presentation and 6.0 mL for multidose presentation) (Refer to Figure 1) and inject into the
19 vial containing the lyophilized vaccine (Refer to Figure 2). Swirl the vial until the vaccine is
20 thoroughly dissolved (Refer to Figure 3). When reconstituted, the vaccine should be a clear,
21 colorless liquid.

22

1 Parenteral drug products should be inspected visually for particulate matter and discoloration
2 prior to administration, whenever solution and container permit. If either of these conditions
3 exists, the vaccine should not be administered.

4

5 Using a suitable sterile needle and syringe and aseptic technique, withdraw (Refer to Figure 4)
6 and administer a 0.5 mL dose of Menomune – A/C/Y/W-135 vaccine by subcutaneous injection.

7 Use a separate sterile needle and syringe and aseptic technique for each dose withdrawn from the
8 multidose vial.

9

10 Do not administer this product intravenously or intramuscularly.

11


12 The preferred site of administration is the deltoid region.

13

14 **Menomune vaccine: Instructions for reconstitution**


15 **Figure 1**

16 For the **single-dose presentation**, withdraw **0.6 mL** of the supplied diluent; for the **multidose**
17 **presentation**, withdraw **6.0 mL**.




3 **Figure 2**

4 Insert the syringe needle through the stopper of the vial of lyophilized Menomune vaccine
5 component and inject the diluent into the vial.



8 **Figure 3**

9 Swirl the vial until the lyophilized vaccine component is thoroughly dissolved.



1


2

3 **Figure 4**

4 After reconstitution, withdraw 0.5 mL of Menomune vaccine and administer subcutaneously.

5 Use the single-dose presentation immediately after reconstitution. The multidose presentation

6 may be used for up to 35 days after reconstitution if stored at 2° to 8°C.



7

8

9 The vaccine should not be combined through reconstitution or mixed with any other vaccine.

10

1 Vaccine supplied in single dose vials should be used immediately after reconstitution. Vaccine
2 supplied in multidose vials may be used for up to 35 days after reconstitution if stored at 2° to
3 8°C (35° to 46°F). [See *How Supplied/Storage and Handling* (16.2).]
4

5 **2.2. Primary Immunization**

6 Primary immunization with Menomune – A/C/Y/W-135 vaccine consists of a single 0.5 mL dose
7 administered subcutaneously.
8

9 The ACIP (Advisory Committee on Immunization Practices) has specific recommendations for
10 use of meningococcal vaccines. (1) (2) (3)
11

12 **2.3. Revaccination**

13 The ACIP has recommendations for revaccination against meningococcal disease for persons at
14 high risk who were previously vaccinated with Menomune – A/C/Y/W-135 vaccine. (1) (3) If
15 Menomune – A/C/Y/W-135 vaccine is used for revaccination, the dose is 0.5 mL administered
16 subcutaneously.
17

18 **3. DOSAGE FORMS AND STRENGTHS**

19 Menomune – A/C/Y/W-135 vaccine is supplied as a lyophilized vaccine, in a single dose or a
20 multidose (10 dose) vial, with corresponding single dose or multidose vial of diluent. The
21 lyophilized vaccine is reconstituted with the diluent [see *Dosage and Administration* (2.1)]. After

1 reconstitution, each dose consists of a 0.5 mL solution for injection. See *Description* (11) for the
2 complete listing of ingredients.

3

4 **4. CONTRAINDICATIONS**

5 **4.1. Hypersensitivity**

6 Do not administer to anyone with a history of a severe allergic reaction (e.g., anaphylaxis) to
7 Menomune – A/C/Y/W-135 vaccine or any component of the vaccine [see *Description* (11)].

8

9 **5. WARNINGS AND PRECAUTIONS**

10 **5.1. Latex**

11 The stopper to the vial of lyophilized vaccine contains natural rubber latex which may cause
12 allergic reactions.

13

14 **5.2. Management of Acute Allergic Reactions**

15 Appropriate medical treatment must be available to manage possible anaphylactic reactions
16 following administration of the vaccine.

17

18 **5.3. Moderate or Severe Acute Illness**

19 To avoid diagnostic confusion between manifestations of an acute illness and possible vaccine
20 adverse effects, vaccination with Menomune – A/C/Y/W-135 vaccine should be postponed in
21 persons with moderate or severe acute illness. (4)

22

1 **5.4. Limitations of Vaccine Effectiveness**

2 Menomune – A/C/Y/W-135 vaccine may not protect all recipients.

3

4 **5.5. Altered Immunocompetence**

5 Persons who are immunosuppressed, including persons receiving immunosuppressive therapy,
6 may have a diminished immune response to Menomune – A/C/Y/W-135 vaccine [see *Drug*
7 *Interactions (7)*].

8

9 **6. ADVERSE REACTIONS**

10 **6.1. Data from Clinical Studies**

11 Because clinical trials are conducted under widely varying conditions, adverse reaction rates
12 observed in the clinical trials of a vaccine cannot be directly compared to rates in the clinical
13 trials of another vaccine and may not reflect the rates observed in clinical practice.

14

15 In three clinical trials primarily designed to assess the safety and immunogenicity of another
16 vaccine, Menactra® [Meningococcal (Groups A, C, Y and W-135) Polysaccharide Diphtheria
17 Toxoid Conjugate Vaccine], participants were randomized to receive Menactra or Menomune –
18 A/C/Y/W-135 vaccine, which was used as a control vaccine. In these three trials, 1519 children
19 2-10 years of age, 972 persons 11-18 years of age, and 1170 adults 18-55 years of age,
20 respectively, received a dose of Menomune – A/C/Y/W-135 vaccine. Overall, of the children 2-
21 10 years of age who received Menactra or Menomune – A/C/Y/W-135 vaccine, 68% were
22 enrolled at US sites and 32% were enrolled at a Chilean site. The median ages of US and Chilean

1 children were 6 and 5 years, respectively; overall, 50.5% were males and 92.0% were Caucasian.
2 Among participants 11-55 years of age who received Menactra or Menomune – A/C/Y/W-135
3 vaccine, all were enrolled at US sites; 54.8% were female; 87.7% were Caucasian.

4
5 Solicited local and systemic reactions were monitored daily for 7 days post-vaccination using a
6 diary card. Information on serious adverse events was collected at interim clinic visits and by
7 telephone interview conducted 6 months post-vaccination. At least 94% of participants from the
8 three studies completed the 6-month follow-up.

9 10 **Serious Adverse Events**

11 Across the three studies, serious adverse events within 6-months following Menomune –
12 A/C/Y/W-135 vaccine were reported in 0.7% of 1519 children 2-10 years of age, 0.6% of 972
13 persons 11-18 years of age, and 1.7% of 1170 persons 18-55 years of age.

14 15 **Solicited Adverse Events^a**

16 The most commonly reported solicited adverse events in US children 2-10 years of age were
17 injection site pain, irritability, and diarrhea. (Table 1)

18
19 The most commonly reported solicited adverse events in adolescents, ages 11-18 years, and
20 adults, ages 18-55 years, were injection site pain, headache, and fatigue. (Table 2)

21 _____

^a Events in Table 1 and Table 2 were collected in the clinical trials under “Solicited local and systemic reactions.”

- 1 **Table 1: Percentage of US Participants 2-10 Years of Age Reporting Solicited Adverse**
- 2 **Events Within 7 Days Following Administration of Menomune – A/C/Y/W-135 Vaccine**

	N=1019-1027*		
Event	Any	Moderate	Severe
	%	%	%
<i>General disorders and administration site conditions</i>			
Injection site reaction			
Pain†	26.1	2.5	0.0
Redness‡	7.9	0.5	0.0
Induration‡	4.2	0.6	0.0
Swelling‡	2.8	0.3	0.0
Systemic events			
Fever§	5.2	1.7	0.2
<i>Gastrointestinal disorders</i>			
Anorexia	8.7	1.3	0.8
Vomiting¶	2.7	0.7	0.6
Diarrhea#	11.8	2.5	0.3
<i>Nervous system disorders</i>			
Drowsiness**	11.2	2.5	0.5
Irritability††	12.2	2.6	0.6
Seizures‡‡	0.0	N/A	N/A
<i>Musculoskeletal and connective tissue disorders</i>			

Arthralgia§§	5.3	0.7	0.0
<i>Skin and subcutaneous disorders</i>			
Rash‡‡	3.0	N/A	N/A

- 1 * N = The total number of participants with reported data. The N is 1027 for all solicited events except
2 fever (N=1019). The percentage is based on N.
- 3 † Moderate: Discomforting, interfered with or limited usual arm movement, Severe: Disabling, child unable
4 to move arm.
- 5 ‡ Moderate: 1.0–2.0 inches; Severe: >2.0 inches.
- 6 § Oral equivalent temperature; Moderate: 38.4-39.4°C, Severe: ≥39.5°C.
- 7 || Moderate: Skipped 2 meals, Severe: skipped ≥3 meals.
- 8 ¶ Moderate: 2 episodes, Severe: ≥3 episodes.
- 9 # Moderate: 3-4 episodes, Severe: ≥5 episodes.
- 10 ** Moderate: Interferes with normal activities, Severe: disabling, unwilling to engage in play or interaction
11 with others.
- 12 †† Moderate: 1-3 hours duration, Severe: >3 hours duration.
- 13 ‡‡ These solicited adverse events were reported as present or absent only.
- 14 §§ Moderate: Decreased range of motion due to pain or discomfort, Severe: unable to move major joints
15 because of pain.
- 16
- 17

- 1 **Table 2: Percentage of Participants 11-55 Years of Age Reporting Solicited Adverse Events**
- 2 **Within 7 Days Following Administration of Menomune – A/C/Y/W-135 Vaccine**

Event	Study 1 N*=970			Study 2 N*=1159		
	Participants 11-18 years of age			Participants 18-55 years of age		
	Any	Moderate	Severe	Any	Moderate	Severe
<i>General disorders and administration site conditions</i>						
Injection site reaction						
Pain†	28.7	2.6	0.0	48.1	3.3	0.1
Redness‡	5.7	0.4	0.0	16.0	1.9	0.1
Induration‡	5.2	0.5	0.0	11.0	1.0	0.0
Swelling‡	3.6	0.3	0.0	7.6	0.7	0.0
Systemic events						
Fatigue§	25.1	6.2	0.2	32.3	6.6	0.4
Malaise§	16.8	3.4	0.4	22.3	4.7	0.9
Chills§	3.5	0.4	0.1	5.6	1.0	0.0
Fever	3.0	0.3	0.1	0.5	0.1	0.0
<i>Gastrointestinal disorders</i>						
Diarrhea¶	10.2	1.3	0.0	14.0	2.9	0.3
Anorexia#	7.7	1.1	0.2	9.9	1.6	0.4
Vomiting**	1.4	0.5	0.3	1.5	0.2	0.4
<i>Nervous system disorders</i>						
Headache††	29.3	6.5	0.4	41.8	8.9	0.9
Seizure‡‡	0.0	N/A	N/A	0.0	N/A	N/A

<i>Musculoskeletal and connective tissue disorders</i>						
Arthralgia§	10.2	2.1	0.1	16.0	2.6	0.1
<i>Skin and subcutaneous disorders</i>						
Rash‡‡	1.4	N/A	N/A	0.8	N/A	N/A

- 1 * N=Total number of participants with data.
- 2 † Moderate: Discomforting, interfered with or limited usual arm movement, Severe: Disabling, unable to
- 3 move arm.
- 4 ‡ Moderate: 1.0–2.0 inches; Severe: >2.0 inches.
- 5 § Moderate: Interferes with normal activities, Severe: disabling, requires bed rest.
- 6 || Oral equivalent temperature. Study 1: Moderate: 38.5-39.4°C, Severe: ≥39.5°C. Study 2: Moderate 39.0-
- 7 39.9°C, Severe: ≥40.0°C.
- 8 ¶ Moderate: 3-4 episodes, Severe: ≥5 episodes.
- 9 # Moderate: Skipped 2 meals, Severe: skipped ≥3 meals.
- 10 ** Moderate: 2 episodes, Severe: ≥3 episodes.
- 11 †† Moderate: discomforting enough to interfere with activities, Severe: disabling requires bed rest and
- 12 analgesics.
- 13 ‡‡ These solicited adverse events were reported as present or absent only.

14

15 6.2. Data from Post-Marketing Experience

16 The following adverse events have been spontaneously reported during post-approval use of

17 Menomune – A/C/Y/W-135 vaccine since 1993 through November 2008. Because these events

18 were reported voluntarily from a population of uncertain size, it is not always possible to reliably

1 estimate their frequency or to establish a causal relationship to Menomune – A/C/Y/W-135
2 vaccine exposure.

3

4 The following adverse events were included based on severity, frequency of reporting or the
5 strength of causal association to Menomune – A/C/Y/W-135 vaccine.

6

7 • Immune System Disorders

8 Hypersensitivity, such as rash, urticaria, pruritus, dyspnea, angioedema

9 • Nervous System Disorders

10 Headache, vasovagal syncope, dizziness, paresthesia, Guillain-Barré syndrome

11 • Gastrointestinal Disorders

12 Nausea, vomiting, diarrhea

13 • Musculoskeletal and Connective Tissue Disorders

14 Myalgia, arthralgia

15 • General Disorders and Administration Site Conditions

16 Fever, injection site reaction, malaise, asthenia, chills, fatigue

17

18 **7. DRUG INTERACTIONS**

19 Separate syringes and injection sites must be used in case of concomitant administration.

20 Do not mix Menomune – A/C/Y/W-135 vaccine with other vaccines in the same syringe or vial.

21

1 Immunosuppressive therapies may reduce the immune response to Menomune – A/C/Y/W-135
2 vaccine.

3

4 No safety and immunogenicity data are available on the concomitant administration of
5 Menomune – A/C/Y/W-135 vaccine with other US licensed vaccines.

6

7 **8. USE IN SPECIFIC POPULATIONS**

8 **8.1. Pregnancy**

9 Pregnancy Category C. Animal reproduction studies have not been conducted with Menomune –
10 A/C/Y/W-135 vaccine. It is also not known whether Menomune – A/C/Y/W-135 vaccine can
11 cause fetal harm when administered to a pregnant woman or can affect reproduction capacity.
12 Menomune – A/C/Y/W-135 vaccine should be given to a pregnant woman only if clearly
13 needed.

14

15 **8.3. Nursing Mothers**

16 It is not known whether Menomune – A/C/Y/W-135 vaccine is excreted in human milk. Because
17 many drugs are excreted in human milk, caution should be exercised when Menomune –
18 A/C/Y/W-135 vaccine is administered to a nursing woman.

19

20 **8.4. Pediatric Use**

21 Safety and effectiveness of Menomune – A/C/Y/W-135 vaccine in children below the age of 2
22 years have not been established.

1
2 During a meningococcal serogroup A epidemic in sub-Saharan Africa, children 3 months to 16
3 years of age were vaccinated with a high molecular weight serogroup A/C meningococcal
4 polysaccharide vaccine. In case-control studies, after 1 year of observation, vaccine efficacy
5 against meningococcal serogroup A disease was estimated to be 87% [90% Confidence Interval
6 (CI), 52% to 96%], overall. After 3 years, efficacy was estimated to be 67% (90% CI, 40% to
7 82%) among children who were 4-16 years of age at the time of vaccination and 8% (90% CI,
8 -102% to 58%) among children who were 1-3 years of age at the time of vaccination. (5)

9
10 The efficacy of a serogroup C meningococcal vaccine in infants and young children was
11 evaluated in a placebo-controlled trial during a serogroup C epidemic in Brazil. Vaccine efficacy
12 was estimated to be 12% (95% CI, -55% to 62%) among children 6 to 23 months of age and 55%
13 (95% CI, -4% to 72%) among children 24 to 36 months of age. (6)

14

15 **8.5. Geriatric Use**

16 Clinical studies of Menomune – A/C/Y/W-135 vaccine did not include sufficient numbers of
17 subjects aged 65 and over to determine whether they respond differently from younger subjects.

18

19 **11. DESCRIPTION**

20 Menomune – A/C/Y/W-135, Meningococcal Polysaccharide Vaccine, Groups A, C, Y and W-
21 135 Combined, is a vaccine for subcutaneous injection. Menomune – A/C/Y/W-135 vaccine
22 consists of a sterile lyophilized preparation of the group-specific polysaccharide antigens from *N*

1 *meningitidis*, Group A, Group C, Group Y, and Group W-135. *N meningitidis* are cultivated on
2 Mueller Hinton casein agar (7) and grown in Watson Scherp casamino acid media (8). The
3 purified polysaccharide is extracted from the *N meningitidis* cells and separated from the media
4 by procedures which include centrifugation, detergent precipitation, alcohol precipitation,
5 solvent or organic extraction, and diafiltration. No preservative is added during manufacture.

6
7 The diluent (0.6 mL) for the single dose presentation contains sterile, pyrogen-free distilled
8 water without preservative. The diluent (6 mL) for the multidose presentation contains sterile,
9 pyrogen-free distilled water and thimerosal, a mercury derivative, which is added as a
10 preservative for the reconstituted vaccine. [See *How Supplied/Storage and Handling* (16).]

11
12 After reconstitution with diluent, the vaccine is a clear colorless liquid solution. Each 0.5 mL
13 dose contains 50 mcg of polysaccharide from each of serogroups A, C, Y, and W-135.

14 Reconstituted vaccine from a multidose vial also contains 25 mcg mercury per dose.

15
16 Each dose of vaccine contains 2.5 mg to 5 mg of lactose added as a stabilizer. (9)

17
18 Potency is evaluated by measuring the molecular size of each polysaccharide component using a
19 column chromatography method as standardized by the US Food and Drug Administration
20 (FDA) and the World Health Organization (WHO) (10) for Meningococcal Polysaccharide
21 Vaccine.

22

1 The vial stopper of the lyophilized vaccine is made with natural rubber latex. The vial stopper of
2 the diluent is not made with natural rubber latex.

3

4 **12. CLINICAL PHARMACOLOGY**

5 **12.1. Mechanism of Action**

6 The presence of bactericidal anti-capsular meningococcal antibodies has been associated with
7 protection from invasive meningococcal disease. (11) (12) Menomune – A/C/Y/W-135 vaccine
8 induces the production of bactericidal antibodies specific to the capsular polysaccharides of
9 serogroups A, C, Y, and W-135.

10

11 **13. NON-CLINICAL TOXICOLOGY**

12 **13.1. Carcinogenesis, mutagenesis, impairment of fertility**

13 Menomune – A/C/Y/W-135 vaccine has not been evaluated for carcinogenic or mutagenic
14 potential or impairment of fertility.

15

16 **14. CLINICAL STUDIES**

17 Effectiveness of Menomune – A/C/Y/W-135 vaccine was inferred by evaluating the proportion
18 of children 2-10 years of age achieving a pre-specified level of serum bactericidal antibody and
19 the proportion of persons 11-55 years of age achieving a 4-fold increase from baseline in serum
20 bactericidal antibody, for each serogroup. Evidence for clinical efficacy against serogroup-
21 specific meningococcal disease in school-age children and adults has been obtained from
22 historical field trials and observational studies with other high molecular weight polysaccharide

1 vaccines containing meningococcal serogroup A and/or C components. (6) No studies have been
2 conducted to evaluate the efficacy of meningococcal polysaccharide vaccines against disease due
3 to serogroups Y and W-135.

4

5 Menomune – A/C/Y/W-135 vaccine was used as the control vaccine in three US, multi-center,
6 clinical trials designed primarily to evaluate the immunogenicity and safety of Menactra vaccine
7 in children (2-10 years old), adolescents (11-18 years old), and adults (18-55 years old),
8 respectively. In these trials, participants in the control arm received a dose of Menomune –
9 A/C/Y/W-135 vaccine. Sera were obtained before and approximately 28 days after vaccination.

10 The Serum Bactericidal Assay (SBA) used to test sera contained an exogenous complement
11 source that was either human (SBA-H) or, when correlated to SBA-H, baby rabbit (SBA-BR).

12 (13)

13

14 Data on immune responses, as measured by SBA-H, following Menomune – A/C/Y/W-135
15 vaccine in a subset of children 2-10 years old are presented in Table 3. Data on immune
16 responses, as measured by SBA-BR, following Menomune – A/C/Y/W-135 vaccine in
17 adolescents and adults are presented in Table 4.

18

1 **Table 3: Bactericidal Antibody Responses* to Menomune – A/C/Y/W-135 Vaccine 28 Days**
2 **After Vaccination for Subsets of Participants Aged 2-3 and 4-10 Years**

Menomune – A/C/Y/W-135 vaccine Aged 2-3 Years N†=50-53				Menomune – A/C/Y/W-135 vaccine Aged 4-10 Years N†=84			
Serogroup			(95% CI) ‡	Serogroup			(95% CI) ‡
A	% ≥1:8	64	50,77	A	% ≥1:8	55	44,66
	GMT	10	7,12		GMT	7	6,9
C	% ≥1:8	38	25,53	C	% ≥1:8	48	37,59
	GMT	11	5,21		GMT	12	7,18
Y	% ≥1:8	73	59,84	Y	% ≥1:8	92	84,97
	GMT	18	11,27		GMT	46	33,66
W-135	% ≥1:8	33	20,47	W-135	% ≥1:8	79	68,87
	GMT	5	3,6		GMT	20	14,27

3 The study was designed to show the safety and immunogenicity of Menactra vaccine are non-inferior to that of
4 Menomune – A/C/Y/W-135 vaccine. The table shows the immune response in Menomune – A/C/Y/W-135 vaccine
5 participants from this study.

6 * Serum Bactericidal Assay with an exogenous human complement (SBA-H) source.

7 † N = Number of subset participants with at least one valid serology result at Day 0 and Day 28.

8 ‡ The 95% CI for the Geometric Mean Titer (GMT) was calculated based on an approximation to the
9 normal distribution.

10

1 In participants 2-3 years of age with undetectable pre-vaccination SBA titers (ie, <4 at Day 0),
2 rates of seroconversion (defined as SBA titer ≥ 8 at Day 28) following Menomune – A/C/Y/W-
3 135 vaccine were 55%, serogroup A (n=16/29); 30%, serogroup C (n=13/43); 57%, serogroup Y
4 (n=17/30); 26%, serogroup W-135 (n=11/43).

5
6 In participants 4-10 years of age with undetectable pre-vaccination SBA titers (ie, <4 at Day 0),
7 rates of seroconversion (defined as SBA titer ≥ 8 at Day 28) following Menomune – A/C/Y/W-
8 135 vaccine were 48%, serogroup A (n=10/21); 38%, serogroup C (n=19/50); 84%, serogroup Y
9 (n=38/45); 68%, serogroup W-135 (n=26/38).

10

11

1 **Table 4: Bactericidal Antibody Responses* to Menomune – A/C/Y/W-135 Vaccine 28 Days**
2 **After Vaccination for Participants Aged 11-18 and 18-55 Years**

Menomune – A/C/Y/W-135 vaccine Aged 11-18 Years N [†] =423				Menomune – A/C/Y/W-135 vaccine Aged 18-55 Years N [†] =1098			
Serogroup			(95% CI) ‡	Serogroup			(95% CI) ‡
A	% ≥4-fold rise§	92.4	(89.5, 94.8)	A	% ≥4-fold rise§	84.6	(82.3, 86.7)
	GMT	3246	(2910, 3620)		GMT	4114	(3832, 4417)
C	% ≥4-fold rise§	88.7	(85.2, 91.5)	C	% ≥4-fold rise§	89.7	(87.8, 91.4)
	GMT	1639	(1406, 1911)		GMT	3469	(3148, 3823)
Y	% ≥4-fold rise§	80.1	(76.0, 83.8)	Y	% ≥4-fold rise§	79.4	(76.9, 81.8)
	GMT	1228	(1088, 1386)		GMT	2449	(2237, 2680)
W-135	% ≥4-fold rise§	95.3	(92.8, 97.1)	W-135	% ≥4-fold rise§	94.4	(92.8, 95.6)
	GMT	1545	(1384, 1725)		GMT	1871	(1723, 2032)

3 Both studies (11-18 and 18-55 years of age) were designed to show the safety and immunogenicity of Menactra
4 vaccine are non-inferior to that of Menomune – A/C/Y/W-135 vaccine. The table shows the immune response in
5 Menomune – A/C/Y/W-135 vaccine participants from these studies.

6 * Serum Bactericidal Assay with baby rabbit complement (SBA-BR).

7 † N = Number of participants with valid serology results at Day 0 and Day 28.

8 ‡ The 95% CI for the GMT was calculated based on an approximation to the normal distribution.

1 § The proportion of subjects with a ≥ 4 -fold rise from baseline in SBA-BR titer for *N meningitidis* for each
2 of the serogroups A, C, Y and W-135, 28 days following vaccination with Menomune – A/C/Y/W-135
3 vaccine.

4
5 In participants 11-18 years of age with undetectable pre-vaccination SBA titers (ie, < 8 at Day 0),
6 rates of seroconversion (defined as a ≥ 4 -fold rise in Day 28 SBA titers) following Menomune –
7 A/C/Y/W-135 vaccine were 100%, serogroup A (n=93/93); 99%, serogroup C (n=151/152);
8 100%, serogroup Y (n=47/47); 99%, serogroup W-135 (n=138/139).

9
10 In participants 18-55 years of age with undetectable pre-vaccination SBA titers (ie, < 8 at Day 0),
11 rates of seroconversion (defined as a ≥ 4 -fold rise in Day 28 SBA titers) following Menomune –
12 A/C/Y/W-135 vaccine were 99%, serogroup A (n=143/144); 98%, serogroup C (n=297/304);
13 97%, serogroup Y (n=221/228); 99%, serogroup W-135 (n=325/328).

14
15

1 **15. REFERENCES**

- 2
- 3 1 Centers for Disease Control and Prevention. Prevention and control of meningococcal
4 disease: recommendations of the Advisory Committee on Immunization Practices (ACIP).
5 MMWR 2005;54 (No. RR - 7): 1-21.
- 6 2 Centers for Disease Control and Prevention. Revised recommendations of the Advisory
7 Committee on Immunization Practices to vaccinate all persons aged 11-18 years with
8 meningococcal conjugate vaccine. MMWR 2007;56:794-5.
- 9 3 Centers for Disease Control and Prevention. Recommendation from the Advisory
10 Committee on Immunization Practices (ACIP) for use of quadrivalent meningococcal
11 conjugate vaccine (MCV4) in children aged 2-10 years at increased risk for invasive
12 meningococcal disease. MMWR 2007;56:1265-6.
- 13 4 Recommendations of the Advisory Committee on Immunization Practices (ACIP) General
14 Recommendations on Immunization. MMWR 2006 December 01;55(RR15):1-48.
- 15 5 Reingold AL, et al. Age-specific differences in duration of clinical protection after
16 vaccination with meningococcal polysaccharide A vaccine. Lancet. 1985;No.8447:114-118.
- 17 6 Granoff DM, et al. Meningococcal vaccines. In: Plotkin SA, Orenstein WA, Offit PA,
18 editors. Vaccines. 5th ed. Philadelphia, PA: WB Saunders Company; 2008:399-434.
- 19 7 Mueller H, et al. A protein-free medium for primary isolation of the gonococcus and
20 meningococcus. Proc Soc Exp Biol Med 1941;48:330.

- 1 8 Watson RG, et al. The specific hapten of group C (group II a) meningococcus. II. Chemical
2 nature. J Immunol. 1958;81:337.
- 3 9 Tiesjema RH, et al. Enhanced stability of meningococcal polysaccharide vaccines by using
4 lactose as a menstruum for lyophilization. Bull WHO 1977;55:43-48.
- 5 10 WHO Technical Report Series, 1981;No.658.
- 6 11 Mäkelä PH, et al. Evolution of conjugate vaccines. Expert Rev Vaccines 2002;1(3):399-
7 410.
- 8 12 Goldschneider I, et al. Human immunity to the meningococcus. I. The Role of Humoral
9 Antibodies. J Exp Med 1969;129:1307-1326.
- 10 13 Maslanka SE, et al. Standardization and a Multilaboratory Comparison of Neisseria
11 meningitidis Serogroup A and C Serum Bactericidal Assays. Clin and Diag Lab Immunol
12 1997;156-167.
- 13

1 **16. HOW SUPPLIED/STORAGE AND HANDLING**

2 **16.1. How Supplied**

3 One single dose vial of lyophilized vaccine (NDC 49281-487-58) with one 0.6 mL vial of diluent
4 (NDC 49281-466-18) (contains no preservative). Supplied as package NDC 49281-489-04.

5

6 One 10 dose vial of lyophilized vaccine (NDC 49281-488-78) with one 6.0 mL vial of diluent
7 (NDC 49281-466-31) (contains preservative). Supplied as package NDC 49281-489-10.

8

9 **16.2. Storage**

10 Store lyophilized vaccine, diluent, and reconstituted vaccine, when not in use, at 2° to 8°C (35° to
11 46°F). Do not freeze.

12

13 Do not use after the expiration date shown on the vial labels of the lyophilized vaccine and
14 diluent.

15

16 Discard remainder of reconstituted vaccine from multidose vials within 35 days after
17 reconstitution. Vaccine from single dose vials should be used immediately after reconstitution.

18

19 **17. PATIENT COUNSELING INFORMATION**

20 Before administration of Menomune – A/C/Y/W-135 vaccine, health-care providers should
21 inform the patient, parent or guardian of the benefits and risks of the vaccine. The health-care
22 provider should provide the Vaccine Information Statements, which are required by the National

1 Childhood Vaccine Injury Act of 1986 to be given with each immunization. Patients, parents, or
2 guardians should be instructed to report adverse reactions to their health-care provider.

3

4

5 Rx only

6

7 Menomune® – A/C/Y/W-135 is a registered trademark of sanofi pasteur and its subsidiaries.

8

9

10 Manufactured by:

11 **Sanofi Pasteur Inc.**

12 Swiftwater PA 18370 USA

13

14

6895

15

16

17

SANOFI PASTEUR 