

**BLUE BIRD
BOV - COCCI
Type B Medicated Liquid Cattle Feed**

For control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in cattle up to 800 lbs.

Active Drug Ingredient: Lasalocid (as Lasalocid sodium).....5 – 80,000 g/ton*

Guaranteed Analysis**

Crude Protein (Min)	_____	%
NPN ^a (Max)	_____	%
Crude Fat (Min)	_____	%
Crude Fiber (Max)	_____	%
Calcium ^a (Min)	_____	%
Calcium ^a (Max)	_____	%
Phosphorus ^a (Min)	_____	%
Potassium ^a (Min)	_____	%
Salt ^a (Min)	_____	%
Salt ^a (Max)	_____	%
Sodium ^b (Min)	_____	%
Sodium ^b (Max)	_____	%
Vitamin A ^a (Min)	_____	IU/lb

pH must be within the range of 4.0 - 8.0

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients: Ingredients as defined by AAFCO.

Mixing Directions

Mix this liquid Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for cattle.

The following table provides examples of mixing rates:

Type B lasalocid concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C lasalocid concentration (g/ton)
100	600	1400	30
1000	120	1880	60
10000	20	1980	100
20000	100	1900	1000
80000	36	1964	1440

[For liquid Type B feeds stored in recirculating tank systems: Recirculate immediately prior to use for no less than 10 minutes, moving not less than one (1) percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used. For liquid Type B feeds stored in mechanical, air or other agitation-type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.]***

The resulting Type C medicated feed should be hand fed at a rate of 1 mg of lasalocid per 2.2 lbs body weight to cattle weighing up to 800 lbs to provide a maximum of 360 mg of lasalocid per head per day.

Caution

Feeding undiluted or mixing errors or inadequate mixing (recirculation or agitation) may result in an excess lasalocid concentration which could be fatal to cattle and sheep. Do not allow horses or other equines access to Type A articles or Type B feeds containing lasalocid as ingestion may be fatal. Safety of lasalocid for use in unapproved species has not been established.

Warning

A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

Manufactured By:
Blue Bird Feed Mill
City, State, Zip

NET WEIGHT ON BAG OR BULK

*The final printed feed label should list only a single drug concentration.

**Nutrient guarantees should conform to AAFCO recommendations for the intended breed or class of cattle

***Recirculation/agitation statements not required for physically stable lasalocid liquid Type B feeds with a pH in the range of 4.0 – 8.0 containing suspending agent(s) sufficient to maintain a viscosity of not less than 300 centipoises per second for 3 months.