

**BLUE BIRD  
BOV - ROG  
Type B Medicated Liquid Pasture Cattle Feed**

For increased rate of weight gain in pasture cattle  
(slaughter, stocker, feeder cattle, and dairy and beef replacement heifers).

**Active Drug Ingredient:** Lasalocid (as Lasalocid sodium).....5 – 80,000 g/ton\*

**Guaranteed Analysis**

Crude Protein (Min).....	%
NPN <sup>a</sup> (Max).....	%
Crude Fat (Min).....	%
Crude Fiber (Max).....	%
Calcium <sup>a</sup> (Min).....	%
Calcium <sup>a</sup> (Max).....	%
Phosphorus <sup>a</sup> (Min).....	%
Potassium <sup>a</sup> (Min).....	%
Salt <sup>a</sup> (Min).....	%
Salt <sup>a</sup> (Max).....	%
Sodium <sup>b</sup> (Min).....	%
Sodium <sup>b</sup> (Max).....	%
Vitamin A <sup>a</sup> (Min).....	IU/lb

pH must be within the range of 4.0 - 8.0

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients:** Ingredients as defined by AAFCO.

**Mixing Directions**

Mix this liquid Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for cattle.

The following table provides examples of mixing rates:

Type B lasalocid concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C lasalocid concentration (g/ton)
100	200	1800	10
500	120	1880	30
1000	60	1940	30
20000	60	1940	600
80000	15	1985	600

[For liquid Type B feeds stored in recirculating tank systems: Recirculate immediately prior to use for no less than 10 minutes, moving not less than one (1) percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used. For liquid Type B feeds stored in mechanical, air or other agitation-type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.]\*\*

The resulting Type C medicated feed should be fed continuously to provide 60 to 300 mg of lasalocid per head per day. Intakes of lasalocid in excess of 200 mg per head per day have not been shown to be more effective than 200 mg per head per day. The drug must be contained in at least one pound of Type C feed.

**Caution**

Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle and sheep. Do not allow horses or other equines access to Type A articles or Type B feeds containing lasalocid as ingestion may be fatal. Safety of lasalocid for use in unapproved species has not been established.

**Warning**

A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

**Manufactured by**  
Blue Bird Feed Mill  
City, State, Zip

**NET WEIGHT ON BAG OR BULK**

\*The final printed feed label should list only a single drug concentration

\*\*Recirculation/agitation label statements not required for physically stable lasalocid liquid Type B feeds with a pH in the range of 4.0 – 8.0 containing suspending agent(s) sufficient to maintain a viscosity of not less than 300 centipoises per second for 3 months.