

Medicated Broiler Chicken Feed

(avilamycin and monensin Type C medicated feed)

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS FOR USE

For the prevention of mortality caused by necrotic enteritis associated with *Clostridium perfringens*; and as an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati* and *E. maxima* in broiler chickens.

ACTIVE INGREDIENTS

Avilamycin ¹	13.6 to 40.9 g/ton *
Monensin ² ,USP.....	90 to 110 g/ton *

GUARANTEED ANALYSIS

Crude Protein, not less than.....		%
Lysine, not less than.....		%
Methionine, not less than.....		%
Crude Fat, not less than.....		%
Crude Fiber, not more than.....		%
Calcium, not less than.....		%
Calcium, not more than.....		%
Phosphorus, not less than.....		%
Salt ³ , not less than.....		%
Salt ³ , not more than.....		%
Sodium ⁴ , not less than.....		%
Sodium ⁴ , not more than.....		%

³If added.

⁴Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed as the sole ration for 21 consecutive days. Feed to chickens that are at risk of developing, but not yet showing clinical signs of, necrotic enteritis associated with *Clostridium perfringens* in broiler chickens.

CAUTIONS

To assure responsible antimicrobial drug use in broiler chickens, treatment administration must begin on or before 10 days of age. The safety of avilamycin has not been established in chickens intended for breeding purposes. Avilamycin has not been demonstrated to be effective in broiler chickens showing clinical signs of necrotic enteritis prior to the start of medication.

The Veterinary Feed Directive (VFD) expiration date must not exceed 90 days from the VFD issuance date. VFDs for avilamycin shall not be refilled.

In the absence of coccidiosis in broiler chickens the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain. Not for replacement chickens intended to become broiler breeding chickens.

WARNINGS

Withdrawal Periods and Residue Warnings



No withdrawal period is required when used according to labeling. Do not feed in laying chickens. Do not feed to chickens over 16 weeks of age.



User Safety Warnings

Not for use in humans. Keep out of reach of children. To report adverse effects in users, to obtain more information, or to obtain a Safety Data Sheet, call 1-800-428-4441.

Animal Safety Warnings

Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal.

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Lot # _____

Expiration Date (90 days from date of manufacture) _____

NADA# 141-465, Approved by FDA
Last revision: September, 2016

*The medicated feed label must state a single drug concentration.

¹ Sourced from Intepriety™, NADA# 141-439

² Sourced from Coban™, NADA# 038-878

Elanco, Intepriety, Coban and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.