

Tylosin Medicated Feed – Swine Ileitis

Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATION

For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis*

ACTIVE DRUG INGREDIENT

Tylosin (as tylosin phosphate) ^a40 or 100 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%
Selenium, not less than.....	_____	ppm
Zinc, not less than.....	_____	ppm

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

CAUTION

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

FEEDING DIRECTIONS

Feed 100 g of tylosin per ton of complete feed for 21 days. Alternatively, feed 100 g of tylosin per ton of complete feed for at least three weeks, followed by 40 g tylosin per ton of complete feed until pigs reach market weight. Feed continuously as the sole ration.

MANUFACTURED BY:
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

^a Sourced from Tylan[®], NADA# 12-491

*The medicated feed label must state a single drug concentration.

Elanco, Tylan and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.