

**Chlortetracycline Turkey Ration I  
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For the control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline.

**ACTIVE DRUG INGREDIENT**

Chlortetracycline.....200 g/ton

**GUARANTEED ANALYSIS**

Crude Protein, not less than.....%  
Crude Fat, not less than.....%  
Crude Fiber, not more than.....%  
Lysine (min).....%  
Methionine (min).....%  
Calcium (min).....%  
Calcium (max).....%  
Phosphorus(min).....%  
Salt<sup>1</sup> (min).....%  
Salt<sup>1</sup> (max).....%  
Sodium<sup>2</sup> (min).....%  
Sodium<sup>2</sup> (max).....%

<sup>1</sup>If added

<sup>2</sup>Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

**INGREDIENTS**

Ingredients as defined by AAFCO

**DIRECTIONS FOR USE**

Feed continuously for 7 to 14 days as the sole ration.

**WARNING:** Do not feed to turkeys producing eggs for human consumption.  
Zero-day withdrawal period.

Net Weight \_\_\_\_\_ lb. ( \_\_\_\_\_ kg.)

Manufactured by  
Blue Bird Feed Company  
Blue bird, MD 00000