

Tylosin/Pyrantel Medicated Feed - Swine Dysentery TUHDW HQMDQG & RQWRO Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS

For treatment and control of swine dysentery (vibrionic) immediately after medicating with tylosin in drinking water; aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; and as an aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections.

ACTIVE INGREDIENTS

Tylosin phosphate	40 or 100 g/ton*
Pyrantel tartrate	0.0106% (96 g/ton)

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____ %
Lysine, not less than.....	_____ %
Crude Fat, not less than.....	_____ %
Crude Fiber, not more than.....	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than.....	_____ %
Phosphorus, not less than.....	_____ %
Salt ¹ , not less than.....	_____ %
Salt ¹ , not more than.....	_____ %
Sodium ² , not less than.....	_____ %
Sodium ² , not more than.....	_____ %
Selenium, not less than.....	_____ ppm
Zinc, not less than.....	_____ ppm

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

CAUTION

Do not use in any finished feed (supplement, concentrate or complete feed) containing bentonite.

FEEDING DIRECTIONS

Feed 40 or 100 g of tylosin per ton of complete feed continuously as the sole ration for two to six weeks after treatment with tylosin in drinking water (250 mg tylosin per gallon) for 3 to 10 days.

WARNING: WITHDRAW 24 HOURS PRIOR TO SLAUGHTER

MANUFACTURED BY:
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Expiration Date: _____

*The medicated feed label must state one of the two drug concentrations.