

# Tylosin/Pyrantel Medicated Feed - Swine Dysentery

## Type C Medicated Feed A

**CAUTION:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

### INDICATIONS FOR USE

For prevention of swine dysentery (vibrionic); aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; and as an aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections.

### ACTIVE INGREDIENTS

Tylosin phosphate..... 40 g/ton  
Pyrantel tartrate..... 0.0106% (96 g/ton)

### GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt <sup>1</sup> , not less than.....	_____	%
Salt <sup>1</sup> , not more than.....	_____	%
Sodium <sup>2</sup> , not less than.....	_____	%
Sodium <sup>2</sup> , not more than.....	_____	%
Selenium, not less than.....	_____	ppm
Zinc, not less than.....	_____	ppm

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

### INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

### CAUTION

Do not use in any finished feed (supplement, concentrate or complete feed) containing bentonite.

### FEEDING DIRECTIONS

Feed continuously as the sole ration until market weight. This feed is for use following the feeding of 100 g per ton tylosin for at least 3 weeks.

### WARNING: WITHDRAW 24 HOURS PRIOR TO SLAUGHTER

MANUFACTURED BY:  
BLUE BIRD FEED MILL  
Any Town, USA 12345

**Net Weight lb (kg) on bag or bulk**

Expiration date: \_\_\_\_\_