

OXYTETRACYCLINE  
TYPE B  
BAG OR BULK

BLUE BIRD  
LOBSTER FEED  
MEDICATED

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

<b>INDICATIONS FOR USE</b>
For control of gaffkemia caused by <i>Aerococcus viridans</i> .

**ACTIVE DRUG INGREDIENT**

Oxytetracycline.....20 g/lb

**GUARANTEED ANALYSIS**

Crude Protein (Min).....%  
Crude Fat (Min).....%  
Crude Fiber (Max).....%  
Phosphorus (Min).....%

**INGREDIENTS**

Ingredients as defined by AAFCO.

**MIXING DIRECTIONS** – Mix 100 lbs. of this Type B with 1900 lbs of feed to manufacture one ton of Type C feed containing 1 g oxytetracycline/lb of feed. Feed the resulting Type C medicated feed as the sole ration for 5 consecutive days.

**► WARNING: Withdraw medicated feed 30 days before harvesting lobsters. ◀**

**FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS.**

\_\_\_lbs(\_\_\_kg) NET WEIGHT

BLUE BIRD FEED MILL  
ROBIN, IN 00000