

LINCOMYCIN - PYRANTEL TARTRATE
TYPE C
BLUE BIRD
SWINE FEED #6
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For treatment and/or control of swine dysentery; for removal and control of large roundworm (*Ascaris suum*) and nodular worm (*Oesophagostomum spp.*) infections.

ACTIVE DRUG INGREDIENT

Lincomycin (lincomycin hydrochloride) 40 g/ton
Pyrantel tartrate 800 g/ton (0.0881%)

GUARANTEED ANALYSIS

Crude Protein (Min) %
Lysine (Min) %
Crude Fat (Min) %
Crude Fiber (Max) %
Calcium (Min) %
Calcium (Max) %
Phosphorus (Min) %
Salt, if added (Min) %
Salt, if added (Max) %
Sodium (Min)* %
Sodium (Max)* %
Selenium (Min) ppm
Zinc (Min) ppm

* Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed as a single therapeutic treatment.
Feed at a rate of 1 lb of feed per 40 lb of body weight for animals up to 200 lb, and 5 lb of feed per head for animals over 200 lb.

CAUTION: Not to be fed to swine that weigh more than 250 pounds. Do not mix in feeds containing bentonite. Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within five to eight days without discontinuing the lincomycin treatment. The effects of lincomycin on swine reproductive performance, pregnancy, and lactation have not been determined. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

WARNING: WITHDRAW 24 HOURS PRIOR TO SLAUGHTER.

_____ lbs (_____ kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000

Expiration Date: _____