

50 LBS. NET WEIGHT

**TYPE C MEDICATED FEED
BLUE BIRD
SWINE FEED
MEDICATED**

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

FENBENDAZOLE IS INDICATED FOR THE REMOVAL OF:

Lungworms: *Metastrongylus apri*, *Metastrongylus pudendotectus*

Gastrointestinal worms: Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworm (*Ascaris suum*); nodular worms (*Oesophagostomum dentatum*, *O. quadrispinulatum*); small stomach worm (*Hyostromylus rubidus*); Adult and larvae (L2, 3, 4 stages - intestinal mucosal forms) whipworm (*Trichuris suis*)

Kidney worm: Adult and larvae *Stephanurus dentatus*.

LINCOMYCIN IS INDICATED FOR: control of swine dysentery; for use in swine on premises with a history of swine dysentery, but where symptoms have not yet occurred.

Active Drug Ingredient

Fenbendazole	10.0 - 80.0 grams per ton
Lincomycin	40 grams per ton

Guaranteed analysis

Crude Protein,	not less than _____%
Crude Fat	not less than _____%
Crude Fiber	not less than _____%

Ingredients

Ground corn	Riboflavin supplement
Ground milo	Calcium pantothenate
Corn gluten meal	Niacin, Vit. B12 supplement
Wheat middlings	Vit. A & D3
Dehydrated alfalfa meal	Ground limestone 1% Salt ½%
Fish meal	Manganese sulfate 0.02%
Animal fat with BHT	Deflourinated phosphate ½%
Soybean oil meal	Potassium iodide 0.002%

Feeding Directions

Calculate the amount of THIS feed that the pigs to be treated should eat within 3 to 12 days to get a total dose of 9 mg of fenbendazole per kilogram body weight (4.08 mg fenbendazole per pound body weight). Feed as the sole feed the calculated amount. When it is fully consumed resume normal feeding.

Lincomycin is combined with fenbendazole in this Type C feed at 40 grams of lincomycin per ton of feed.

CAUTION: Occasionally, swine fed lincomycin may, within the first 2 days after the onset of treatment, develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within 5-8 days without discontinuing the lincomycin treatment. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects. The effects of lincomycin on swine reproductive performance, pregnancy, and lactation have not been determined.

SPECIAL CONSIDERATION: Feeds containing lincomycin are not to be fed to swine that weigh more than 250 lb.

Manufactured by
Blue Bird Feed Mills
Robin, Indiana

June 9, 2016