

OXYTETRACYCLINE  
TYPE B  
BAG OR BULK

BLUE BIRD  
SALMONID and CATFISH FEED

MEDICATED

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

ACTIVE DRUG INGREDIENT

Oxytetracycline.....20 g/lb

INDICATIONS FOR USE
For Salmonids  For control of ulcer disease caused by <i>Hemophilus piscium</i> , furunculosis caused by <i>Aeromonas salmonicida</i> , bacterial hemorrhagic septicemia caused by <i>Aeromonas liquefaciens</i> and pseudomonas disease. <b>► WARNING: Do not liberate or slaughter salmonids for food during treatment or for 21 days following last feeding of medicated feed. ◀</b>
For Catfish  For control of bacterial hemorrhagic septicemia caused by <i>Aeromonas liquefaciens</i> and pseudomonas disease. <b>► WARNING: Do not liberate or slaughter catfish for food during treatment or for 21 days following last feeding of medicated feed. Do not use when water temperature is below 62°F (16.7°C). ◀</b>

GUARANTEED ANALYSIS

Crude Protein (Min).....%  
Crude Fat (Min).....%  
Crude Fiber (Max).....%  
Phosphorus (Min).....%

INGREDIENTS

Ingredients as defined by AAFCO.

**MIXING DIRECTIONS – Salmonids and Catfish**

Note that feeding rates may vary depending on the size and health of the fish. Feeding rates should be chosen to ensure that all fish in the rearing unit are adequately medicated. Use the table below to calculate the amount of Type B feed to prepare Type C feed to use at your feeding rate. Feed oxytetracycline in a mixed ration for 10 days.

Feeding Rate*	To achieve a dose of 2.5 – 3.75 g oxytetracycline/100 pounds of fish:		
pounds/100 pounds (%)	Oxytetracycline in Type C Medicated Feed (g/ton)	Pounds of Type B Medicated Feed per ton of feed	Pounds of total biomass that one ton of Type C Medicated Feed will treat**
1	5,000 – 7,500	250.0 – 375.0	200,000
2	2,500 – 3,750	125.0 – 187.5	100,000
3	1,667 – 2,500	83.3 – 125.0	66,667
4	1,250 – 1,875	62.5 – 93.8	50,000
5	1,000 – 1,500	50.0 – 75.0	40,000
6	833 – 1,250	41.7 – 62.5	33,333
7	714 – 1,071	35.7 – 53.6	28,571
8	625 – 938	31.3 – 46.9	25,000
9	556 – 833	27.8 – 41.7	22,222
10	500 – 750	25.0 – 37.5	20,000
15	333 – 500	16.7 – 25.0	13,333

\*To calculate grams oxytetracycline per ton finished feed at other feed rates: (desired dose rate/feeding rate) x 2000

To calculate Type B per ton at other rated rates: lb Type B per ton finished feed/20

Example: at a feeding rate of 2.5% (2.5 pounds per 100 pounds), with a desired dose of 3.75 g/100 pounds:

3.75 per 100 lb/2.5 lb per 100 lb = 1.5 g per lb x 2000 lb = 3000 g oxytetracycline/ton

3000 g oxytetracycline per ton/20 g oxytetracycline per lb of Type B = 150 pounds Type B per ton of feed

\*\* To calculate total required medicated feed for the 10 day treatment period, multiply total estimated rearing unit biomass by 10 and divide by the pounds of medicated feed needed at your feeding rate

Example: at a feeding rate of 2% in a unit containing 5,000 pounds of fish biomass:

5,000 lb x 10 = 50,000 lb/100,000 lb per ton = 0.5 tons

**FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS.**

\_\_\_lbs(\_\_\_kg) NET WEIGHT

**BLUE BIRD FEED MILL  
ROBIN, IN 00000**