

OXYTETRACYCLINE – SALINOMYCIN SODIUM  
 TYPE C  
 BLUE BIRD  
 CHICKEN FEED  
 MEDICATED

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For chickens as an aid in the reduction of mortality due to air-sacculitis (air sac infection) caused by *Escherichia coli* sensitive to oxytetracycline and for the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati* and *E. maxima*.

**ACTIVE DRUG INGREDIENT**

Oxytetracycline .....500 g/ton  
 Salinomycin Sodium .....40 - 60 g/ton\*

**GUARANTEED ANALYSIS**

Crude Protein (Min).....	_____	%
Crude Lysine (Min).....	_____	%
Methionine (Min).....	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Min).....	_____	%
Calcium (Min).....	_____	%
Calcium (Max).....	_____	%
Phosphorus (Min).....	_____	%
Salt, if added (Min).....	_____	%
Salt, if added (Max).....	_____	%
Total Sodium (Min).....	_____	%
Total Sodium (Max).....	_____	%

(Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee)

**INGREDIENTS**

Ingredients as defined by AAFCO

**FEEDING DIRECTIONS**

Feed for 5 days as sole ration. Treat at first clinical signs of disease.

**WARNING: DO NOT FEED TO LAYING CHICKENS. DO NOT USE IN LOW-CALCIUM FEED CONTAINING LESS THAN 0.55% DIETARY CALCIUM. USE IN SUCH FEEDS MAY RESULT IN VIOLATIVE RESIDUES. WITHDRAW 24 HOURS BEFORE SLAUGHTER.**

**CAUTION: Do not use in feeds containing pellet binders. May be fatal if accidentally fed to adult turkeys or to horses.**

\_\_\_\_\_ lbs (\_\_\_\_\_kg) NET WEIGHT

BLUE BIRD FEED MILL  
 Robin, IN 00000

\* Final printed label on formulated Type C medicated feed must bear a single concentration of drug.