

OXYTETRACYCLINE  
TYPE C  
BAG OR BULK

BLUE BIRD  
SALMONID and CATFISH FEED  
MEDICATED

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

<b>INDICATIONS FOR USE</b>
For Salmonids  For control of ulcer disease caused by <i>Hemophilus piscium</i> , furunculosis caused by <i>Aeromonas salmonicida</i> , bacterial hemorrhagic septicemia caused by <i>Aeromonas liquefaciens</i> and pseudomonas disease.
<b>► WARNING: Do not liberate or slaughter salmonids for food during treatment or for 21 days following last feeding of medicated feed. ◀</b>
For Catfish  For control of bacterial hemorrhagic septicemia caused by <i>Aeromonas liquefaciens</i> and pseudomonas disease.
<b>► WARNING: Do not liberate or slaughter catfish for food during treatment or for 21 days following last feeding of medicated feed. Do not use when water temperature is below 62°F (16.7°C). ◀</b>

ACTIVE DRUG INGREDIENT

Oxytetracycline.....333 to 7,500 g/ton<sup>1</sup>

GUARANTEED ANALYSIS

Crude Protein (Min).....%  
Crude Fat (Min).....%  
Crude Fiber (Max).....%  
Phosphorus (Min).....%

INGREDIENTS

Ingredients as defined by AAFCO.

**FEEDING DIRECTIONS – Salmonids and Catfish**

Note that feeding rates may vary depending on the size and health of the fish. Feeding rates should be chosen to ensure that all fish in the rearing unit are adequately medicated. Use the table below to calculate the amount of Type C to use at your feeding rate. Feed for 10 days.

Feeding Rate <sup>2</sup> pounds/100 pounds (%)	To achieve a dose of 2.5 – 3.75 g oxytetracycline/100 pounds of fish Oxytetracycline in Finished Feed g/ton	Total biomass that one ton of medicated feed will treat <sup>3</sup> (pounds)
1	5,000 – 7,500	200,000
2	2,500 – 3,750	100,000
3	1,667 – 2,500	66,667
4	1,250 – 1,875	50,000
5	1,000 – 1,500	40,000
6	833 – 1,250	33,333
7	714 – 1,071	28,571
8	625 – 938	25,000
9	556 – 833	22,222
10	500 – 750	20,000
15	333 – 500	13,333

<sup>1</sup>The final printed label must include only a single drug concentration

<sup>2</sup>To calculate grams oxytetracycline per ton finished feed at other feed rates: (desired dose rate/feeding rate) x 2000

Example: at a feeding rate of 2.5% (2.5 pounds per 100 pounds), with a desired dose of 3.75 g/100 pounds:  
 $3.75 \text{ per } 100 \text{ lb} / 2.5 \text{ lb per } 100 \text{ lb} = 1.5 \text{ g per lb} \times 2000 \text{ lb} = 3000 \text{ g oxytetracycline/ton}$

<sup>3</sup>To calculate total required medicated feed for the 10 day treatment period, multiply total estimated rearing unit biomass by 10 and divide by the pounds of medicated feed needed at your feeding rate

Example: at a feeding rate of 2% in a unit containing 5,000 pounds of fish biomass:  
 $5,000 \text{ lb} \times 10 = 50,000 \text{ lb} / 100,000 \text{ lb per ton} = 0.5 \text{ tons}$

**FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS.**

\_\_\_lbs(\_\_\_kg) NET WEIGHT

**BLUE BIRD FEED MILL  
ROBIN, IN 00000**