

Sulfadimethoxine and Ormetoprim^a Poultry Concentrate
Type B Medicated Feed
(sulfadimethoxine and ormetoprim Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Chickens [broiler and replacement (breeder and layer)]: As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*; as an aid in the prevention of bacterial infections caused by *Haemophilus gallinarum* (infectious coryza), *Escherichia coli* (colibacillosis) and *Pasteurella multocida* (fowl cholera).

Turkeys: As an aid in the prevention of coccidiosis caused by *Eimeria adenoides*, *E. gallopavonis*, and *E. meleagrimitis*; for the prevention of bacterial infections caused by *Pasteurella multocida* (fowl cholera).

Ducks, including breeding ducks: As an aid in the control of bacterial infections due to *Pasteurella multocida* (fowl cholera) in ducks including breeding ducks.

Ducks: As an aid in the control of bacterial infections due to *Escherichia coli*, *Riemerella anatipestifer* (strains 2, 3, & 4) and severe challenge of *P. multocida* (fowl cholera).

Chukar Partridges: For the prevention of coccidiosis caused by *Eimeria kofoidi* and *E. legionensis*.

Active Drug Ingredients^a

Sulfadimethoxine.....	1135 – 11,350 g/ton**	
	[0.0.5675 g/lb (0.125%) – 5.675 g/lb (1.25%)]	
Ormetoprim	681 – 6,810 g/ton**	
	[0.3405 g/lb (0.075%) – 3.405 g/lb (0.75%)]	

**The final printed feed label should list only a single drug concentration.

GUARANTEED ANALYSIS

Crude Protein, Min	_____ %
Lysine, Min	_____ %
Methionine, Min	_____ %
Crude Fat, Min	_____ %
Crude Fiber, Max	_____ %
Calcium, Min	_____ %
Calcium, Max	_____ %
Phosphorus, Min	_____ %
Salt ¹ , Min	_____ %
Salt ¹ , Max	_____ %
Sodium ² , Min	_____ %
Sodium ² , Max	_____ %

¹ Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

² Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Ingredients as defined by AAFCO

Directions for Use

Chickens: Thoroughly mix 20 – 200 lbs. of this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for broiler and replacement chickens (breeder and layer) containing 113.5 g/ton (0.0125%) sulfadimethoxine and 68.1 g/ton (0.0075%) ormetoprim. Feed the resulting Type C medicated feed continuously as the sole ration.

Turkeys: Thoroughly mix 10 to 100 lbs. of this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for turkeys containing 56.75 g/ton (0.00625%) sulfadimethoxine and 34.05 g/ton (0.00375%) ormetoprim. Feed the resulting Type C medicated feed continuously as the sole ration.

Ducks:

As an aid in the control of *P. multocida* (fowl cholera) in ducks, including breeding ducks, thoroughly mix 40 to 400 lbs. of this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed containing 227 g/ton (0.025%) sulfadimethoxine and 136.2 g/ton (0.015%) ormetoprim. Feed the resulting Type C medicated feed for 7 days as the sole ration.

As an aid in the control of *E. coli*, *R. anatipestifer* and severe challenge of *P. multocida* (fowl cholera), thoroughly mix 80 to 800 lbs. of this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed containing 454 g/ton (0.05%) sulfadimethoxine and 272.4 g/ton (0.03%) ormetoprim. Feed the resulting Type C medicated feed for 7 days as the sole ration.

Chukar partridges: Thoroughly mix 20 to 200 of this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of a Type C medicated feed containing 113.5 g/ton (0.0125%) sulfadimethoxine and 68.1 g/ton (0.0075%) ormetoprim. Feed the resulting Type C medicated feed continuously as the sole ration to young birds up to 8 weeks of age.

Caution

The safety of feeding sulfadimethoxine and ormetoprim at levels higher than 0.04% (total drug) in breeding ducks has not been established. For control of duck disease, medication should be started at the first sign of infection. The safety of feeding RofenAid to Chukar partridge breeding stock has not been established.

Warning

Do not feed to turkeys and ducks producing eggs for food. Withdraw 5 days prior to slaughter of broiler and replacement chickens, turkeys, and ducks. Do not feed to chickens over 16 weeks (112 days) of age.

Lot Number (if applicable): _____

MANUFACTURED BY
Blue Bird Feed Company
City, State, Zip

NET WEIGHT ON BAG OR BULK

^aRofenAid® 40 is the proprietary name of sulfadimethoxine and ormetoprim formulation Type A medicated article (NADA 040-209).

Zoetis Inc. [7 March 2016]