

**Type C Medicated Cattle Supplement M-T+**  
**Type C Medicated Feed**  
(melengestrol acetate and tylosin Type C top-dress medicated feed)

**For Use in Heifer Feeds Only**

**Caution:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

**INDICATIONS**

Heifers being fed in confinement for slaughter: For increased rate of weight gain, improved feed efficiency, suppression of estrus (heat), and for reduction of incidence of liver abscesses associated with *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*.

**ACTIVE DRUG INGREDIENTS**

Melengestrol acetate (MGA<sup>®</sup>)<sup>a</sup>.....0.0000276 to 0.00022% (0.25 to 2 g/ton)\*  
Tylosin (as tylosin phosphate) (Tylan<sup>™</sup>)<sup>b</sup>.....90 to 360 g/ton\*

**GUARANTEED ANALYSIS**

Crude Protein, not less than.....	_____	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than .....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt <sup>2</sup> , not less than.....	_____	%
Salt <sup>2</sup> , not more than.....	_____	%
Sodium <sup>3</sup> , not less than.....	_____	%
Sodium <sup>3</sup> , not more than.....	_____	%
Potassium, not less than.....	_____	%
Vitamin A <sup>2,4</sup> , not less than.....	_____	I.U./lb

<sup>1</sup>When added.

<sup>2</sup>If added.

<sup>3</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup>Other than precursors of Vitamin A.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**FEEDING DIRECTIONS**

Each pound contains 0.125 to 1.0 mg melengestrol acetate and 45 to 180 mg of tylosin. Feed to heifers at a rate of 0.5 to 2.0 pounds per head per day to provide 0.25 to 0.5 mg melengestrol acetate and 60 to 90 mg tylosin per head per day. Prior to feeding, this Type C product must be top-dressed onto a complete feed or mixed into the amount of complete feed consumed by an animal per day.

**Caution**

Feed continuously as sole ration.

MGA is for use only in heifers being fed in confinement for slaughter. Not effective in steers and spayed heifers.

---

**Warning:** No withdrawal period is required.

When mixing and handling Tylan™, use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Lot Number (if applicable): \_\_\_\_\_

**MANUFACTURED BY**

Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

\*Final printed label on formulated Type C medicated feed must bear a single concentration for each drug.

<sup>a</sup>MGA® is the proprietary name of melengestrol acetate Type A medicated article (NADA 039-402).

<sup>b</sup>Tylan™ is the proprietary name of tylosin phosphate Type A medicated article (NADA 012-491).

Zoetis Inc. [7 September 2016]