

**Sulfaquinoxaline CC 0.1  
Chicken Ration  
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

As an aid in controlling outbreaks of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, and *E. brunetti*.

**ACTIVE DRUG INGREDIENT**

Sulfaquinoxaline ..... 909 g/ton (0.1%)

**GUARANTEED ANALYSIS**

Crude Protein, not less than .....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than .....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than .....	_____	%
Calcium, not less than .....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt <sup>1</sup> , not less than.....	_____	%
Salt <sup>1</sup> , not more than.....	_____	%
Sodium <sup>2</sup> , not less than.....	_____	%
Sodium <sup>2</sup> , not more than.....	_____	%

<sup>1</sup> If added.

<sup>2</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**DIRECTIONS FOR USE**

Feed for 48 to 72 hours. Skip 3 days and medicate with 0.05% Sulfaquinoxaline for 2 days, skip 3 days, and medicate with 0.05% for another 2 days. If bloody droppings recur, feed 0.05% for another 2 days.

**CAUTION:** For control of outbreaks of disease, medication should be initiated as soon as the diagnosis is determined. Medicated chickens must actually consume enough medicated feed which provides a recommended dose of approximately 3.5 to 60 mg/lb/day in chickens, depending upon the age, class of animal, ambient temperature, and other factors. Prolonged administration of sulfaquinoxaline at higher doses may result in depressed feed intake, deposition of sulfaquinoxaline crystals in the kidney, and interference with normal blood clotting.

▶ **WARNING:** Do not treat within 10 days of slaughter. Do not medicate chickens producing eggs for human consumption. ◀

**MANUFACTURED BY  
BLUE BIRD FEED MILL  
Anytown, USA 12345**

Net weight \_\_\_lb (\_\_\_kg) on bag or bulk