

**Sulfaquinoxaline CP-AE
Chicken Ration
Type C Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

As an aid in preventing outbreaks of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, and *E. brunetti* under average conditions of exposure.

ACTIVE DRUG INGREDIENT

Sulfaquinoxaline 136 g/ton (0.015%)

GUARANTEED ANALYSIS

Crude Protein, not less than	_____ %
Lysine, not less than	_____ %
Methionine, not less than	_____ %
Crude Fat, not less than	_____ %
Crude Fiber, not more than	_____ %
Calcium, not less than	_____ %
Calcium, not more than	_____ %
Phosphorus, not less than	_____ %
Salt ¹ , not less than	_____ %
Salt ¹ , not more than	_____ %
Sodium ² , not less than	_____ %
Sodium ² , not more than	_____ %

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

DIRECTIONS FOR USE

Feed continuously from the time birds are placed on litter and continue past the age when coccidiosis is ordinarily a hazard. Do not change litter unless absolutely necessary.

CAUTION: If death losses caused by coccidiosis exceed 0.5% in a 2-day period, use the sulfaquinoxaline levels recommended for control of outbreaks, returning to the original prevention dosage schedule after the outbreak has subsided. Losses may result from intercurrent disease, other conditions affecting drug intake, or variant strains of coccidia species which can contribute to the virulence of coccidiosis under field conditions.

Medicated chickens must actually consume enough medicated feed which provides a recommended dose of approximately 3.5 to 60 mg/lb/day in chickens, depending upon the age, class of animal, ambient temperature, and other factors. Prolonged administration of sulfaquinoxaline at higher doses may result in depressed feed intake, deposition of sulfaquinoxaline crystals in the kidney, and interference with normal blood clotting.

▶ **WARNING:** Do not treat within 10 days of slaughter. Do not medicate chickens producing eggs for human consumption. ◀

**MANUFACTURED BY
BLUE BIRD FEED MILL
Anytown, USA 12345**

Net weight ___lb (___kg) on bag or bulk