

**Sulfaquinoxaline CP-SE  
Chicken Ration  
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

As an aid in preventing outbreaks of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, and *E. brunetti* where excessive exposure to coccidia is increased due to overcrowding or other management factors.

**ACTIVE DRUG INGREDIENT**

Sulfaquinoxaline ..... 159 g/ton (0.0175%)

**GUARANTEED ANALYSIS**

Crude Protein, not less than .....	_____ %
Lysine, not less than .....	_____ %
Methionine, not less than .....	_____ %
Crude Fat, not less than .....	_____ %
Crude Fiber, not more than .....	_____ %
Calcium, not less than .....	_____ %
Calcium, not more than .....	_____ %
Phosphorus, not less than .....	_____ %
Salt <sup>1</sup> , not less than .....	_____ %
Salt <sup>1</sup> , not more than .....	_____ %
Sodium <sup>2</sup> , not less than .....	_____ %
Sodium <sup>2</sup> , not more than .....	_____ %

<sup>1</sup> If added.

<sup>2</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**DIRECTIONS FOR USE**

Feed continuously from the time birds are placed on litter and continue past the age when coccidiosis is ordinarily a hazard. Do not change litter unless absolutely necessary.

**CAUTION:** If death losses caused by coccidiosis exceed 0.5% in a 2-day period, use the sulfaquinoxaline levels recommended for control of outbreaks, returning to the original prevention dosage schedule after the outbreak has subsided. Losses may result from intercurrent disease, other conditions affecting drug intake, or variant strains of coccidia species which can contribute to the virulence of coccidiosis under field conditions.

Medicated chickens must actually consume enough medicated feed which provides a recommended dose of approximately 3.5 to 60 mg/lb/day in chickens, depending upon the age, class of animal, ambient temperature, and other factors. Prolonged administration of sulfaquinoxaline at higher doses may result in depressed feed intake, deposition of sulfaquinoxaline crystals in the kidney, and interference with normal blood clotting.

▶ **WARNING:** Do not treat within 10 days of slaughter. Do not medicate chickens producing eggs for human consumption. ◀

**MANUFACTURED BY  
BLUE BIRD FEED MILL  
Anytown, USA 12345**

Net weight \_\_\_lb (\_\_\_kg) on bag or bulk