Sulfaquinoxaline AFC FT 0.1 Chicken Ration Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

As an aid in the control of acute fowl cholera caused by *Pasteurella multocida* susceptible to sulfaquinoxaline and fowl typhoid caused by *Salmonella gallinarum* susceptible to sulfaquinoxaline.

ACTIVE DRUG INGREDIENT Sulfaquinoxaline 909 g/ton (0.1%) GUARANTEED ANALYSIS Crude Protein, not less than % Lysine, not less than % Methionine, not less than % Crude Fat, not less than % Crude Fiber, not more than % Calcium, not less than % Calcium, not more than % Phosphorus, not less than % Salt¹, not less than % Salt¹, not more than % Sodium², not less than % Sodium², not more than % Sodium², not more than %

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

DIRECTIONS FOR USE

Feed for 48 to 72 hours. Mortality should be brought under control. After medication, move birds to clean ground or to a clean house.

CAUTION: For control of outbreaks of disease, medication should be initiated as soon as the diagnosis is determined. Medicated chickens must actually consume enough medicated feed which provides a recommended dose of approximately 3.5 to 60 mg/lb/day in chickens, depending upon the age, class of animal, ambient temperature, and other factors. Prolonged administration of sulfaquinoxaline at higher doses may result in depressed feed intake, deposition of sulfaquinoxaline crystals in the kidney, and interference with normal blood clotting.



WARNING: Do not treat within 10 days of slaughter. Do not medicate chickens producing eggs for human consumption.

•

MANUFACTURED BY BLUE BIRD FEED MILL Anytown, USA 12345

Net weight ___lb (___kg) on bag or bulk

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.