

BLUE BIRD LABELING
chlortetracycline hydrochloride
Type B Medicated Turkey Feed – HEX

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of hexamitiasis caused by *Hexamita meleagridis* susceptible to chlortetracycline in turkeys.

Active Drug Ingredient

Chlortetracycline* 401 – 80,000 g/ton**

Guaranteed Analysis

Crude Protein (Min) _____ %
 Lysine (Min) _____ %
 Methionine (Min) _____ %
 Crude Fat (Min) _____ %
 Crude Fiber (Max) _____ %
 Calcium (Min) _____ %
 Calcium (Max) _____ %
 Phosphorus (Min) _____ %
 Salt^a (Min) _____ %
 Salt^a (Max) _____ %
 Sodium^b (Min) _____ %
 Sodium^b (Max) _____ %

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Ingredients as defined by AAFCO.

Mixing Directions

Thoroughly mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for turkeys containing 400 grams chlortetracycline per ton.

The following table provides examples of mixing rates:

Type B CTC concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C CTC concentration (g/ton)
10,000 (5 g/lb)	80	1920	400
20,000 (10 g/lb)	40	1960	400
40,000 (20 g/lb)	20	1980	400
80,000 (40 g/lb)	10	1990	400

The resulting Type C medicated feed should be fed continuously for 7 to 14 days.

Warning

Do not feed to turkeys producing eggs for human consumption.

Manufactured By:

Blue Bird Feed Mill

City, State, Zip

NET WEIGHT ON BAG OR BULK

*Deracin is the trade name of chlortetracycline hydrochloride formulation (Type A Medicated Article).

**The final printed feed label should list only a single drug concentration.