

**Sulfaquinoxaline TCP
Turkey Ration
Type C Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

As an aid in preventing outbreaks of coccidiosis caused by *Eimeria meleagritidis* and *E. adenoeides*.

ACTIVE DRUG INGREDIENT

Sulfaquinoxaline 159 g/ton (0.0175%)

GUARANTEED ANALYSIS

Crude Protein, not less than	_____ %
Lysine, not less than.....	_____ %
Methionine, not less than	_____ %
Crude Fat, not less than.....	_____ %
Crude Fiber, not more than	_____ %
Calcium, not less than	_____ %
Calcium, not more than.....	_____ %
Phosphorus, not less than.....	_____ %
Salt ¹ , not less than	_____ %
Salt ¹ , not more than.....	_____ %
Sodium ² , not less than.....	_____ %
Sodium ² , not more than.....	_____ %

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

DIRECTIONS FOR USE

Feed continuously during time birds are closely confined. May be continued for week to 10 days after flock is transferred to range to reduce danger of an outbreak following moving of the flock. Do not change litter unless absolutely necessary.

CAUTION: Medicated turkeys must actually consume enough medicated feed which provides a recommended dose of approximately 2.5 to 100 mg/lb/day in turkeys, depending upon the age, class of animal, ambient temperature, and other factors. Prolonged administration of sulfaquinoxaline at higher doses may result in depressed feed intake, deposition of sulfaquinoxaline crystals in the kidney, and interference with normal blood clotting.

▶ **WARNING:** Do not treat within 10 days of slaughter. Do not medicate turkeys producing eggs for human consumption. ◀

**MANUFACTURED BY
BLUE BIRD FEED MILL
Anytown, USA 12345**

Net weight ___lb (___kg) on bag or bulk