

**Sulfaquinoxaline RP
Growing Rabbit Ration
Type C Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

As an aid in preventing coccidiosis caused by *Eimeria stiedae* in rabbits.

ACTIVE DRUG INGREDIENT

Sulfaquinoxaline 227 g/ton (0.025%)

GUARANTEED ANALYSIS

Crude Protein, not less than	_____ %
Crude Fat, not less than	_____ %
Crude Fiber ¹ , not more than	_____ %
Crude Fiber ¹ , not less than	_____ %
Calcium ² , not less than	_____ %
Calcium ² , not more than	_____ %
Phosphorus ² , not less than	_____ %
Salt ² , not less than	_____ %
Salt ² , not more than	_____ %
Sodium ³ , not less than	_____ %
Sodium ³ , not more than	_____ %
Vitamin A ² , not less than	_____ I.U./lb

¹ The maximum crude fiber shall not exceed the minimum by more than 5.0 units.

² If added.

³ Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Start treatment after weaning. Feed continuously for 30 days or feed medicated feed for 2 days out of every week until marketing.

CAUTION: Medicated rabbits must actually consume enough medicated feed which provides a recommended dose of approximately 2.8 to 68 mg/lb/day depending upon the age, class of animal, ambient temperature, and other factors. Prolonged administration of sulfaquinoxaline at higher doses may result in depressed feed intake, deposition of sulfaquinoxaline crystals in the kidney, and interference with normal blood clotting.

▶ **WARNING:** Do not treat within 10 days of slaughter. ◀

MANUFACTURED BY

BLUE BIRD FEED MILL
Anytown, USA 12345

Net weight ____ lb (____ kg) bag or bulk