

BLUE BIRD CTC*
Type B Cattle Feed/Bacterial enteritis/Bacterial pneumonia
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Calves, Beef and Non-Lactating Dairy Cattle: For treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to Chlortetracycline.

ACTIVE DRUG INGREDIENT

Chlortetracycline 2-40 g/lb **

GUARANTEED ANALYSIS

Crude Protein, Min..... %
 NPN, Max %
 Crude Fat, Min..... %
 Crude Fiber, Min..... %
 Calcium, Min..... %
 Calcium, Max..... %
 Phosphorus, Min..... %
 Potassium, Min..... %
 Salt, Min (if added) %
 Salt, Max (if added) %
 Sodium, Min (if exceeding salt guarantee)..... %
 Sodium, Max (if exceeding salt guarantee)..... %
 Vitamin A, Min..... IU/lb

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to manufacture a complete feed. The following table provides examples of mixing and feeding rates:

Type B Medicated Feed Concentration (g/ton)	Type B Medicated Feed to make one ton of Type C Medicated Feed (lb)	Drug Level in Type C Medicated Feed (g/ton)	Type C Medicated Feed Consumption (lbs. of feed per 100 lb body weight per day)
10000	400	2000	1
10000	200	1000	2
8000	250	1000	2
10000	133.3	666.7	3
8000	166.7	666.7	3

The resulting Type C medicated feed should be fed at a rate to provide 10 mg/lb body weight per day. Treat for not more than 5 days.

Withdrawal Periods and Residue Warnings: Withdraw 10 days prior to slaughter. A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

BLUE BIRD FEED COMPANY
CITY. STATE. ZIP

NET WEIGHT ON BAG OR BULK

* This Bluebird label refers to Chloratet™ brand of Chlortetracycline.

**Final label for Type B medicated feed must state a single concentration of drug which will vary with body weight and feed consumption to provide 10 mg/lb body weight per day in the Type C feed.

Lot No.